

Research Article

The Dirichlet Problem for Second-Order Divergence Form Elliptic Operators with Variable Coefficients: The Simple Layer Potential Ansatz

Alberto Cialdea, Vita Leonessa, and Angelica Malaspina

Department of Mathematics, Computer Science and Economics, University of Basilicata, Viale dell'Ateneo Lucano 10, 85100 Potenza, Italy

Correspondence should be addressed to Alberto Cialdea; cialdea@email.it

Received 9 June 2015; Accepted 18 October 2015

Academic Editor: Giovanni P. Galdi

Copyright © 2015 Alberto Cialdea et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate the Dirichlet problem related to linear elliptic second-order partial differential operators with smooth coefficients in divergence form in bounded connected domains of \mathbb{R}^m ($m \geq 3$) with Lyapunov boundary. In particular, we show how to represent the solution in terms of a simple layer potential. We use an indirect boundary integral method hinging on the theory of reducible operators and the theory of differential forms.

1. Introduction

As remarked in [1, p. 121], elliptic operators with variable coefficients naturally arise in several areas of physics and engineering. In this paper, we study the Dirichlet problem related to a scalar elliptic second-order differential operator with smooth coefficients in divergence form in a bounded simply connected domain of \mathbb{R}^m ($m \geq 3$) with Lyapunov boundary.

This is a classical problem which nowadays can be treated in several ways. In particular, different potential methods have been developed for such operators (see, e.g., [1–6]).

In the present paper, we obtain the solution of the Dirichlet problem by means of a simple layer potential instead of the classical double layer potential (see, e.g., [6, pp. 73–75]). We use an indirect boundary integral method introduced for the first time in [7] for the m -dimensional Laplacian. It requires neither the knowledge of pseudodifferential operators nor the use of hypersingular integrals, but it hinges on the theory of singular integral operators and the theory of differential forms (for details of the method, see, e.g., [8, Section 2]). The method has been also used to treat different boundary value problems in simply connected domains: the Neumann problem for Laplace equation (via a double layer potential), the Dirichlet problem for the Lamé and Stokes systems, the four

boundary value problems of the theory of thermoelastic pseudooscillations, the traction problem for Lamé and Stokes systems, the four basic boundary value problems arising in couple-stress elasticity, and the two boundary value problems of the linear theory of viscoelastic materials with voids (see [9, 10] and the references therein). The method can be applied also in multiply connected domains, as shown for the Laplacian, the linearized elastostatics, and the Stokes system (see [11] and the references therein).

The present paper is organized as follows.

In Section 2, after giving preliminary results, we make use of Fichera's construction of a principal fundamental solution [12] and we prove some identities for the related nuclear double form.

Section 3 is devoted to the study of the Dirichlet problem. It contains the main results concerning the reduction of a certain singular integral operator acting in spaces of differential forms and the integral representation of the solution of the Dirichlet problem by means of a simple layer potential.

2. Preliminary Results

Let Ω be a bounded domain (open connected set) of \mathbb{R}^m ($m \geq 3$).

In this paper, we deal with the Dirichlet problem:

$$\begin{aligned} Eu &= 0 \quad \text{in } \Omega, \\ u &= f \quad \text{on } \Sigma, \end{aligned} \tag{1}$$

where E is a scalar second-order differential operator (throughout this paper, we use the Einstein summation convention):

$$Eu(x) = \frac{\partial}{\partial x^i} \left(a^{ij}(x) \frac{\partial u(x)}{\partial x^j} \right). \tag{2}$$

We suppose that the coefficients a^{ij} are defined on \bar{T} , T being an open ball containing $\bar{\Omega}$, and we assume that they belong to $C^{2,\lambda}(\bar{T})$, $0 < \lambda \leq 1$.

Moreover, assume that $A = (a^{ij})_{i,j=1,\dots,m}$ is a symmetric contravariant positive-definite tensor. Then, E is a uniform elliptic operator; that is, there exists $c > 0$ such that $a^{ij}(x)\xi_i\xi_j \geq c|\xi|^2$, for every $(\xi_1, \dots, \xi_m) \in \mathbb{R}^m$ and for any $x \in T$.

For the sake of simplicity, we suppose that the determinant $|A|$ of A is equal to 1.

It is known that to the contravariant tensor A there corresponds a covariant tensor $A^{-1} = (a_{ij})_{i,j=1,\dots,m}$ such that

$$a^{ij}a_{jh} = \delta_h^i, \quad \text{for every } i, h = 1, \dots, m, \tag{3}$$

δ_h^i being the Kronecker delta.

A differential form of degree k (in short a k -form) on T is a function defined on T whose values are in the k -covectors space of \mathbb{R}^m . A k -form u can be represented as

$$u = \frac{1}{k!} u_{s_1 \dots s_k} dx^{s_1} \dots dx^{s_k} \tag{4}$$

with respect to an admissible coordinate system (x_1, \dots, x_m) , where $u_{s_1 \dots s_k}$ are the components of a skew-symmetric covariant tensor (for details about differential forms, we refer to [13, 14]).

The symbol $C_k^h(T)$ means the space of all k -forms whose components are continuously differentiable up to the order h in a coordinate system of class C^{h+1} (and then in every coordinate system of class C^{h+1}).

If $u \in C_k^1(T)$, the differential of u is a $(k+1)$ -form defined as

$$du = \frac{1}{k!} \frac{\partial u_{s_1 \dots s_k}}{\partial x^j} dx^j dx^{s_1} \dots dx^{s_k}. \tag{5}$$

Further, if $u \in C_k^0(T)$, the adjoint of u is the following $(m-k)$ -form:

$$\begin{aligned} *u &= \frac{1}{k!(m-k)!} \delta_{j_1 \dots j_k i_{k+1} \dots i_m}^{1 \dots m} \\ &\cdot a^{s_1 j_1} \dots a^{s_k j_k} u_{s_1 \dots s_k} dx^{i_{k+1}} \dots dx^{i_m}, \end{aligned} \tag{6}$$

where $\delta_{q_1 \dots q_r}^{p_1 \dots p_r}$ is the generalized Kronecker delta ($r \leq m$). We recall that (see, e.g., [15, p. 127])

$$\delta_{h_1 \dots h_s h_{s+1} \dots h_m}^{j_1 \dots j_s j_{s+1} \dots j_m} \delta_{k_{s+1} \dots k_m}^{h_{s+1} \dots h_m} = (m-s)! \delta_{h_1 \dots h_s k_{s+1} \dots k_m}^{j_1 \dots j_s j_{s+1} \dots j_m}. \tag{7}$$

We remark that (see, e.g., [13, p. 285])

$$**u = (-1)^{k(m+1)} u. \tag{8}$$

If $u \in C_k^1(T)$, we define the codifferential of u as the following $(k-1)$ -form:

$$\delta u = (-1)^{m(k+1)+1} * d * u. \tag{9}$$

A differential double form $u_{h,k}(x, y)$ of degree h with respect to x and of degree k with respect to y (in short a double (h, k) -form) is represented as

$$\begin{aligned} u_{h,k}(x, y) &= \frac{1}{h!k!} u_{s_1 \dots s_h j_1 \dots j_k}(x, y) dx^{s_1} \dots dx^{s_h} dy^{j_1} \dots dy^{j_k}. \end{aligned} \tag{10}$$

If $h = k$, we denote it briefly by $u_k(x, y)$.

We proceed to introduce the following double k -form (see [13, p. 204]) defined, for every $x, y \in \bar{T}$, $x \neq y$, as

$$\begin{aligned} \lambda_k(x, y) &= \frac{1}{(k!)^2} L(x, y) \\ &\cdot a_{s_1 \dots s_k i_1 \dots i_k}(y) dx^{s_1} \dots dx^{s_k} dy^{i_1} \dots dy^{i_k}, \end{aligned} \tag{11}$$

where, for $k \leq m$,

$$a_{s_1 \dots s_k i_1 \dots i_k} = \begin{vmatrix} a_{s_1 i_1} & \dots & a_{s_1 i_k} \\ \vdots & \ddots & \vdots \\ a_{s_k i_1} & \dots & a_{s_k i_k} \end{vmatrix} = \delta_{i_1 \dots i_k}^{s_1 \dots s_k} a_{s_1 i_1} \dots a_{s_k i_k}, \tag{12}$$

$$\begin{aligned} L(x, y) &= \frac{1}{(m-2)\omega_m} [a_{ij}(y)(x^i - y^i)(x^j - y^j)]^{(2-m)/2} \end{aligned} \tag{13}$$

(ω_m being the hypersurface measure of the unit sphere in \mathbb{R}^m) is a parametrix in the sense of Hilbert and E.E. Levi for the operator E . We recall that (if we write $u_{h,k}(x, y) = \mathcal{O}(|x - y|^\alpha)$, $u_{h,k}(x, y)$ being a double (h, k) -form, we mean that all its components are $\mathcal{O}(|x - y|^\alpha)$)

$$L(x, y) = \mathcal{O}(|x - y|^{2-m}), \tag{14}$$

$$d_x L(x, y) = \mathcal{O}(|x - y|^{1-m}), \tag{15}$$

$$d_y L(x, y) = \mathcal{O}(|x - y|^{1-m})$$

(see [13, Section 9]).

The next results provide other properties of L and λ_k .

Lemma 1. For every $p = 1, \dots, m$,

$$\frac{\partial L(x, y)}{\partial y^p} = -\frac{\partial L(x, y)}{\partial x^p} + M(x, y), \tag{16}$$

$$x, y \in \bar{T}, \quad x \neq y,$$

where $M(x, y) = \mathcal{O}(|x - y|^{2-m})$.

Proof. Taking definition (13) into account, we have

$$\begin{aligned} \frac{\partial L(x, y)}{\partial x^p} &= -\frac{1}{2\omega_m} [a_{ij}(y)(x^i - y^i)(x^j - y^j)]^{-m/2} \\ &\cdot \{a_{ij}(y) [\delta_p^i(x^j - y^j) + \delta_p^j(x^i - y^i)]\} \\ &= -\frac{1}{\omega_m} [a_{ij}(y)(x^i - y^i)(x^j - y^j)]^{-m/2} \\ &\cdot [a_{pj}(y)(x^j - y^j)]. \end{aligned} \tag{17}$$

On the other hand,

$$\begin{aligned} \frac{\partial L(x, y)}{\partial y^p} &= -\frac{1}{2\omega_m} [a_{ij}(y)(x^i - y^i)(x^j - y^j)]^{-m/2} \\ &\cdot \left[\frac{\partial a_{ij}(y)}{\partial y^p} (x^i - y^i)(x^j - y^j) \right] \\ &- \frac{1}{2\omega_m} [a_{ij}(y)(x^i - y^i)(x^j - y^j)]^{-m/2} \\ &\cdot \{a_{ij}(y) [-\delta_p^i(x^j - y^j) - \delta_p^j(x^i - y^i)]\} \\ &= M(x, y) + \frac{1}{\omega_m} [a_{ij}(y)(x^i - y^i)(x^j - y^j)]^{-m/2} \\ &\cdot [a_{pj}(y)(x^j - y^j)] = M(x, y) - \frac{\partial L(x, y)}{\partial x^p} \end{aligned} \tag{18}$$

and this yields the claim. \square

The identities proved in the next proposition generalize the ones obtained by Colautti [16, p. 309] for the Laplacian.

Proposition 2. *Let λ_k be the double k -form defined by (11). Then, for every $x \neq y$, the following properties hold:*

$$\begin{aligned} *_x \lambda_k(x, y) &= (-1)^{k(m-k)} *_y \lambda_{m-k}(x, y) \\ &+ \tau_{m-k, k}(x, y), \quad k \leq m, \end{aligned} \tag{19}$$

where

$$\tau_{m-k, k}(x, y) = \mathcal{O}(|x - y|^{3-m}), \tag{20}$$

$$\begin{aligned} *_x d_x \lambda_k(x, y) &= (-1)^{mk+1} *_y d_y \lambda_{m-k-1}(x, y) \\ &+ \gamma_{m-k-1, k}(x, y), \quad k < m, \end{aligned} \tag{21}$$

where $\gamma_{m-k-1, k}(x, y) = \mathcal{O}(|x - y|^{2-m})$; and

$$\delta_x \lambda_{k+1}(x, y) = d_y \lambda_k(x, y) + \epsilon_{k, k+1}(x, y), \quad k < m, \tag{22}$$

where

$$\epsilon_{k, k+1}(x, y) = \mathcal{O}(|x - y|^{2-m}). \tag{23}$$

Proof. First, we prove (19). It follows from (12), (3), and (7) that

$$\begin{aligned} *_y \lambda_{m-k}(x, y) &= \frac{(-1)^{k(m-k)}}{k!(m-k)!} \\ &\cdot \delta_{p_1 \dots p_k q_1 \dots q_{m-k}}^{1 \dots m} L(x, y) dx^{q_1} \dots dx^{q_{m-k}} dy^{p_1} \dots dy^{p_k}. \end{aligned} \tag{24}$$

On the other hand,

$$\begin{aligned} *_x \lambda_k(x, y) &= \frac{1}{(m-k)!(k!)^2} \\ &\cdot \delta_{j_1 \dots j_k q_1 \dots q_{m-k}}^{1 \dots m} [(a^{s_1 j_1} \dots a^{s_k j_k})(x) \\ &- (a^{s_1 j_1} \dots a^{s_k j_k})(y)] \\ &\cdot L(x, y) a_{s_1 \dots s_k p_1 \dots p_k}(y) dx^{q_1} \dots dx^{q_{m-k}} dy^{p_1} \dots dy^{p_k} \\ &+ \frac{1}{(m-k)!(k!)^2} \delta_{j_1 \dots j_k q_1 \dots q_{m-k}}^{1 \dots m} (a^{s_1 j_1} \dots a^{s_k j_k})(y) \\ &\cdot L(x, y) a_{s_1 \dots s_k p_1 \dots p_k}(y) dx^{q_1} \dots dx^{q_{m-k}} dy^{p_1} \dots dy^{p_k}. \end{aligned} \tag{25}$$

From (12), (3), and (7), we have that

$$\begin{aligned} *_x \lambda_k(x, y) &= \frac{1}{(m-k)!(k!)^2} \\ &\cdot \delta_{j_1 \dots j_k q_1 \dots q_{m-k}}^{1 \dots m} [(a^{s_1 j_1} \dots a^{s_k j_k})(x) \\ &- (a^{s_1 j_1} \dots a^{s_k j_k})(y)] L(x, y) \\ &\cdot a_{s_1 \dots s_k p_1 \dots p_k}(y) dx^{q_1} \dots dx^{q_{m-k}} dy^{p_1} \dots dy^{p_k} \\ &+ \frac{1}{(m-k)!k!} \\ &\cdot \delta_{p_1 \dots p_k q_1 \dots q_{m-k}}^{1 \dots m} L(x, y) dx^{q_1} \dots dx^{q_{m-k}} dy^{p_1} \dots dy^{p_k} \\ &= \tau_{m-k, k}(x, y) + (-1)^{k(m-k)} *_y \lambda_{m-k}(x, y), \end{aligned} \tag{26}$$

where $\tau_{m-k, k}(x, y)$ satisfies (20) on account of

$$(a^{s_1 j_1} \dots a^{s_k j_k})(x) - (a^{s_1 j_1} \dots a^{s_k j_k})(y) = \mathcal{O}(|x - y|) \tag{27}$$

and (8).

Now we pass to show (21). With calculations analogue to (26), we have that

$$\begin{aligned}
*_x d_x \lambda_k(x, y) &= \frac{1}{(m-k-1)!(k!)^2} \delta_{j_1 \dots j_k i_{k+2} \dots i_m}^{1 \dots m} \\
&\cdot \left[(a^{s_j} a^{s_1 j_1} \dots a^{s_k j_k})(x) \right. \\
&- (a^{s_j} a^{s_1 j_1} \dots a^{s_k j_k})(y) \left. \right] \frac{\partial L(x, y)}{\partial x^s} a_{s_1 \dots s_k i_1 \dots i_k}(y) \\
&\cdot dx^{i_{k+2}} \dots dx^{i_m} dy^{i_1} \dots dy^{i_k} + \frac{1}{(m-k-1)!(k!)^2} \\
&\cdot \delta_{j_1 \dots j_k i_{k+2} \dots i_m}^{1 \dots m} (a^{s_j} a^{s_1 j_1} \dots a^{s_k j_k})(y) \frac{\partial L(x, y)}{\partial x^s} \\
&\cdot a_{s_1 \dots s_k i_1 \dots i_k}(y) dx^{i_{k+2}} \dots dx^{i_m} dy^{i_1} \dots dy^{i_k} \\
&= \gamma'_{m-k-1, k}(x, y) + \frac{1}{(m-k-1)!k!} \\
&\cdot \delta_{j_{k+2} \dots i_m i_1 \dots i_k}^{1 \dots m} a^{s_j}(y) \cdot \frac{\partial L(x, y)}{\partial x^s} \\
&\cdot dx^{i_{k+2}} \dots dx^{i_m} dy^{i_1} \dots dy^{i_k},
\end{aligned} \tag{28}$$

where $\gamma'_{m-k-1, k}(x, y) = \mathcal{O}(|x-y|^{1-m})$ thanks to (15) and (27). Moreover,

$$\begin{aligned}
*_y d_y \lambda_{m-k-1}(x, y) &= \frac{1}{k! [(m-k-1)!]^2} \delta_{j_{q_1} \dots q_{m-k-1} i_1 \dots i_k}^{1 \dots m} \\
&\cdot (a^{s_j} a^{p_1 q_1} \dots a^{p_{m-k-1} q_{m-k-1}})(y) \\
&\cdot \frac{\partial}{\partial y^s} [L(x, y) a_{s_1 \dots s_{m-k-1} p_1 \dots p_{m-k-1}}(y)] \\
&\cdot dx^{s_1} \dots dx^{s_{m-k-1}} dy^{i_1} \dots dy^{i_k}.
\end{aligned} \tag{29}$$

Arguing again as in (26) and taking Lemma 1 into account, we get

$$\begin{aligned}
*_y d_y \lambda_{m-k-1}(x, y) &= \frac{(-1)^{k(m-k-1)}}{k!(m-k-1)!} \\
&\cdot \delta_{j_1 \dots i_k s_1 \dots s_{m-k-1}}^{1 \dots m} a^{s_j}(y) \frac{\partial L(x, y)}{\partial y^s} \\
&\cdot dx^{s_1} \dots dx^{s_{m-k-1}} dy^{i_1} \dots dy^{i_k} + \frac{1}{k! [(m-k-1)!]^2} \\
&\cdot \delta_{j_{q_1} \dots q_{m-k-1} i_1 \dots i_k}^{1 \dots m} (a^{s_j} a^{p_1 q_1} \dots a^{p_{m-k-1} q_{m-k-1}})(y) \\
&\cdot L(x, y) \frac{\partial}{\partial y^s} a_{s_1 \dots s_{m-k-1} p_1 \dots p_{m-k-1}}(y) \\
&\cdot dx^{s_1} \dots dx^{s_{m-k-1}} dy^{i_1} \dots dy^{i_k} = -\frac{(-1)^{mk}}{k!(m-k-1)!} \\
&\cdot \delta_{j_1 \dots i_k s_1 \dots s_{m-k-1}}^{1 \dots m} a^{s_j}(y)
\end{aligned}$$

$$\begin{aligned}
&\cdot \frac{\partial L(x, y)}{\partial x^s} dx^{s_1} \dots dx^{s_{m-k-1}} dy^{i_1} \dots dy^{i_k} \\
&+ \frac{(-1)^{mk}}{k!(m-k-1)!} \delta_{j_1 \dots i_k s_1 \dots s_{m-k-1}}^{1 \dots m} a^{s_j}(y) \\
&\cdot M(x, y) dx^{s_1} \dots dx^{s_{m-k-1}} dy^{i_1} \dots dy^{i_k} \\
&+ \gamma''_{m-k-1, k}(x, y) = (-1)^{mk+1} *_x d_x \lambda_k(x, y) \\
&+ \gamma'''_{m-k-1, k}(x, y) + \gamma''''_{m-k-1, k}(x, y),
\end{aligned} \tag{30}$$

where both $\gamma''_{m-k-1, k}(x, y)$ and $\gamma'''_{m-k-1, k}(x, y)$ are $\mathcal{O}(|x-y|^{2-m})$. Then, we obtain the claim by setting

$$\begin{aligned}
\gamma_{m-k-1, k}(x, y) \\
&= (-1)^{mk} (\gamma''_{m-k-1, k}(x, y) + \gamma'''_{m-k-1, k}(x, y)).
\end{aligned} \tag{31}$$

Finally, we prove (22). Thanks to (9) and (19), we have

$$\begin{aligned}
\delta_x \lambda_{k+1}(x, y) &= (-1)^{m-k} *_x d_x *_y \lambda_{m-k-1}(x, y) \\
&+ (-1)^{mk+1} *_x d_x \tau_{m-k-1, k+1}(x, y) \\
&= (-1)^{m-k} *_y *_x d_x \lambda_{m-k-1}(x, y) \\
&+ \epsilon'_{k, k+1}(x, y),
\end{aligned} \tag{32}$$

where $\epsilon'_{k, k+1}(x, y) = \mathcal{O}(|x-y|^{2-m})$. Now, by using (21) and (8), we get

$$\begin{aligned}
\delta_x \lambda_{k+1}(x, y) &= (-1)^{m-k-km+1} *_y *_y d_y \lambda_k(x, y) \\
&+ (-1)^{m-k} *_y \gamma_{k, m-k-1}(x, y) \\
&+ \epsilon'_{k, k+1}(x, y) \\
&= d_y \lambda_k(x, y) + \epsilon_{k, k+1}(x, y),
\end{aligned} \tag{33}$$

and hence the claim with

$$\begin{aligned}
\epsilon_{k, k+1}(x, y) &= (-1)^{m-k} *_y \gamma_{k, m-k-1}(x, y) \\
&+ \epsilon'_{k, k+1}(x, y).
\end{aligned} \tag{34}$$

□

Proposition 3. If $u \in C_k^2(T)$, then

$$(\delta d + d\delta)u = -Eu + Fu, \tag{35}$$

where F is a linear first-order differential operator whose coefficients depend only on first- and second-order derivatives of entries of the tensor A .

In particular,

$$(\delta_x d_x + d_x \delta_x) \lambda_k(x, y) = F_x[\lambda_k(x, y)], \quad x \neq y. \quad (36)$$

Proof. We begin by observing that

$$\begin{aligned} d\delta u &= (-1)^{m(k+1)+1} d \frac{1}{(k-1)!k!(m-k)!} \delta_{hh_{k+1} \dots h_m i_2 \dots i_k}^{1 \dots m} a^{jh} a^{i_{k+1} h_{k+1}} \dots a^{i_m h_m} \delta_{j_1 \dots j_k i_{k+1} \dots i_m}^{1 \dots m} a^{s_1 j_1} \dots a^{s_k j_k} \frac{\partial}{\partial x^j} u_{s_1 \dots s_k} dx^{i_2} \dots dx^{i_k} \\ &+ (-1)^{m(k+1)+1} d \frac{1}{(k-1)!k!(m-k)!} \delta_{hh_{k+1} \dots h_m i_2 \dots i_k}^{1 \dots m} a^{jh} a^{i_{k+1} h_{k+1}} \dots a^{i_m h_m} \cdot \delta_{j_1 \dots j_k i_{k+1} \dots i_m}^{1 \dots m} \frac{\partial}{\partial x^j} (a^{s_1 j_1} \dots a^{s_k j_k}) \\ &\cdot u_{s_1 \dots s_k} dx^{i_2} \dots dx^{i_k}. \end{aligned} \quad (37)$$

Since A is symmetric and $|A| = 1$, we get

$$\begin{aligned} &\delta_{j_1 \dots j_k i_{k+1} \dots i_m}^{1 \dots m} a^{s_1 j_1} \dots a^{s_k j_k} a^{i_{k+1} h_{k+1}} \dots a^{i_m h_m} \\ &= \delta_{1 \dots m}^{s_1 \dots s_k h_{k+1} \dots h_m} \end{aligned} \quad (38)$$

$$\begin{aligned} &\cdot \frac{\partial}{\partial x^{i_1}} \left[a^{jh} a^{i_{k+1} h_{k+1}} \dots a^{i_m h_m} \frac{\partial}{\partial x^j} (a^{s_1 j_1} \dots a^{s_k j_k}) \right. \\ &\cdot \left. u_{s_1 \dots s_k} \right] dx^{i_1} dx^{i_2} \dots dx^{i_k}. \end{aligned} \quad (39)$$

and, keeping in mind (7), we get

$$\begin{aligned} d\delta u &= -\frac{1}{(k-1)!k!} \delta_{hi_2 \dots i_k}^{s_1 \dots s_k} \frac{\partial}{\partial x^{i_1}} \left(a^{jh} \right. \\ &\cdot \left. \frac{\partial u_{s_1 \dots s_k}}{\partial x^j} \right) dx^{i_1} dx^{i_2} \dots dx^{i_k} \\ &+ (-1)^{m(k+1)+1} \frac{1}{(k-1)!k!(m-k)!} \\ &\cdot \delta_{hh_{k+1} \dots h_m i_2 \dots i_k}^{1 \dots m} \delta_{j_1 \dots j_k i_{k+1} \dots i_m}^{1 \dots m} \end{aligned}$$

On the other hand,

$$\begin{aligned} \delta du &= (-1)^{m(k+2)+1} \frac{1}{(m-k-1)!(k!)^2} \\ &\cdot \delta_{qj_{k+2} \dots j_m i_1 \dots i_k}^{1 \dots m} a^{pq} a^{i_{k+2} j_{k+2}} \dots a^{i_m j_m} \delta_{hh_1 \dots h_k i_{k+2} \dots i_m}^{1 \dots m} \\ &\cdot \frac{\partial}{\partial x^p} \left(a^{jh} a^{s_1 h_1} \dots a^{s_k h_k} \frac{\partial u_{s_1 \dots s_k}}{\partial x^j} \right) dx^{i_1} \dots dx^{i_k}. \end{aligned} \quad (40)$$

Then,

$$\begin{aligned} d\delta u + \delta du &= -\frac{1}{(k-1)!k!} \delta_{hi_2 \dots i_k}^{s_1 \dots s_k} a^{jh} \frac{\partial^2 u_{s_1 \dots s_k}}{\partial x^{i_1} \partial x^j} dx^{i_1} dx^{i_2} \dots dx^{i_k} + (-1)^{m(k+2)+1} \frac{1}{(m-k-1)!(k!)^2} \delta_{qj_{k+2} \dots j_m i_1 \dots i_k}^{1 \dots m} \\ &\cdot a^{pq} a^{i_{k+2} j_{k+2}} \dots a^{i_m j_m} \cdot \delta_{hh_1 \dots h_k i_{k+2} \dots i_m}^{1 \dots m} a^{jh} a^{s_1 h_1} \dots a^{s_k h_k} \frac{\partial^2 u_{s_1 \dots s_k}}{\partial x^p \partial x^j} dx^{i_1} \dots dx^{i_k} - \frac{1}{(k-1)!k!} \delta_{hi_2 \dots i_k}^{s_1 \dots s_k} \frac{\partial a^{jh}}{\partial x^{i_1}} \frac{\partial u_{s_1 \dots s_k}}{\partial x^j} \\ &\cdot dx^{i_1} dx^{i_2} \dots dx^{i_k} + (-1)^{m(k+1)+1} \frac{1}{(k-1)!k!(m-k)!} \delta_{hh_{k+1} \dots h_m i_2 \dots i_k}^{1 \dots m} \delta_{j_1 \dots j_k i_{k+1} \dots i_m}^{1 \dots m} a^{jh} a^{i_{k+1} h_{k+1}} \dots a^{i_m h_m} \\ &\cdot \frac{\partial}{\partial x^j} (a^{s_1 j_1} \dots a^{s_k j_k}) \frac{\partial u_{s_1 \dots s_k}}{\partial x^{i_1}} dx^{i_1} dx^{i_2} \dots dx^{i_k} + (-1)^{m(k+2)+1} \frac{1}{(m-k-1)!(k!)^2} \\ &\cdot \delta_{qj_{k+2} \dots j_m i_1 \dots i_k}^{1 \dots m} a^{pq} a^{i_{k+2} j_{k+2}} \dots a^{i_m j_m} \delta_{hh_1 \dots h_k i_{k+2} \dots i_m}^{1 \dots m} \frac{\partial}{\partial x^p} (a^{jh} a^{s_1 h_1} \dots a^{s_k h_k}) \frac{\partial u_{s_1 \dots s_k}}{\partial x^j} dx^{i_1} \dots dx^{i_k} + (-1)^{m(k+1)+1} \\ &\cdot \frac{1}{(k-1)!k!(m-k)!} \delta_{hh_{k+1} \dots h_m i_2 \dots i_k}^{1 \dots m} \delta_{j_1 \dots j_k i_{k+1} \dots i_m}^{1 \dots m} \frac{\partial}{\partial x^{i_1}} (a^{jh} a^{i_{k+1} h_{k+1}} \dots a^{i_m h_m}) \frac{\partial}{\partial x^j} (a^{s_1 j_1} \dots a^{s_k j_k}) u_{s_1 \dots s_k} dx^{i_1} dx^{i_2} \dots dx^{i_k} \\ &+ (-1)^{m(k+1)+1} \frac{1}{(k-1)!k!(m-k)!} \delta_{hh_{k+1} \dots h_m i_2 \dots i_k}^{1 \dots m} \delta_{j_1 \dots j_k i_{k+1} \dots i_m}^{1 \dots m} a^{jh} a^{i_{k+1} h_{k+1}} \dots a^{i_m h_m} \frac{\partial^2}{\partial x^{i_1} \partial x^j} (a^{s_1 j_1} \dots a^{s_k j_k}) \\ &\cdot u_{s_1 \dots s_k} dx^{i_1} dx^{i_2} \dots dx^{i_k} = -\frac{1}{k!} \frac{\partial}{\partial x^p} \left(a^{jp} \frac{\partial u_{s_1 \dots s_k}}{\partial x^j} \right) dx^{s_1} \dots dx^{s_k} + \frac{m! - (m-k)!}{m!k!} \frac{\partial a^{jp}}{\partial x^p} \frac{\partial u_{s_1 \dots s_k}}{\partial x^j} dx^{s_1} \dots dx^{s_k} \\ &- \frac{m! - (m-k)!}{m!(k-1)!k!} \delta_{hi_2 \dots i_k}^{s_1 s_2 \dots s_k} \frac{\partial a^{jh}}{\partial x^{i_1}} \frac{\partial u_{s_1 \dots s_k}}{\partial x^j} dx^{i_1} \dots dx^{i_k} - \frac{(m-k+1)!}{m!(k-1)!^2} \delta_{hi_2 \dots i_k}^{j s_2 \dots s_k} \frac{\partial a^{s_1 h}}{\partial x^j} \frac{\partial u_{s_1 \dots s_k}}{\partial x^{i_1}} dx^{i_1} \dots dx^{i_k} + \frac{(m-k)!}{m!(k-1)!k!} \delta_{i_1 i_2 \dots i_k}^{j s_2 \dots s_k} \end{aligned}$$

$$\begin{aligned} & \cdot \frac{\partial a^{s_1, h}}{\partial x^h} \frac{\partial u_{s_1, \dots, s_k}}{\partial x^j} dx^{i_1} \dots dx^{i_k} - \frac{(m-k)!}{m!(k-1)!^2} \delta_{hi_2, \dots, i_k}^{js_2, \dots, s_k} \frac{\partial a^{s_1, h}}{\partial x^{i_1}} \frac{\partial u_{s_1, \dots, s_k}}{\partial x^j} dx^{i_1} \dots dx^{i_k} - \frac{(m-k+1)!}{m!(k-1)!} \delta_{hi_2, \dots, i_k}^{js_2, \dots, s_k} \frac{\partial^2 a^{s_1, h}}{\partial x^{i_1} \partial x^j} u_{s_1, \dots, s_k} dx^{i_1} \dots dx^{i_k} \\ & = -\frac{1}{k!} E u_{s_1, \dots, s_k} dx^{s_1} \dots dx^{s_k} + \frac{1}{k!} F u_{s_1, \dots, s_k} dx^{s_1} \dots dx^{s_k} \end{aligned} \tag{41}$$

and this proves (35). Finally, (36) follows from (35). \square

Finally, following Fichera we employ the parametrix L to construct a principal fundamental solution of the differential operator E (see [12]).

Lemma 4. *There exists $\zeta(x, y)$ such that the function*

$$S(x, y) = L(x, y) + \zeta(x, y), \quad x \in \bar{T}, \quad y \in T, \tag{42}$$

is a principal fundamental solution of E . In particular, we have

$$\begin{aligned} E_x S(x, y) &= 0, \quad x \in T, \quad y \in T, \quad x \neq y, \\ S(x, y) &= 0, \quad x \in \partial T, \quad y \in T, \\ S(x, y) &= \mathcal{O}(|x - y|^{2-m}). \end{aligned} \tag{43}$$

Moreover, for every $x \neq y$,

$$d_x \zeta(x, y) = \mathcal{O}(|x - y|^{2-m-\gamma}) \tag{44}$$

for some $0 < \gamma \leq 1$.

Proof. The existence of $\zeta(x, y)$ can be obtained as the solution of a certain integral equation (see [12, §2]). In [12] properties (43) are proved and $\zeta(x, y)$ is written as

$$\zeta(x, y) = G(x, y) + \int_T L(x, w) R(w, y) dw, \tag{45}$$

where G is a smooth function on T and, for some $0 < \gamma \leq 1$,

$$R(w, y) = \mathcal{O}(|w - y|^{1-m-\gamma}). \tag{46}$$

Then, by (15),

$$\begin{aligned} d_x \zeta(x, y) &= d_x G(x, y) + \int_T d_x L(x, w) R(w, y) dw \\ &= \mathcal{O}(|x - y|^{2-m-\gamma}). \end{aligned} \tag{47}$$

\square

3. The Dirichlet Problem

In this section, we suppose that the domain Ω is such that $\mathbb{R}^m \setminus \bar{\Omega}$ is connected and such that its boundary $\Sigma = \partial\Omega$ is a Lyapunov surface (i.e., $\Sigma \in C^{1,\lambda}$, $0 < \lambda \leq 1$).

By $n(y) = (n_1(y), \dots, n_m(y))$, we denote the outwards unit normal vector at the point $y \in \Sigma$ and by $\nu(y) = (\nu_1(y), \dots, \nu_m(y))$ we denote the conormal vector at the point

$y \in \Sigma$ associated with the operator E and defined as $\nu_i(y) = a^{ij}(y)n_j(y)$ ($i = 1, \dots, m$). By $\partial u / \partial \nu$ we denote the conormal derivative

$$\frac{\partial u}{\partial \nu} = a^{ij} n_j \frac{\partial u}{\partial y^i}. \tag{48}$$

As usual, the symbols $L^p(\Sigma)$ and $W^{1,p}(\Sigma)$ ($1 < p < +\infty$) stand for the classical Lebesgue and Sobolev spaces, respectively.

By $L^p_k(\Sigma)$, we denote the space of all k -forms whose components are L^p real-valued functions in a coordinate system of class C^1 (and then in every coordinate system of class C^1).

We will look for the solution of the Dirichlet problem for the operator E in the domain Ω in the form of a simple layer potential. To this end, we introduce the space \mathcal{S}^p .

Definition 5. The function u belongs to \mathcal{S}^p if and only if there exists $\varphi \in L^p(\Sigma)$ such that it can be represented by means of a simple layer potential; that is,

$$u(x) = \int_{\Sigma} \varphi(y) S(x, y) d\sigma_y, \quad x \in \Omega. \tag{49}$$

Specifically our aim is to give an existence and uniqueness theorem for the Dirichlet problem

$$\begin{aligned} u &\in \mathcal{S}^p, \\ Eu &= 0 \quad \text{in } \Omega, \\ u &= f \quad \text{on } \Sigma, \quad f \in W^{1,p}(\Sigma). \end{aligned} \tag{50}$$

First, we prove the following formula.

Proposition 6. *For any $u \in W^{1,p}(\Sigma)$,*

$$\begin{aligned} & \frac{\partial}{\partial \nu_z} \left(\int_{\Sigma} u(x) \frac{\partial}{\partial \nu_x} L(z, x) d\sigma_x \right) d\sigma_z \\ &= d_z \int_{\Sigma} du(x) \wedge \lambda_{m-2}(z, x) \\ &+ \int_{\Sigma} u(x) \wedge F_z [\lambda_{m-1}(z, x)] \\ &- \int_{\Sigma} u(x) \wedge \eta_{m-1}(z, x), \quad z \in \Sigma, \end{aligned} \tag{51}$$

where F is the linear first-order differential operator considered in Proposition 3 and

$$\eta_{m-1}(z, x) = \mathcal{O}(|z - x|^{1-m}). \tag{52}$$

Proof. Set, for every $z \notin \Sigma$,

$$\begin{aligned} U(z) &= \int_{\Sigma} du(x) \wedge \lambda_{m-2}(z, x), \\ V(z) &= \int_{\Sigma} u(x) \wedge d_z [\lambda_{m-1}(z, x)]. \end{aligned} \tag{53}$$

On account of (22) and (36), we get

$$\begin{aligned} dU(z) &= - \int_{\Sigma} u(x) \wedge d_z d_x [\lambda_{m-2}(z, x)] \\ &= - \int_{\Sigma} u(x) \wedge d_z \delta_z [\lambda_{m-1}(z, x)] \\ &\quad + \int_{\Sigma} u(x) \wedge d_z [\epsilon_{m-2, m-1}(z, x)] \\ &= \int_{\Sigma} u(x) \wedge \delta_z d_z [\lambda_{m-1}(z, x)] \\ &\quad - \int_{\Sigma} u(x) \wedge F_z [\lambda_{m-1}(z, x)] \\ &\quad + \int_{\Sigma} u(x) \wedge d_z [\epsilon_{m-2, m-1}(z, x)] \\ &= \delta V(z) - \int_{\Sigma} u(x) \wedge F_z [\lambda_{m-1}(z, x)] \\ &\quad + \int_{\Sigma} u(x) \wedge \eta_{m-1}(z, x) \end{aligned} \tag{54}$$

and (52) follows from (23).

On the other hand, if A_i^j is the minor of A^{-1} obtained deleting the i th row and the j th column, for $z \in \Omega$, $x \in \Sigma$ we get

$$\begin{aligned} d_z [\lambda_{m-1}(z, x)] &= d_z L(z, x) \\ &\cdot |A_i^j(x)| dz^1 \cdots \widehat{i} \cdots dz^m dx^1 \cdots \widehat{j} \cdots dx^m \\ &= \frac{\partial L(z, x)}{\partial z^i} (-1)^{i-j} \\ &\cdot |A_i^j(x)| dz^1 \cdots dz^m (-1)^{j-1} dx^1 \cdots \widehat{j} \cdots dx^m \\ &= \frac{\partial L(z, x)}{\partial z^i} a^{ij}(x) n_j(x) d\sigma_x dz^1 \cdots dz^m = -a^{ij}(x) \\ &\cdot n_j(x) \frac{\partial L(z, x)}{\partial x^i} d\sigma_x dz^1 \cdots dz^m \\ &= -\frac{\partial L(z, x)}{\partial v_x} d\sigma_x dz^1 \cdots dz^m. \end{aligned} \tag{55}$$

Therefore,

$$\begin{aligned} V(z) &= - \int_{\Sigma} u(x) \frac{\partial L(z, x)}{\partial v_x} d\sigma_x dz^1 \cdots dz^m \\ &= V_0(z) dz^1 \cdots dz^m, \\ \delta V(z) &= (-1)^{m(m+1)+1} * d * V(z) = - * dV_0(z) \\ &= - * \frac{\partial V_0(z)}{\partial z^j} dz^j \\ &= -\frac{1}{(m-1)!} \delta_{hk_2 \cdots k_m}^{1 \cdots m} a^{jh}(z) \frac{\partial V_0(z)}{\partial z^j} dz^{k_2} \cdots dz^{k_m} \\ &= -a^{jh}(z) \frac{\partial V_0(z)}{\partial z^j} (-1)^{h-1} dz^1 \cdots \widehat{h} \cdots dz^m \\ &= -a^{jh}(z) n_h(z) \frac{\partial V_0(z)}{\partial z^j} d\sigma_z = -\frac{\partial V_0(z)}{\partial v_z} d\sigma_z. \end{aligned} \tag{56}$$

Then, if $z \in \Sigma$,

$$\lim_{z' \rightarrow z} \delta V(z') = -\frac{\partial V_0(z)}{\partial v_z} d\sigma_z \tag{57}$$

and this concludes the proof. \square

Remark 7. We note that (51) generalizes the following identity (see [7] [8, Proposition 2.2]):

$$\begin{aligned} \frac{\partial}{\partial n_z} \left(\int_{\Sigma} u(x) \frac{\partial}{\partial n_x} s(z, x) d\sigma_x \right) d\sigma_z \\ = d_z \int_{\Sigma} du(x) \wedge s_{m-2}(z, x), \quad u \in W^{1,p}(\Sigma), \end{aligned} \tag{58}$$

where $s(z, x)$ and $s_k(z, x)$ denote the fundamental solution for Laplace equation and the double k -form associated with $s(z, x)$, respectively.

We recall that if B and \tilde{B} are two Banach spaces and $C : B \rightarrow \tilde{B}$ is a continuous linear operator, we say that C can be reduced on the left if there exists a continuous linear operator $C' : \tilde{B} \rightarrow B$ such that $C'C = I + K$, where I stands for the identity operator on B and $K : B \rightarrow B$ is compact. One of the main properties of such operators is that equation $C\alpha = \beta$ has a solution if and only if $\langle \gamma, \beta \rangle = 0$ for any γ such that $C^*\gamma = 0$, C^* being the adjoint of C (see [17, 18]).

Theorem 8. Let $\tilde{J} : L^p(\Sigma) \rightarrow L^p_1(\Sigma)$ be the singular integral operator defined as

$$\begin{aligned} \tilde{J}\varphi(x) &= \int_{\Sigma} \varphi(y) d_x [S(x, y)] d\sigma_y, \\ \varphi &\in L^p(\Sigma), \quad x \in \Sigma. \end{aligned} \tag{59}$$

Then, \tilde{J} can be reduced on the left by the operator $J' : L^p_1(\Sigma) \rightarrow L^p(\Sigma)$:

$$J'\psi(z) = * \int_{\Sigma} \psi(x) \wedge d_z [\lambda_{m-2}(z, x)], \quad z \in \Sigma, \tag{60}$$

where the symbol \int_{Σ}^* means that if $\omega = \omega_0 d\sigma$ is an $(m-1)$ -form on Σ , then $\int_{\Sigma}^* \omega = \omega_0$.

Proof. We start with the observation that

$$\begin{aligned} \tilde{J}\varphi(x) &= \int_{\Sigma} \varphi(y) d_x [L(x, y)] d\sigma_y \\ &\quad + \int_{\Sigma} \varphi(y) d_x [\zeta(x, y)] d\sigma_y \\ &= J\varphi(x) + Z\varphi(x) \end{aligned} \tag{61}$$

and then

$$J'\tilde{J}\varphi = J'J\varphi + J'Z\varphi. \tag{62}$$

The operator $J'Z$ is compact because of (44). Concerning $J'J$, keeping in mind Proposition 6 and setting $u(x) = \int_{\Sigma} \varphi(y)L(x, y)d\sigma_y$, we get

$$\begin{aligned} J'J\varphi(z) &= \int_{\Sigma}^* \int_{\Sigma} \varphi(y) d_x [L(x, y)] d\sigma_y \wedge d_z [\lambda_{m-2}(z, x)] \\ &= \int_{\Sigma}^* du(x) \wedge d_z [\lambda_{m-2}(z, x)] \\ &= \frac{\partial}{\partial v_z} \int_{\Sigma} u(x) \frac{\partial}{\partial v_x} L(z, x) d\sigma_x \\ &\quad - \int_{\Sigma}^* u(x) \wedge F_z [\lambda_{m-1}(z, x)] \\ &\quad + \int_{\Sigma}^* u(x) \wedge \eta_{m-1}(z, x) \\ &= \frac{\partial}{\partial v_z} \int_{\Sigma} u(x) \frac{\partial}{\partial v_x} L(z, x) d\sigma_x + Q\varphi(z). \end{aligned} \tag{63}$$

Since $F_z[\lambda_{m-1}(z, x)] = \mathcal{O}(|z-x|^{1-m})$ and in view of (52), Q is a compact operator from $L^p(\Sigma)$ into itself.

In view of the Stokes formula for u and on account of known properties of potentials (see, e.g., [6, p. 35]), we get

$$\begin{aligned} &\frac{\partial}{\partial v_z} \int_{\Sigma} u(x) \frac{\partial}{\partial v_x} L(z, x) d\sigma_x \\ &= \frac{\partial}{\partial v_z} \left[u(z) + \int_{\Sigma} \frac{\partial}{\partial v_x} u(x) L(z, x) d\sigma_x \right] \\ &= \left(1 - \frac{1}{2}\right) \frac{\partial}{\partial v_z} u(z) + \int_{\Sigma} \frac{\partial}{\partial v_x} u(x) \frac{\partial}{\partial v_z} L(z, x) d\sigma_x \\ &= \frac{1}{2} \left(-\frac{1}{2}\varphi(z) + \int_{\Sigma} \varphi(y) \frac{\partial}{\partial v_z} L(z, y) d\sigma_y \right) \\ &\quad + \int_{\Sigma} \left[-\frac{1}{2}\varphi(x) + \int_{\Sigma} \varphi(y) \frac{\partial}{\partial v_x} L(x, y) d\sigma_y \right] \frac{\partial}{\partial v_z} L(z, x) d\sigma_x \\ &= -\frac{1}{4}\varphi(z) + \int_{\Sigma} \varphi(y) d\sigma_y \int_{\Sigma} \frac{\partial}{\partial v_x} L(x, y) \frac{\partial}{\partial v_z} L(z, x) d\sigma_x. \end{aligned} \tag{64}$$

Then,

$$\begin{aligned} J'J\varphi(z) &= -\frac{1}{4}\varphi(z) \\ &\quad + \int_{\Sigma} \varphi(y) d\sigma_y \int_{\Sigma} \frac{\partial}{\partial v_x} L(x, y) \frac{\partial}{\partial v_z} L(z, x) d\sigma_x \\ &\quad + Q\varphi(z) = -\frac{1}{4}\varphi(z) + K^2\varphi(z) + Q\varphi(z). \end{aligned} \tag{65}$$

Since $\partial/\partial v_x L(x, y) = \mathcal{O}(|x-y|^{1-m+\lambda})$, K is a compact operator.

Thus,

$$J'\tilde{J}\varphi = J'J\varphi + J'Z\varphi = -\frac{1}{4}\varphi + (K^2 + Q + J'Z)\varphi \tag{66}$$

is a Fredholm operator and the assertion is proved. \square

Theorem 9. Given $\omega \in L_1^p(\Sigma)$, there exists a solution of the singular integral equation

$$\tilde{J}\varphi(x) = \omega(x), \quad \varphi \in L^p(\Sigma), \quad x \in \Sigma \tag{67}$$

if and only if

$$\int_{\Sigma} \gamma \wedge \omega = 0 \tag{68}$$

for every weakly closed form $\gamma \in L_{m-2}^q(\Sigma)$ ($1/p + 1/q = 1$).

Proof. Denote by $\tilde{J}^* : L_{m-2}^q(\Sigma) \rightarrow L^q(\Sigma)$ the adjoint of \tilde{J} ; that is,

$$\tilde{J}^*\gamma(x) = \int_{\Sigma} \gamma(y) \wedge d_y [S(x, y)], \quad x \in \Sigma. \tag{69}$$

From Theorem 8, it follows that operator \tilde{J} can be reduced on the left; therefore, (67) admits a solution $\varphi \in L^p(\Sigma)$ if and only if

$$\int_{\Sigma} \gamma \wedge \omega = 0, \quad \forall \gamma \in L_{m-2}^q(\Sigma), \quad \tilde{J}^*\gamma = 0. \tag{70}$$

On the other hand, $\tilde{J}^*\gamma = 0$ if and only if γ is a weakly closed form; that is,

$$\int_{\Sigma} \gamma \wedge dg = 0, \quad \forall g \in C^\infty(\mathbb{R}^m). \tag{71}$$

In fact, if

$$\int_{\Sigma} \gamma(y) \wedge d_y [S(x, y)] = 0, \quad \text{a.e. } x \in \Sigma, \tag{72}$$

we have

$$\int_{\Sigma} p(x) d\sigma_x \int_{\Sigma} \gamma(y) \wedge d_y [S(x, y)] = 0, \tag{73}$$

$$\forall p \in C^\lambda(\Sigma)$$

and then

$$0 = \int_{\Sigma} \gamma(y) \wedge d_y \int_{\Sigma} p(x) S(x, y) d\sigma_x = \int_{\Sigma} \gamma \wedge du \quad (74)$$

for any smooth solution u of $Eu = 0$ in $\bar{\Omega}$. Therefore, we have

$$\int_{\Sigma} \gamma(y) \wedge d_y [S(x, y)] = 0, \quad \forall x \in T \setminus \bar{\Omega}. \quad (75)$$

Let us consider

$$z(x) = \int_{\Sigma} \gamma(y) \wedge d_y [S(x, y)], \quad x \in T. \quad (76)$$

If $v \in C^\infty(T)$ and $\eta \in C^1(\bar{\Omega}) \cap C^2(\Omega)$ are such that $E\eta = Ev$ in Ω and $\eta = 0$ on Σ , we have

$$\begin{aligned} \int_{\Omega} zEv \, dx &= \int_{\Omega} zE\eta \, dx \\ &= \int_{\Omega} E\eta(x) \, dx \int_{\Sigma} \gamma(y) \wedge d_y [S(x, y)] \\ &= \int_{\Sigma} \gamma(y) \wedge d_y \int_{\Omega} E\eta(x) S(x, y) \, dx. \end{aligned} \quad (77)$$

From the Green formulas we have

$$\int_{\Omega} S(x, y) E\eta(x) \, dx = \int_{\Sigma} S(x, y) \frac{\partial \eta}{\partial \nu}(x) \, d\sigma_x, \quad (78)$$

$y \in \Sigma.$

In view of (72), we find

$$\begin{aligned} \int_{\Omega} zEv \, dx &= \int_{\Sigma} \gamma(y) \wedge d_y \int_{\Omega} E\eta(x) S(x, y) \, dx \\ &= \int_{\Sigma} \gamma(y) \wedge d_y \int_{\Sigma} \frac{\partial \eta}{\partial \nu}(x) S(x, y) \, d\sigma_x \\ &= \int_{\Sigma} \frac{\partial \eta}{\partial \nu}(x) \, d\sigma_x \int_{\Sigma} \gamma(y) \wedge d_y [S(x, y)] \\ &= 0. \end{aligned} \quad (79)$$

We have proved that $z = 0$ on Σ , $z = 0$ in Ω , and then $z = 0$ in T . Therefore,

$$\begin{aligned} 0 &= \int_T zE\varphi \, dx = \int_T E\varphi \, dx \int_{\Sigma} \gamma(y) \wedge d_y [S(x, y)] \\ &= \int_{\Sigma} \gamma(y) \wedge d_y \int_T E\varphi(x) S(x, y) \, dx \\ &= \int_{\Sigma} \gamma(y) \wedge d\varphi(y), \end{aligned} \quad (80)$$

for any $\varphi \in \dot{C}^\infty(T)$. This implies (71) and the theorem is proved. \square

Lemma 10. For every $f \in W^{1,p}(\Sigma)$, there exists a solution of the boundary value problem

$$\begin{aligned} w &\in \mathcal{S}^p, \\ Ew &= 0 \quad \text{in } \Omega, \\ dw &= df \quad \text{on } \Sigma. \end{aligned} \quad (81)$$

Its solution w is a simple layer potential (49) whose density φ solves $\bar{f}\varphi = df$ (see (59)).

Proof. Consider the following singular integral equation:

$$\int_{\Sigma} \varphi(y) \, d_x [S(x, y)] \, d\sigma_y = df(x), \quad x \in \Sigma, \quad (82)$$

in which the unknown is $\varphi \in L^p(\Sigma)$ and the datum is $df \in L^p_1(\Sigma)$. With conditions (68) being satisfied, in view of Theorem 9 there exists a solution φ of (82). \square

Lemma 11. Let \mathcal{A} be the eigenspace of the Fredholm integral equation

$$-\frac{1}{2}\psi(x) + \int_{\Sigma} \psi(y) \frac{\partial}{\partial \nu_x} S(x, y) \, d\sigma_y = 0, \quad (83)$$

a.e. $x \in \Sigma.$

The dimension of \mathcal{A} is 1.

Proof. The Fredholm equation (83) has the same number of linearly independent solutions of the following equation:

$$-\frac{1}{2}\gamma(x) + \int_{\Sigma} \gamma(y) \frac{\partial}{\partial \nu_y} S(x, y) \, d\sigma_y = 0, \quad \text{a.e. } x \in \Sigma. \quad (84)$$

Obviously, the constant functions are eigensolutions of (84). We want to show that they are the only ones. Let γ_1 and γ_2 be two linearly independent eigensolutions of (84) and set

$$u_i(x) = \int_{\Sigma} \gamma_i(y) S(x, y) \, d\sigma_y, \quad i = 1, 2. \quad (85)$$

We note that γ_1 and γ_2 are Hölder continuous functions. With potentials u_i being smooth solutions of the problem

$$\begin{aligned} Eu &= 0 \quad \text{in } \Omega, \\ \frac{\partial u}{\partial \nu^+} &= 0 \quad \text{on } \Sigma, \end{aligned} \quad (86)$$

we get $u_i = \alpha_i$ in Ω . Choose $(c_1, c_2) \neq (0, 0)$ such that $c_1\alpha_1 + c_2\alpha_2 = 0$ and set

$$u(x) = \int_{\Sigma} (c_1\gamma_1(y) + c_2\gamma_2(y)) S(x, y) \, d\sigma_y. \quad (87)$$

Since $u = c_1\alpha_1 + c_2\alpha_2 = 0$ in Ω , u satisfies the following boundary value problem:

$$\begin{aligned} Eu &= 0 \quad \text{in } T \setminus \bar{\Omega}, \\ u &= 0 \quad \text{on } \Sigma, \\ u &= 0 \quad \text{on } \partial T. \end{aligned} \quad (88)$$

By Green's formula, $u = 0$ in $T \setminus \Omega$ and therefore $u = 0$ in T . This implies $c_1\gamma_1 + c_2\gamma_2 = 0$, which is a contradiction. \square

Lemma 12. *Given $c \in \mathbb{R}$, there exists a solution of the following boundary value problem:*

$$\begin{aligned} v &\in \mathcal{S}^p, \\ Ev &= 0 \quad \text{in } \Omega, \\ v &= c \quad \text{on } \Sigma. \end{aligned} \tag{89}$$

It is given by

$$v(x) = c \int_{\Sigma} \psi_0(y) S(x, y) d\sigma_y, \quad x \in \Omega, \tag{90}$$

where ψ_0 is the unique element of \mathcal{A} such that

$$\int_{\Sigma} \psi_0(y) S(x, y) d\sigma_y = 1, \quad \forall x \in \bar{\Omega}. \tag{91}$$

Proof. Let $\psi \in \mathcal{A}$, $\psi \neq 0$. Setting

$$P\psi(x) = \int_{\Sigma} \psi(y) S(x, y) d\sigma_y \tag{92}$$

we have that $P\psi = c$ in Ω . As in Lemma 11, this implies that if $c = 0$, we have that $\psi = 0$. Then, $c \neq 0$. Function $\psi_0 = (1/c)\psi$ satisfies (91) and v given by (90) is solution of (89). \square

Theorem 13. *The Dirichlet problem (50) has a unique solution for every $f \in W^{1,p}(\Sigma)$. In particular, the density φ of u can be written as $\varphi = \varphi_0 + \psi$, where φ_0 solves the singular integral system*

$$\int_{\Sigma} \varphi_0(y) d_x [S(x, y)] d\sigma_y = df(x), \quad a.e. x \in \Sigma \tag{93}$$

and $\psi \in \mathcal{A}$.

Proof. Let w be a solution of the boundary value problem (81). Since $dw = df$ on Σ , $w = f - c$ on Σ for some $c \in \mathbb{R}$. Function $u = w + v$, where v is given by (90), solves problem (50).

Consider now two solutions of the same problem (50):

$$u(x) = \int_{\Sigma} \varphi(y) S(x, y) d\sigma_y, \tag{94}$$

$$u'(x) = \int_{\Sigma} \varphi'(y) S(x, y) d\sigma_y.$$

Therefore, the potential

$$v(x) = \int_{\Sigma} \psi(y) S(x, y) d\sigma_y, \tag{95}$$

where $\psi = \varphi - \varphi'$, solves the problem

$$\begin{aligned} v &\in \mathcal{S}^p, \\ Ev &= 0 \quad \text{in } \Omega, \\ v &= 0 \quad \text{on } \Sigma. \end{aligned} \tag{96}$$

Since

$$\int_{\Sigma} \psi(y) d_x [S(x, y)] d\sigma_y = 0 \quad \text{on } \Sigma, \tag{97}$$

we have $J'\tilde{J}\psi = 0$ (see (66)). By standard arguments, ψ is Hölder continuous and then $v \in C^0(\bar{\Omega}) \cap C^2(\Omega)$. The weak maximum principle (see, e.g., [19, p. 32]) shows that $v = 0$ in Ω ; that is, $u = u'$. \square

We end this section by observing that when we study the Dirichlet problem (50), we need to solve the singular integral equation $\tilde{J}\varphi = df$, $\varphi \in L^p(\Sigma)$. We have proved that this equation can be reduced to a Fredholm one by means of the operator J' . This reduction is not an equivalent reduction in the usual sense (see, e.g., [18, pp. 19-20]); that is, it is not true that $\mathcal{N}(J') = \{0\}$, $\mathcal{N}(J')$ being the kernel of the operator J' . However, if the condition

$$\mathcal{N}(J'\tilde{J}) = \mathcal{N}(\tilde{J}) \tag{98}$$

is true, J' still provides equivalence in a certain sense. In fact, we have the following lemma.

Lemma 14. *If condition (98) holds, the singular integral equation (82) is equivalent to the Fredholm equation $J'\tilde{J}\varphi = J'df$.*

Proof. Condition (98) implies that if g is such that there exists a solution φ of the equation $\tilde{J}\varphi = g$, then this equation is satisfied if and only if $J'\tilde{J}\varphi = J'g$. Since the equation $\tilde{J}\varphi = df$ is solvable (see Lemma 10), we have that $\tilde{J}\varphi = df$ if and only if $J'\tilde{J}\varphi = J'df$. \square

Condition (98) is satisfied, for example, if the differential operator E has constant coefficients. This can be proved as in [20, Remark 1, p. 1045], replacing the Laplacian and the normal derivative by E and the conormal derivative, respectively.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1] O. Chkadua, S. E. Mikhailov, and D. Natroshvili, "Analysis of segregated boundary-domain integral equations for variable-coefficient problems with cracks," *Numerical Methods for Partial Differential Equations*, vol. 27, no. 1, pp. 121-140, 2011.
- [2] O. Chkadua, S. E. Mikhailov, and D. Natroshvili, "Localized boundarydomain singular integral equations based on harmonic parametrix for divergence-form elliptic PDEs with variable matrix coefficients," *Integral Equations and Operator Theory*, vol. 76, no. 4, pp. 509-547, 2013.
- [3] G. C. Hsiao and W. L. Wendland, *Boundary Integral Equations*, Springer, Berlin, Germany, 2008.
- [4] M. Kohr, M. Lanza De Cristoforis, and W. L. Wendland, "Poisson problems for semilinear Brinkman systems on Lipschitz domains in \mathbb{R}^n ," *Zeitschrift für Angewandte Mathematik und Physik*, vol. 66, no. 3, pp. 833-864, 2015.
- [5] W. McLean, *Strongly Elliptic Systems and Boundary Integral Equations*, Cambridge University Press, Cambridge, UK, 2000.

- [6] C. Miranda, *Partial Differential Equations of Elliptic Type*, Springer, New York, NY, USA, 1970.
- [7] A. Cialdea, "On oblique derivate problem for Laplace equation and connected topics," *Rendiconti della Accademia Nazionale delle Scienze detta dei XL: Memorie di Matematica e Applicazioni Serie 5*, vol. 12, no. 1, pp. 181–200, 1988.
- [8] A. Cialdea, E. Dolce, V. Leonessa, and A. Malaspina, "On the potential theory in Cosserat elasticity," *Bulletin of TICMI*, vol. 18, no. 2, pp. 67–81, 2014.
- [9] A. Cialdea, E. Dolce, V. Leonessa, and A. Malaspina, "New integral representations in the linear theory of viscoelastic materials with voids," *Publications de l'Institut Mathematique*, vol. 96, no. 110, pp. 49–65, 2014.
- [10] A. Cialdea, E. Dolce, and A. Malaspina, "A complement to potential theory in the Cosserat elasticity," *Mathematical Methods in the Applied Sciences*, vol. 38, no. 3, pp. 537–547, 2015.
- [11] A. Cialdea, V. Leonessa, and A. Malaspina, "On the dirichlet problem for the stokes system in multiply connected domains," *Abstract and Applied Analysis*, vol. 2013, Article ID 765020, 12 pages, 2013.
- [12] G. Fichera, "Linear elliptic equations of higher order in two independent variables and singular equations, with applications to anisotropic inhomogeneous elasticity," in *Partial Differential Equations and Continuum Mechanics*, R. E. Langer, Ed., pp. 55–80, University of Wisconsin Press, Madison, Wis, USA, 1961.
- [13] G. Fichera, "Spazi lineari di k -misure e di forme differenziali," in *Proceedings of the International Symposium on Linear Spaces*, pp. 175–226, Hebrew University of Jerusalem, Jerusalem, Israel, July 1960.
- [14] H. Flanders, *Differential Forms with Applications to the Physical Science*, Academic Press, New York, NY, USA, 1963.
- [15] D. Lovelock and H. Rund, *Tensor, Differential Forms, and Variational Principles. Pure and Applied Mathematics*, Wiley-Interscience, New York, NY, USA, 1975.
- [16] M. P. Colautti, "Sui problemi variazionali di un corpo elastico incompressibile," *Atti della Accademia Nazionale dei Lincei. Memorie. Classe di Scienze Fisiche, Matematiche e Naturali. Sezione Ia. Matematica, Meccanica, Astronomia, Geodesia e Geofisica. Serie VIII*, vol. 8, no. 8, pp. 291–343, 1967.
- [17] G. Fichera, "Una introduzione alla teoria delle equazioni integrali singolari," *Rendiconti di Matematica e Delle sue Applicazioni*, vol. 19, no. 17, pp. 82–191, 1958, Reprinted in *Integrali Singolari e Questioni Connesse*, "Singular integrals and related questions" *Lectures from the Centro Internazionale Matematico Estivo (C.I.M.E.) Summer School held in Varenna*, June 10–19, 1957. Edited by G. Fichera and E. Magenes (Italian).
- [18] S. G. Mikhlin and S. Prössdorf, *Singular Integral Operators*, Springer, Berlin, Germany, 1986.
- [19] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Classics in Mathematics, Springer-Verlag, Berlin, Germany, 2001.
- [20] A. Cialdea, V. Leonessa, and A. Malaspina, "On the Dirichlet and the Neumann problems for Laplace equation in multiply connected domains," *Complex Variables and Elliptic Equations*, vol. 57, no. 10, pp. 1035–1054, 2012.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

