



Long-term thinning effects on tree growth, drought response and water use efficiency at two Aleppo pine plantations in Spain

Àngela Manrique-Alba^{a,*}, Santiago Beguería^a, Antonio J. Molina^b, María González-Sanchis^b, Miquel Tomàs-Burguera^a, Antonio D. del Campo^b, Michele Colangelo^{c,d}, J. Julio Camarero^c

^a Estación Experimental Aula Dei (EEAD-CSIC), Zaragoza, Spain

^b Research Group in Forest Science and Technology (Re-ForeST), Research Institute of Water and Environmental Engineering (IIAMA), Universitat Politècnica de València, Valencia, Spain


^c Instituto Pirenaico de Ecología (IPE-CSIC), Zaragoza, Spain

^d School of Agricultural, Forest, Food and Environmental Sciences, University of Basilicata, Potenza, Italy

HIGHLIGHTS

- Thinning enhanced tree growth on Mediterranean Aleppo pine forests.
- Thinning reduced growth dependence on water stress (SPEI).
- Thinning improved drought resilience, making pines better adapted to unfavourable conditions.
- Effects of thinning remained at least for 20 years, especially in the heavy thinning treatment.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 28 January 2020

Received in revised form 2 April 2020

Accepted 5 April 2020

Available online 18 April 2020

Editor: Jose Ignacio Querejeta

Keywords:

Pinus halepensis

Thinning

Water use efficiency

Oxygen isotopes

Drought stress

Dendroecology

ABSTRACT

In Mediterranean areas where drought-induced forest dieback and tree mortality have been widely reported, it is still under debate how the likely risks of climate change will affect tree growth and consequently forest productivity. Increasing tree mortality has been associated not only to increased drought, but also to a lack of management in many dense pine forests and plantations, where warming may intensify tree-to-tree competition for soil water. This emphasizes the need of using silviculture to adapt dense stands of Mediterranean pine reforestations to warmer and drier conditions. Here we combined dendrochronology and C and O isotope analyses of wood in two Aleppo pine (*Pinus halepensis*) plantations, growing under semiarid conditions and experimentally thinned at high and moderate intensities along with control. The main aim was to understand the responses of radial growth and water use efficiency (WUEi) to different thinning intensities, and to analyze the effectiveness of thinning to enhance post-drought growth resilience. Thinning had a positive effect on growth, produced an increase of $\delta^{18}\text{O}$, reduced growth sensitivity to drought and decreased WUEi, suggesting a reduction of drought stress. These results were consistent across sites, and were significant even 20 years after the intervention took place. Considering the climate effects on growth through the SPEI drought index to calculate resistance and recovery indices, an increase of resistance after thinning was observed. We conclude that high thinning intensity (50%

* Corresponding author.

E-mail address: amanrique@eead.csic.es (À. Manrique-Alba).

of basal area removed) is a useful silviculture intervention on Mediterranean Aleppo pine plantations that enhances their growth, and makes them less dependent on harsh climatic conditions, improving their resilience against drought and consequently making them better adapted to more unfavourable conditions.

© 2020 Elsevier B.V. All rights reserved.

1. Introduction

Aleppo pine (*Pinus halepensis* Mill.) has been used extensively in afforestation programs across the Mediterranean Basin during the last 100 years (Maestre and Cortina, 2004). Covering over 25,000 km² in the Mediterranean Basin (Quézel, 2000). It is one of the most used tree species for reforestation under semiarid conditions (Pemán et al., 2009), being for instance representative in a 70% of the reforested area in eastern Spain (Vallejo and Alloza, 1998). Aleppo pine forests provide hydrological cycle stabilization, soil protection (Fady et al., 2003) and wood products. In some cases, afforestation with *P. halepensis* has promoted the establishment of high-density plantations characterized by slow-growing rates which are prone to fire and pest attacks (Maestre and Cortina, 2004), and more predisposed to forest dieback following extreme drought events (Carnicer et al., 2011; García de la Serrana et al., 2015). Severe canopy dieback and increasing tree mortality in pine plantations has been associated not only to increased drought but also to a lack of forest management, which may intensify inter-tree competition (Sánchez-Salguero et al., 2012, 2015).

Understanding how forest structure modulates the physiological response of trees to drought could help managers to adopt strategies for improving the resistance and resilience against drought (Moreno-Gutiérrez et al., 2011). This is especially critical in Mediterranean areas within the context of climate change, where is still debated how climate change impacts will affect tree growth and forest productivity in addition to more recurrent events of tree mortality and forest dieback (Martinez-Vilalta and Pinol, 2002; Allen et al., 2010; McDowell and Allen, 2015).

Forest density reduction treatments (i.e., thinning) offer a strategy for potentially mitigating these effects not only by improving forest health but also the hydrological balance through increased soil infiltration and streamflow (del Campo et al., 2019; Tague et al., 2019). By means of reducing the competition for water, light and nutrients, thinning has been observed to increase both the resistance (i.e., the capacity to maintain growth levels during drought) and the resilience (i.e., the ability to recover to the pre-drought growth levels) of trees in case of drought (Aussenac, 2000; Giuggiola et al., 2013; Sohn et al., 2013; Olivar et al., 2014). However, the response of trees to drought varied with thinning intensity (Molina and del Campo, 2012), stand density, age, the site climate conditions (Sohn et al., 2016), and the time since thinning intervention (del Campo et al., 2014). Other policy-relevant questions remain open, such as what the optimum thinning intensity is and how long the effects last. In order to answer these questions, long-term thinning experiments with different intervention intensities are required. For example, Moreno-Gutiérrez et al. (2011) assessed the impacts of thinning on growth in *P. halepensis* in a very short-term scale (12 months after thinning), and del Campo et al. (2018, 2019) for a period of 3 years after thinning. Long-term thinning studies are scarce and limited to other pine species, generally growing in high-productivity regions (Mäkinen and Isomäki, 2004; Prescott et al., 2019). Also, these studies focused on stem production instead of on understanding physiological mechanisms related to growth. To the authors' knowledge, no long-term results have been reported specifically for Aleppo pine.

Analysis of radial-growth rates using tree rings offers a convenient way to characterize tree responses to environmental stressors and it can be used as a vitality indicator, as growth decline can be considered an early-warning signal of forest dieback (Dobbertin, 2005; Camarero

et al., 2015). In addition, the analysis of stable isotopes (C, O) in tree-ring wood may allow to obtain complementary information about the water and carbon economy of trees (Scheidegger et al., 2000; McCarroll and Loader, 2004; Gessler et al., 2018). The composition of carbon isotopes ($\delta^{13}\text{C}$) is a proxy of intrinsic water-use efficiency (WUEi) in C3 plants (Farquhar et al., 1989; Klein et al., 2005). The WUEi is the ratio between the net photosynthetic rate (A) and the stomatal conductance (g_s) (Farquhar et al., 1989). Reduction of WUEi after a thinning treatment can be expected as a result of decreasing competition for soil water, and such an effect has been observed in several empirical studies regarding pine species (McDowell et al., 2006; Giuggiola et al., 2016; Navarro-Cerrillo et al., 2019). However, other authors reported no variation in WUEi after thinning (Martin-Benito et al., 2010; Moreno-Gutiérrez et al., 2011), so the question remains open.

Several physiological adaptations determine WUEi adjustments after thinning: (i) variations in g_s at constant A ; (ii) variations in A at constant g_s ; and (iii) variations of both, with that of g_s or A being stronger. The dual-isotope approach based on the simultaneous analysis of $\delta^{13}\text{C}$ and oxygen stable isotopes ($\delta^{18}\text{O}$) has been proposed as a means to disentangle the influences of A and g_s on WUEi, since $\delta^{18}\text{O}$ is related to g_s (Scheidegger et al., 2000; Barbour, 2007), especially in drought-sensitive tree species (Battipaglia et al., 2014). However, while $\delta^{13}\text{C}$ responses to thinning have been described in previous studies considering different pine species and a variety of precipitation regimes, only a few of them combined $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ (McDowell et al., 2006; Martin-Benito et al., 2010; Moreno-Gutiérrez et al., 2011; Giuggiola et al., 2016; Fernandes et al., 2016), with varying responses and interpretation of the results.

We used dendrochronology and stable isotopes ($\delta^{13}\text{C}$, $\delta^{18}\text{O}$) analyses to evaluate the growth and drought-climate relationships and to assess the physiological responses under different thinning intensities (medium and high intensity thinning treatments) and different times after thinning intervention (ten and twenty years after the thinning treatments) in two planted stands of Aleppo pine in Spain. We selected two sites with similar Mediterranean climatic conditions and performed the same analyses in order to determine the robustness of the results (that is, we treated the sites as replicates). The main objectives of this research were: (i) to assess the response of radial growth and WUEi to different thinning intensities and different time since thinning intervention and their relationship with climate; (ii) to disentangle the effects of thinning on WUEi through complementary measurements of oxygen stable isotopes; and (iii) to analyze the effectiveness of thinning on improving growth indices. We hypothesize that thinning will improve radial growth and resilience of trees against drought, attributable to a relaxation of drought stress. In thinned plots, we expect a decrease of WUEi, caused by a reduction of tree competition for soil water and nutrients. Regarding the physiological mechanisms, we hypothesize that combined analysis of $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ would allow us to determine the relative influence of changes in A and g_s on WUEi and growth.

2. Material and methods

2.1. Study sites and thinning experiments

We selected two areas afforested with *P. halepensis* located in eastern Spain: Tarazona (41° 52' N, 1° 50' W, 798 m a.s.l.) and La Hundo (39° 05' N, 1° 12' W, 950 m a.s.l.; Fig. 1). The two sites are characterized by similar Mediterranean climatic conditions. To describe the climatic



Fig. 1. Location of sites at eastern Spain and view of the thinning treatments studied in Tarazona and La Hundo. Note that pictures were taken in 2018 for Tarazona while in 2008 for La Hundo (with time periods of 20 years and 1 month from thinning intervention, respectively).

characteristics of both sites we used the 1.1-km² gridded climate data for the 1981 to 2010 period of temperature, precipitation (Vicente-Serrano et al., 2017) and reference evapotranspiration (Tomas-Burguera et al., 2019). Mean annual temperature ranges from 12.4 °C (s.d. of 0.5 °C) at Tarazona to 13.2 °C (s.d. of 0.6 °C) at La Hundo. In Tarazona and La Hundo, the mean annual Penman-Monteith reference evapotranspiration is 1121 mm and 1217 mm, respectively, whereas the mean annual precipitation varies from 415 mm (s.d. of 95 mm) to 484 mm (s.d. of 143 mm). The hydrologic balance (annual precipitation minus potential evapotranspiration) is −706 mm (s.d. 117 mm) in Tarazona, and −733 mm (s.d. 167 mm) in La Hundo. Both locations have similar seasonal temperature and precipitation distribution (Fig. S2), characterized by a strong intra-annual variation with a dry period in summer and two maxima in spring and fall. The maximum rainfall in La Hundo occurs in fall (October, 62 mm) due to its closer location to the Mediterranean Sea, while Tarazona shows its maximum in spring (May, 58 mm), which is a typical feature of a more continental Mediterranean climate subjected to Atlantic influences. In fact, following the Köppen classification, only La Hundo adjusts to a purely Mediterranean climate ('Csa'), while Tarazona is classified as Temperate Oceanic ('Cfb'). Despite this difference in the Köppen classification, both sites were considered as Mediterranean as both showed a marked arid summer period.

The studied *P. halepensis* plantations were established at both sites between the late 1940s and the early 1960s with the main objectives of improving the ecological status and reducing soil erosion. No forest management has been carried out until the thinning experiments took place. In the Tarazona site the plots contained scattered sclerophyllous shrubland vegetation (*Quercus coccifera* L., *Juniperus thurifera* L., *Rhamnus alaternus* L., and *Rhamnus lycioides* L.), whereas in La Hundo the plots presented scant presence of understory species (e.g., *Quercus ilex* sbsp. *Ballota* (Desf.) Samp., *Juniperus oxycedrus* L. and *Juniperus phoenicea* L.). At both sites understory was not removed during the thinning intervention nor afterwards. The soils, according to the World Reference Base for Soil Resources (WRB, 2014), correspond to Cambisols and Leptosols for Tarazona and La Hundo, respectively. The soil depth is around 60 cm in both sites. In both study sites the slope gradient is <5%.

Experimental thinning treatments took place in different years and included several intensities at both sites. In Tarazona the experimental thinning was carried out in 1998 while in La Hundo was performed in 2008. In Tarazona the initial canopy cover of 81% (control plot, C) was

diminished to 62% (moderate intensity plot, M) and 40% (high intensity plot, H). In La Hundo the initial canopy cover was reduced from approximately 84% (control plot, C) to 68% (moderate thinning intensity plot, M) and 50% (high thinning intensity plot, H). For additional details of thinning intervention in La Hundo see previous works by Molina and del Campo (2012), and del Campo et al. (2014) and for Tarazona see Hernández and Arrechea (2017). In both sites the plots show similar forest structure and tree characteristics (Table 1).

2.2. Dendrochronology

In 2018 we selected 15 *P. halepensis* trees per site and per treatment and extracted 2 cores per tree at breast height (1.3 m) separated by 180° using Pressler borers. A total of 180 samples from 90 trees (two samples × 15 trees × three treatments × two sites) were thus obtained. Samples were taken from dominant and mature trees showing no visible signs of damage. The diameter at 1.3 m of all sample trees and the total height of 5 selected trees per plot were measured using tapes and clinometers, respectively.

The wood samples were visually cross-dated, and total tree-ring width (RW) was measured under a stereo microscope with an accuracy of 0.01 mm using the TSAP-Win program LINTAB measuring device (Rinntech, Heidelberg, Germany). The dating was checked using the COFECHA program (Holmes, 1983). The RW was transformed into

Table 1

Characteristics of the *Pinus halepensis* stands and trees immediately after thinning interventions in the study sites (values are means). Note that relative changes when comparing forest structure metrics among control and thinned treatments are consistent between the sites.

Site/thinning treatment	Diameter at 1.3 m (cm)	Tree height (m)	Canopy cover (%)	Density (trees ha ⁻¹)	Basal area (m ² ha ⁻¹)
Tarazona					
Control	13.6	9.2	81	2000	31.0
Moderate	14.5	9.3	62	1050	21.3
High	14.6	8.9	40	975	15.5
La Hundo					
Control	16.9	11.5	84	1489	40.1
Moderate	17.7	12.2	68	744	27.2
High	17.5	11.3	50	478	18.2

basal area increment (BAI) which is biologically more meaningful to quantify growth variations between years and treatments. BAI values were calculated annually from each RW measurements assuming a circular shape of stems as the difference between consecutive annual cross-sectional basal areas. We are aware that such assumption may introduce additional uncertainty since stems are not perfectly circular. However, BAI is still a better biological descriptor of growth than tree-ring width. The mean BAI of the two cores for each tree was computed, and the resulting mean BAI series were further used in all statistical analyses. The years prior to 1965 were not considered due to the juvenile age of the trees.

2.3. Climatic data and identification of drought years

The Standardized Precipitation Evapotranspiration Index (SPEI, unitless), was used to characterize drought conditions at several time scales (1 to 48 months). The SPEI is a multi-scalar drought index based on anomalies of the climatic water balance with respect to the normal conditions at a given site, and it allows determining the onset and magnitude of droughts (Vicente-Serrano et al., 2010). The SPEI is computed as the difference between precipitation and potential evapotranspiration (the climatic water balance) accumulated over a given time period (the so called 'time scale'). The values are standardized after fitting a log-logistic distribution to the data, so the SPEI is finally expressed as standard deviations following a standard Gaussian distribution (Beguería et al., 2014). SPEI weekly time series were downloaded from <http://monitordesequia.csic.es> (Vicente-Serrano et al., 2017) using the sites' coordinates.

Since the SPEI data consisted on weekly values, and there are different variants of the index depending on the time scale (i.e., the period over which the climatic water balance is calculated), we undertook a preliminary analysis in order to determine the time scale and week of the year that had the best correlation with the BAI. Pearson correlation coefficients between alternative SPEI and BAI time series (from October of the previous year to December of the growing year, that is a period of 15 months) were thus computed, allowing to determine the best combination at each site. Based on the SPEI data, the 1986–1987, 1994–1995, 2005–2006 and 2012 droughts were considered as the most severe in the two study sites. In addition, for La Hundo 1999–2000 and 2014 were also included as droughts.

2.4. Resistance, recovery and resilience indices

The set of selected droughts included events before and after the thinning treatments. In order to quantify the responses of BAI to the selected droughts and to explore the effects of management on growth, we followed Lloret et al. (2011) who defined resistance, recovery and resilience indices. The resistance index (Rt) quantifies the capacity to maintain growth levels during a drought; the recovery index (Rc) measures the growth increase after a drought; and the resilience index (Rs) reports the ability to recover pre-drought growth levels after a drought. The indices are computed as follows:

$$\text{Resistance (Rt)} = \text{BAI}_D / \text{BAI}_{\text{Pre-D}} \quad (1)$$

$$\text{Recovery (Rc)} = \text{BAI}_{\text{Post-D}} / \text{BAI}_D \quad (2)$$

$$\text{Resilience (Rs)} = \text{BAI}_{\text{Post-D}} / \text{BAI}_{\text{Pre-D}} \quad (3)$$

where BAI_D corresponds to BAI during the drought year, and the $\text{BAI}_{\text{Pre-D}}$ and $\text{BAI}_{\text{Post-D}}$ correspond to the average BAI of the 2 years previous to the drought event and after the drought, respectively. Sohn et al. (2016) suggested to use 3 years before and after the drought event, however we selected 2-years long for the pre- and post-drought periods in order to avoid drought events overlap and following Camarero et al. (2018). This is also in agreement with 2-year period established as a

global reference for post-drought recovery by Anderegg et al. (2015). The resilience indices were calculated for the selected drought episodes.

2.5. Stable isotope analysis and intrinsic water use efficiency

Wood tissue samples were taken for C and O isotope analyses, as studies comparing whole wood and cellulose show similar trends (Saurer et al., 2004). For each treatment (Control and Heavy thinning), we selected the five trees whose cross-dated tree-ring width series presented the highest correlation with the mean series. For the five trees selected per treatment, rings were separated in years using scalpels, starting in 1992 and sampling every year onwards. Then, year pools were obtained by making composite samples of the five selected trees. The resulting wood samples were homogenized to a fine powder using a ball mixer mill (Retsch MM301, Haan, Germany). Wood aliquots (0.001 g) were weighted on a microbalance (AX205 Mettler Toledo, OH, USA). In the case of $\delta^{13}\text{C}$, samples were stored in tin foil capsules and combusted to CO_2 using a Flash EA-1112 elemental analyser interfaced with a Finnigan MAT Delta C isotope ratio mass spectrometer (Thermo Fisher Scientific Inc., MA, USA). In the case of $\delta^{18}\text{O}$, samples were stored in silver foil capsules and combusted in an elementary PyroCube (Elementar Analysensysteme GmbH, Hanau, Germany) interfaced to a PDZ Europa 20-20 isotope ratio mass spectrometer (Sercon Ltd., Cheshire, UK). Isotope analyses were performed at the Stable Isotope Facility of the University of California at Davis (USA). Stable isotope ratios were expressed as per mil (‰) deviations using the δ notation relative to VPDB (for carbon) and VSMOW (for oxygen) standards. The standard deviation for repeated analyses was $\pm 0.06\text{‰}$ and $\pm 0.25\text{‰}$ for $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$, respectively.

We calculated WUEi values from tree-ring wood C isotopes using the following framework. We calculated C isotope discrimination in wood ($\Delta^{13}\text{C}$) based on changes in $\delta^{13}\text{C}$ of atmospheric CO_2 ($\delta^{13}\text{C}_a$) and wood $\delta^{13}\text{C}$ ($\delta^{13}\text{C}_w$) following Farquhar and Richards (1984):

$$\Delta^{13}\text{C} = (\delta^{13}\text{C}_a - \delta^{13}\text{C}_w) / (1 + \delta^{13}\text{C}_w / 1000) \quad (4)$$

$\delta^{13}\text{C}_a$ was obtained from McCarroll and Loader (2004). Then, we calculated WUEi (in $\mu\text{mol mol}^{-1}$) using the equation proposed by Farquhar et al. (1982):

$$\text{WUEi} = c_a [1 - (c_i / c_a)] 0.625 = (c_a - c_i) / 1.6 \quad (5)$$

where c_a is the atmospheric CO_2 concentration, c_i is the CO_2 concentration in the sub-stomatal cavity of leaves, and 0.625 is the relation among the conductance of H_2O compared to the conductance of CO_2 . To determine c_i we followed Francey and Farquhar (1982):

$$c_i = [(\Delta^{13}\text{C} - a) \times c_a] / (b - a) \quad (6)$$

where a is the diffusion fractionation across the boundary layer and the stomata (4.4‰) and b is the Rubisco fractionation (27.0‰).

The relationships between $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ were analysed by means of the differences between thinned and control plots ($\Delta\delta^{13}\text{C}$ and $\Delta\delta^{18}\text{O}$, respectively), during the pre- and post-thinning periods.

2.6. Statistical analyses

Univariate ANOVA was used for testing for differences in BAI, $\delta^{18}\text{O}$ and WUEi comparing between pre- and post-treatment periods and between treatment levels. In Tarazona, differences were tested for two post-treatment periods of ten years each. ANOVA was also used to test for differences between treatments in resistance, recovery and resilience indices for each drought year, using the Tukey's significant difference post hoc test. For multi-year droughts the last year was used, and prior to the analysis we examined all the indices for normality and homoscedasticity.

One problem with ANOVA was that it did not allow for controlling for the influence of other relevant factors besides the period or the treatment levels, such as the climate. Climate has a strong influence on both the growth and water use efficiency of trees, and differences in climate between pre and post thinning periods could alter the results when comparing based on ANOVA. Therefore, we used a set of multivariate linear models in order to test for treatment differences in BAI, WUEi and $\delta^{18}\text{O}$ while controlling for other relevant factors such as differences in climate (SPEI) or random differences among plots or trees. This also allowed us to test for differences in the sensitivity to the SPEI between periods or among treatments.

In the case of BAI we fitted a series of generalized linear mixed-effects models (GLMM) including the treatment intensity (control, C; medium, M; and high, H) and the time period (previous to thinning, *preT*; and posterior to thinning, *postT*) and the SPEI as fixed factors, and the tree labels as random factors. A Gamma GLMM with a logarithmic link was chosen as the model better fitting the skewed shape of the data. Separated models were fitted for each site, leading to two different sets of parameters. Model configurations of increasing complexity were fitted:

$$\begin{aligned} \text{M0: } \mu(t, p, n) &= \beta_0(t, p) + \beta_1 \text{ SPEI}_i + v(n) + \varepsilon \\ \text{M1: } \mu(t, p, n) &= \beta_0(t, p) + \beta_1(p) \text{ SPEI}_i + v(n) + \varepsilon \\ \text{M2: } \mu(t, p, n) &= \beta_0(t, p) + \beta_1(t) \text{ SPEI}_i + v(n) + \varepsilon \\ \text{M3: } \mu(t, p, n) &= \beta_0(p) + \beta_1(t, p) \text{ SPEI}_i + v(n) + \varepsilon \\ \text{M4: } \mu(t, p, n) &= \beta_0(t, p) + \beta_1(t, p) \text{ SPEI}_i + v(n) + \varepsilon \end{aligned} \quad (7)$$

where $\mu(t, p, n) = \log(\text{BAI}(t, p, n))$ is the dependent variable for a given site, treatment (t), time period (p) and tree (n); β_0 is an intercept; β_1 is the climate effect (SPEI); v is the individual-level random effect following a Normal distribution with mean 0; and ε is a Gamma-distributed random error. The subscripts (t, p) refer to interactions between the model coefficients (β) and the treatment and time period factors, so for instance $\beta_1 \text{ SPEI}$ indicates that the coefficient does not vary between periods and treatments, while $\beta_1(t, p) \text{ SPEI}$ indicates a full-interaction configuration in which the coefficient varies between both treatments and periods. The different models include, thus, configurations with equal slopes and different intercepts; equal intercepts and different slopes; and different slopes and intercepts. The models were fitted using the `lme4` R package (Bates et al., 2015), and the Akaike's Information Criterion (AIC) was used to compare between them. The AIC was used because it provides a balance between goodness-of-fit and model complexity. It must be noted that mixed models do not allow for computing p -values of the fixed-effects model coefficients due to the existence of random effects, so we used the overlapping of the 95% confidence bands around the model coefficients to determine the significance of differences between treatment levels and between periods.

For $\delta^{18}\text{O}$ and WUEi we fitted linear models, since there were no data for individual trees (as stated, a pooled sample was used for the isotopic analysis), so using a mixed-effects model was not possible. Similar model configurations were compared:

$$\begin{aligned} \text{M0: } \mu(t, p) &= \beta_0(t, p) + \beta_1 \text{ SPEI}_i + \varepsilon \\ \text{M1: } \mu(t, p) &= \beta_0(t, p) + \beta_1(p) \text{ SPEI}_i + \varepsilon \\ \text{M2: } \mu(t, p) &= \beta_0(t, p) + \beta_1(t) \text{ SPEI}_i + \varepsilon \\ \text{M3: } \mu(t, p) &= \beta_0(p) + \beta_1(t, p) \text{ SPEI}_i + \varepsilon \\ \text{M4: } \mu(t, p) &= \beta_0(t, p) + \beta_1(t, p) \text{ SPEI}_i + \varepsilon \end{aligned} \quad (8)$$

where $\mu(t, p)$ equals $\partial\text{O}^{18}(t, p)$ or WUEi(t, p), respectively, and the error term ε follows a Normal distribution.

3. Results

3.1. Thinning effects on radial growth

The BAI series showed very similar patterns for all treatment plots before thinning as indicated by the non-significant differences (Fig. 2 and Table 2). As expected, lower values of BAI were recorded during drought years (e.g. 1994–1995, 2005–2006, 2012), while high BAI values were observed during wet years such as 1997 (Fig. S2). The lowest values of BAI occurred in 1994 in Tarazona (0.72 cm^2) and in 2005 in La Hundo (0.80 cm^2).

Differences between treatments were apparent after the thinning in both sites. BAI was enhanced in both thinning intensities plots, although the greatest increases occurred on the high intensity thinning plots (H). On the other hand, the moderate intensity treatment (M) in Tarazona showed increasing BAI annual values in the *postT10* period, while in the last period the differences with the control disappeared, decreasing with the time elapsed since thinning (Fig. 2). In La Hundo, differences were significant compared with the *postT* period. On the contrary, in the control plots the differences between the periods were negligible.

3.2. Combined effects of drought and thinning on radial growth

In both sites the highest correlation between BAI and SPEI time series in control plots was found during summer (see Fig. S1 in the Supporting Information). The SPEI at the time scale of 9 months and week 31 of the year (mid-August) had the highest correlation in Tarazona, with a positive and significant Pearson correlation coefficient of 0.75, while in La Hundo the highest correlation was found in week 24 (mid-June) at the same time scale of 9 months, with a positive and significant correlation coefficient of 0.55 (Fig. S3). SPEI data for those weeks and time scales were therefore used in the regression analysis as independent variables.

The results of the linear regression analysis confirmed that BAI was significantly affected by the SPEI and the thinning treatments (Table S1, Fig. 3). It is interesting to compare the model coefficients between *preT* and *postT* periods for a given treatment, while the comparison between the two sampling sites confirms the robustness of the results.

Between the different model configurations tested, the lowest AIC was obtained by model M4 with full interaction between the intercept and the slope coefficients and the two factors (period and treatment). The relationship between BAI and SPEI was positive in all cases, so high values of the SPEI, indicative of a wetter-than-normal year, were associated with high values of BAI and vice versa (Fig. 3a). The regression coefficients for SPEI were in general higher in Tarazona than in La Hundo, indicating a stronger influence of climate on BAI. In the *postT* period the coefficients were lower in the thinned plots than in the control, although this difference was only significant in La Hundo between control and heavy thinning. While the coefficients were not significantly different between both periods in the control plots, they were significantly different in the thinned plots, in both sites.

There were differences in the model intercepts, too, indicative of differences in mean BAI for controlled SPEI conditions (since a value of $\text{SPEI} = 0$ represents average water balance). The coefficients were significantly higher for thinned plots in the *postT* period, providing evidence that BAI increased on average after thinning (Fig. 3a). This increase was moderately higher (albeit not significantly so) in the heavy thinned than in the moderately thinned plots. Thinning increased BAI by around 0.50 to 0.75 cm^2 in moderately and heavy thinned plots in Tarazona, and by 0.75 and 1.25 cm^2 in La Hundo. For comparison of the magnitude of the effects, this is a similar difference as between moderately favourable ($\text{SPEI} = 0.5$) and unfavourable ($\text{SPEI} = -0.5$) years in the control plot.

Fig. 3b shows the resulting GLM models of BAI against SPEI, only for the post period. The higher sensitivity of BAI to SPEI in Tarazona is reflected by the steeper lines, while differences in BAI between




Fig. 2. Basal area increment (BAI) series of *Pinus halepensis* plantations in Tarazona and La Hunde for control plots (squares), moderately (grey triangles) and heavily thinned (black circles) plots. Bars represent standard errors. Vertical dashed lines indicate the date of the thinning interventions (1998 in Tarazona and 2008 in La Hunde). Significant differences in mean BAI values among treatments before thinning (*preT*) and after thinning (*postT*) are indicated with letters (p -value < 0.05) in the right-hand plots.

treatment levels are also apparent. The regression line for the heavy thinned plot in La Hunde is slightly negative, but the coefficient's confidence interval did not exclude zero so the line can be considered as flat, indicating that a negligible influence of SPEI on BAI existed.

Inspection of the random effects revealed that there were remarkable differences between individuals, with extreme values ranging between -0.2 and 0.2 cm^2 . In general, thinning did not seem to modify significantly the variability between trees in the *postT* period.

3.3. Resistance, recovery and resilience indices

Drought resistance (i.e., the ratio between BAI values of the drought year and the two previous years, R_t) tended to be, as expected, lower

than one, indicating reduced growth during a drought. There were exceptions to this, however, such as in the 1995 episode in Tarazona where values were unexpectedly higher than one. In general, R_t was increased by thinning, but only significantly so in La Hunde (Fig. 4a, b). Differences were also found during the *preT* period in both sites, with thinning plots having lower R_t in general (but only significantly so in one drought episode on each site).

Drought recovery (i.e. the ratio between BAI values posterior to and during the drought, R_c) was higher than one in all cases, indicating a good ability to recover to the pre-drought situation in general. Values between 2.5 and 5 were normal, although in some episodes (2006 in Tarazona, 2000 in La Hunde) R_c was almost one implying very limited growth recovery. No significant differences were found between

Table 2

Basal Area Increment (BAI), intrinsic water-use efficiency (WUEi) and oxygen isotope ratio ($\delta^{18}\text{O}$) values calculated for the two study sites and treatments (*preT* and *postT* refer to the pre- and post-thinning periods, *postT10* and *postT20* in Tarazona). Values are means $\pm$ SE.

Thinning intensity and period*	Tarazona			La Hundé	
	<i>preT</i> (1965–1997)	<i>postT10</i> (1998–2007)	<i>postT20</i> (2008–2017)	<i>preT</i> (1965–2007)	<i>postT</i> (2008–2017)
BAI (cm^2)					
Control	3.51 $\pm$ 0.30	2.77 $\pm$ 0.29a	3.50 $\pm$ 0.47a	3.45 $\pm$ 0.19	3.24 $\pm$ 0.26a
Moderate	3.26 $\pm$ 0.26	4.97 $\pm$ 0.59b	4.91 $\pm$ 0.48a	4.00 $\pm$ 0.23	7.45 $\pm$ 0.55b
Heavy	2.60 $\pm$ 0.26	5.52 $\pm$ 0.73b	6.93 $\pm$ 0.61b	4.16 $\pm$ 0.26	9.87 $\pm$ 0.10c
WUEi ($\mu\text{mol mol}^{-1}$)					
Control	109.10 $\pm$ 3.04	107.84 $\pm$ 1.96	107.74 $\pm$ 2.29	102.36 $\pm$ 1.56b	102.83 $\pm$ 2.19b
Thinned	106.80 $\pm$ 2.08	103.49 $\pm$ 2.17	101.45 $\pm$ 1.94	95.25 $\pm$ 1.05a	95.59 $\pm$ 1.95a
$\delta^{18}\text{O}$ (‰)					
Control	27.47 $\pm$ 0.11b	27.45 $\pm$ 0.10	26.93 $\pm$ 0.16a	27.86 $\pm$ 0.12b	27.80 $\pm$ 0.12a
Thinned	26.58 $\pm$ 0.26a	27.73 $\pm$ 0.15	27.57 $\pm$ 0.13b	27.44 $\pm$ 0.14a	28.41 $\pm$ 0.25b

* Letters (a, b, c) indicate significant differences between treatments at the 95% confidence level (p -value $<$ 0.05), according to ANOVA using post hoc Tukey test for multiple comparisons.

treatments after thinning (*postT*), although the values were consistently lower on thinned plots suggesting reduced recovery (Fig. 4c, d). This interpretation is reinforced by the fact that, before thinning (*preT* period), R_c was higher in the thinning plots, especially in Tarazona.

Regarding the resilience index (i.e. the ratio between BAI values posterior to and before the drought. R_s), values higher than one were the norm although there were exceptions in some droughts and plots. Significant differences between treatments were found after eight and four years in Tarazona and La Hundé, respectively, with lower resilience corresponding to thinned plots (Fig. 4e, f). In Tarazona, again, differences between plots were also found before thinning (*preT* period), although the variability between trees was very high and these values must be considered with care.

3.4. Combined effects of thinning and SPEI on WUEi and $\delta^{18}\text{O}$

Analysis of the time series of WUEi showed lower average values on thinned plots even during the *preT* period (Fig. 5 and Table 2), most notably in La Hundé. This means that there were differences between the plots (control and thinned) before the treatment took place, so the most valuable comparison has to be between *preT* and *postT* periods. In Tarazona WUEi decreased from 106.8 to 103.49 $\mu\text{mol mol}^{-1}$ in *postT10* and to 101.45 $\mu\text{mol mol}^{-1}$ *postT20* for thinned trees, while in La Hundé the differences were lower. At the same time, WUEi on the control plot remained remarkably constant during the two periods. In all cases the WUEi increased during dry years in both sites (e.g. 1994–1995, 2005–2006, 2014), especially in the control plots.

The best linear model configuration according to the AIC statistic corresponded to model M4, with full interaction between the intercept and the slope and the treatment and period factors. The SPEI had a negative effect on WUEi, indicating increased water use efficiency during

dry periods (Fig. 6a and Table S3). There were differences in the SPEI model coefficients between treatments both during the *preT* and *postT* periods in both sites, although their confidence ranges highly overlapped. Differences in the SPEI coefficients between *preT* and *postT* periods were only significant in the thinned plots, and especially so in the case of Tarazona, with more negative values in the latter period suggesting that thinning increased the sensitivity of WUEi to the SPEI. The model intercepts were also different between treatments both during the *preT* and *postT* periods, and especially so in La Hundé. Only in Tarazona the intercept was around 4 $\mu\text{mol mol}^{-1}$ lower in the *postT* period with respect to the *preT* period in the thinned plot, while in the control plot, as well as in La Hundé for both plots, the intercepts remained fairly constant during the two periods.

The relationship between WUEi and SPEI during the *postT* period is more clearly seen in Fig. 6b. The range of variation of WUEi with SPEI was similar in both sites, ranging roughly between 115 and 90 mmol mol^{-1} for SPEI values of -2 (very dry conditions) and 2 (very wet conditions), respectively. WUEi was consistently lower in the thinned plots in both sites, and only in Tarazona the slope of the model was slightly different, so during droughts (negative SPEI) the difference in WUEi was small but increased rapidly as the conditions improved. This suggests a more elastic response of thinned trees to the climatic conditions, at least on this site, although as it was mentioned before this difference was not significant since the confidence bands of the coefficients slightly overlapped.

With respect to $\delta^{18}\text{O}$, there were also differences between control and thinned plots even before the thinning treatment (*preT* period), with lower values in the thinned plots in both sites (Fig. 7 and Table 2). However, in both sites $\delta^{18}\text{O}$ increased in the thinning plots after the thinning treatment, while in the control plots it remained constant or even decreased (Tarazona). In Tarazona $\delta^{18}\text{O}$ was significantly




Fig. 3. Model coefficients for the fixed effects (mean values and 95% confidence intervals) for basal area increment (BAI) in Tarazona and La Hundé, for three treatment levels during pre-thinning (*preT*) and post-thinning (*postT*) periods (a), and generalized linear models of BAI against SPEI for different treatment levels in the *postT* period (b).




Fig. 4. Temporal evolution of tree resistance (a, b), recovery (c, d) and resilience (e, f) indices in response to selected droughts calculated for Tarazona (left panels) and La Hundo (right panels) growth data (BAI) considering control plots (C), moderately (M) and heavily thinned (H) plots. Values are shown before (circles, Pre) and after thinning (squares, Post) treatments. Vertical dashed lines indicate the date of thinning treatments. Bars represent standard deviations. Significant differences among thinning treatments are indicated with letters (p -value < 0.05).

higher ($27.57 \pm 0.13\%$, p -value < 0.05) for thinned trees in the *postT20* period and was especially responsive during dry periods (2005–2006 and 2012). In La Hundo it was also significantly higher ($28.41 \pm 0.25\%$, p -value < 0.05) for thinned trees, and remarkably higher during the dry year 2012.

Regression analyses revealed a negative relationship between $\delta^{18}\text{O}$ and SPEI, implying that $\delta^{18}\text{O}$ decreased during favourable years (positive SPEI) and increased during droughts (Table S3, Fig. 8). The exception to that was Tarazona during the *preT* period, which had a positive coefficient. There were marked differences between sites in the evolution between *preT* and *postT* periods: while in Tarazona the SPEI coefficient experienced a decrease, in La Hundo the opposite was found. No significant differences were found between thinned and control plots in the slope of the regression, and in fact model M1, with no interaction between β_1 and treatment, was selected by the AIC statistic as the best configuration. Once the effect of the climatic conditions (SPEI) were accounted for we found significant evidence of increasing $\delta^{18}\text{O}$ values after thinning, as revealed by the model intercepts. Thus, while in the control plots the model intercepts decreased in the period after thinning (only significantly in Tarazona), the thinned trees experienced a strong and significant increase in both sites. The difference between control and thinned trees was about 0.5‰ in both sites, as shown in Fig. 8b.

We analysed the changes in isotopic concentrations after thinning according to the dual isotope interpretation of Scheidegger et al. (2000). Fig. 9 shows annual and mean differences in $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ with respect to the control plot ($\Delta\delta^{13}\text{C}$ and $\Delta\delta^{18}\text{O}$, respectively) after thinning. The predominant signal was a positive movement in the

$\Delta\delta^{18}\text{O}$ axis, while in the $\Delta\delta^{13}\text{C}$ there was larger variability between years and the average difference was almost zero. There was no relationship between dry/wet years and the positive or negative sign of $\Delta\delta^{13}\text{C}$, and the only effect seen at the annual level was a tendency of the year or years after the thinning to have null or negative values of $\Delta\delta^{18}\text{O}$, as it was the case of 1998 in Tarazona or 2008 and 2009 in La Hundo.

4. Discussion

4.1. Thinning effects on radial growth

Thinning enhanced radial growth at both experimental sites, and slightly more so in the heavy thinning plot than in the moderately thinned plot. Reduction of intra-species competition as induced by thinning clearly benefited the remaining trees, a result that can be explained by the fact that resource availability (water, nutrients, light) increases with the growing space (Breda et al., 1995; Aussenac, 2000). The positive effect of thinning on BAI has already been reported for *P. halepensis* in semiarid conditions and similar thinning intensity reductions of basal area by Martín-Benito et al. (2010), del Campo et al. (2014), Fernandes et al. (2016) and Tsamir et al. (2019), where the increase of BAI represented a mean annual increment of $7.01 \text{ cm}^2 \text{ year}^{-1}$ for heavy thinning treatments and $2.85 \text{ cm}^2 \text{ year}^{-1}$ for moderate thinning treatments. An increase of growth after thinning was also observed in other planted pine species such as *P. nigra* (Navarro-Cerrillo et al.,




Fig. 5. Time series of the intrinsic water-use efficiency (WUEi) at Tarazona and La Hundo according to thinning treatments: control (empty triangles) and heavy thinning (filled dots). In both cases, $n = 5$ trees per treatment. Vertical dashed lines indicate the date of thinning treatments (1998 in Tarazona and 2008 in La Hundo). Right margin plots show mean and standard errors for both treatments during pre- (*preT*) and post-thinning (*postT*) periods, with significant (p -value < 0.05) differences among treatments during each period indicated with letters.

2019), *P. sylvestris* (Giuggiola et al., 2016; Navarro-Cerrillo et al., 2019) and *Pinus ponderosa* (McDowell et al., 2003; McDowell et al., 2006).

It is crucial to examine the positive effect of thinning in long-term experiments to assess the persistence of the effects of thinning with time elapsed since the intervention (Bottero et al., 2017). We found that the effect of thinning enhanced growth even 20 years after the treatment in the heavy thinning treatment in Tarazona, while in the

moderate treatment the effects started to fade (the growth difference with respect to control was not significant, albeit it was still positive). This persistent effect has already been reported for *Pinus contorta* where the enhanced growth was sustained after 18 years (Prescott et al., 2019), and for *P. pinaster* throughout all the study period (15 years) (Navarro-Cerrillo et al., 2016). The maintenance of benefits of thinning are common for short-term trials and have been described




Fig. 6. Model coefficients for the fixed effects (mean values and 95% confidence intervals) for intrinsic water use efficiency (WUEi) in Tarazona and La Hundo, for two treatment levels during pre- (*preT*) and post-thinning (*postT*) periods (a), and generalized linear models of WUEi against SPEI for control and heavy thinning treatment levels in the *postT* period (b).




Fig. 7. Time series of $\delta^{18}\text{O}$ values at Tarazona and La Hundo according to thinning treatments: control (empty triangles) and heavy thinning (filled dots). In both cases, $n = 5$ trees per treatment. Vertical dashed lines indicate the date of thinning treatments, and margin plots show the mean and standard deviation for each treatment during pre- (*preT*) and post-thinning (*postT*) periods, with significant ($p\text{-value} < 0.05$) differences among treatments during each period indicated with letters.

from 5 to 10 years and in several pine species, such as *P. halepensis* (Ruano et al., 2013; Tsamir et al., 2019), *P. sylvestris* and *P. nigra* (Navarro-Cerrillo et al., 2019) and *P. ponderosa* and *P. resinosa* (Bottero et al., 2017).

As a novelty with respect to previous studies, we also considered the effect of climate (SPEI drought index) on growth (BAI) under the different treatments. Growth was reduced during dry years (negative SPEI

values) as expected, but we also found a lower sensitivity of BAI to SPEI after thinning, with regression coefficients closer to zero (and more so in the heavy thinning than in the moderate thinning). This can be interpreted as the climate affecting less in the thinned plots where soil water availability increases. Lower climate sensitivity implies reduced stress during non-favourable periods (droughts), which have important positive consequences for forest vitality and health. We also




Fig. 8. Model coefficients for the fixed effects (mean values and 95% confidence intervals) for $\delta^{18}\text{O}$ in Tarazona and La Hundo, for two treatment levels during pre- (*preT*) and post-thinning (*postT*) periods (a), and generalized linear models of $\delta^{18}\text{O}$ against SPEI for control and heavy thinning treatment levels in the *postT* period (b).




Fig. 9. $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ differences from the control (thinned - control). White squares represent the mean values of $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ during the *preT* period. Grey arrows represent the deviations with respect to these means for all the years in the *postT* period, and the black arrow represents the mean value of the *postT* period. Thinning years (*italics*).

found differences between sites, as the SPEI had a stronger effect in Tarazona than in La Hundé, suggesting that drought represents a harder constraint in the former site. Despite the two sites having similar Mediterranean climatic conditions (Fig. S2), there are other environmental differences that may help to explain why the trees were more climate-sensitive in one place than in the other. The growing season is on average shorter in Tarazona due to lower winter and spring temperatures. Additionally, growth rates may also be affected by other factors such as different crown classes (Martín-Benito et al., 2008), nutrient limitation, plant acclimation to microclimate conditions or reallocation of carbohydrates after thinning (Peñuelas et al., 2011), which could differ between sites.

Several studies have described climate-growth relationships on Mediterranean pine species (De Luis et al., 2011; Camarero et al., 2013). A strong relationship between BAI and spring-summer precipitation has been reported frequently in water-limited Mediterranean ecosystems (Martín-Benito et al., 2010; Fernandes et al., 2016; Navarro-Cerrillo et al., 2019). There are fewer studies, however, that included the effects of climate and management together. Martín-Benito et al. (2010) found in *P. nigra* that, after thinning, tree growth had a lower correlation with precipitation than before thinning, i.e. lower sensitivity, while trees in their control plot did not show any difference, with the correlation even increasing after thinning. Sohn et al. (2016) and Giuggiola et al. (2016) suggested a weaker growth-climate relationship as thinning intensity increases, at least for *P. sylvestris*. In the case of *P. halepensis*, Fernandes et al. (2016) found in La Hundé experimental site that thinning made trees less sensitive to soil water shortage, showing that trees in the unthinned plots needed to rely more heavily on current year precipitation than those in thinned plots. Here we used the SPEI instead of only precipitation data, since it provides a more reliable measure of drought severity by considering also the temperature-driven evaporative demand by the atmosphere (Vicente-Serrano et al., 2010; Beguería et al., 2014), and we also found evidence of a reduced sensitivity to climate induced by thinning.

4.2. Thinning effects on water-use efficiency (WUEi)

We found differences in WUEi between control and thinned plots before the thinning took place. Despite this, WUEi was significantly

reduced after thinning, while it remained fairly constant in the control plots when comparing the pre- and post-thinning periods. This result was consistent across sites. The reduction of WUEi on thinned stands coincides with the results of Navarro-Cerrillo et al. (2019) for *P. nigra* and *P. sylvestris* and Fernandes et al. (2016) for *P. halepensis*, depicted the last work a reduction from $94.4 \mu\text{mol mol}^{-1}$ to $88.7 \mu\text{mol mol}^{-1}$ over 10 years after thinning. McDowell et al. (2003) also found a reduction from $112 \mu\text{mol mol}^{-1}$ to $97 \mu\text{mol mol}^{-1}$ over 6 years after thinning in *P. ponderosa*. Here we found a reduction between 5 and $10 \mu\text{mol mol}^{-1}$ for average climatic conditions (SPEI = 0) in Tarazona and La Hundé, respectively, which is slightly lower. Interestingly, we found that the effect of thinning persisted in Tarazona even 20 years after the treatment.

The decrease in WUEi and the increase of BAI in thinned stands indicate that water deficit due to competition limited tree growth on these plantations (Moreno-Gutiérrez et al., 2012). This interpretation is also supported when analyzing the effect of climate, since WUEi increased during droughts as the trees are affected by water shortage (Moreno-Gutiérrez et al., 2012) and it decreased during wet years. In terms of the regression model, WUEi had a negative relationship with the SPEI implying that lower water use efficiency is related to a reduction in water stress (Klein et al., 2005). In one site (Tarazona), we found that the sensitivity of WUEi to SPEI tended to become higher (i.e., more negative regression coefficient) in the thinned plot, indicating a more elastic response of trees to the climatic conditions. However, this difference was only marginally significant, and this effect was not found in La Hundé, where the regression coefficients were not significantly different between control and thinned plots.

Thinning effects on isotopic composition and physiological interpretation.

Our finding of increasing $\delta^{18}\text{O}$ and constant $\delta^{13}\text{C}$ after thinning in both sites coincides with the results reported by Fernandes et al. (2016) for *P. halepensis*, Martín-Benito et al. (2010) for *P. nigra*, and Brooks and Mitchell (2011) for *Pseudotsuga menziesii*. Giuggiola et al. (2016) found increasing $\delta^{18}\text{O}$ but decreasing $\delta^{13}\text{C}$ after thinning for *P. sylvestris*, and other combinations including constant (Powers et al., 2010) or even decreasing $\delta^{18}\text{O}$ have also been reported by McDowell et al. (2006) for *P. ponderosa* and Moreno-Gutiérrez et al. (2011) for *P. halepensis*.

According to the dual-isotope theoretical model of Scheidegger et al. (2000) (Fig. 9), increasing $\delta^{18}\text{O}$ and constant $\delta^{13}\text{C}$ after thinning should be interpreted as the result of a simultaneous reduction of stomatal conductance (g_s) and photosynthetic capacity (A). This interpretation, however, would contradict our findings of enhanced BAI and reduced WUEi after thinning. As it has been pointed out by Roden and Siegwolf (2012), this model has some limitations and should only be applied under certain conditions (Barnard et al., 2012). Variability of $\delta^{18}\text{O}$ in plants includes the combined effects of the isotopic composition of the soil water; leaf water enrichment due to transpiration; and biochemical fractionation during the synthesis of organic matter (Scheidegger et al., 2000; Gessler et al., 2014). As their authors point out, the model is based on the assumptions of constant isotopic composition of soil water and atmospheric vapour. We shall examine the validity of these assumptions in our specific case, and in general in the case of thinning treatments.

Especially in dry environments, it is very likely that thinning induces a hotter micro-climate with increased turbulence and air interchange at the stand level, leading to reduced air humidity (Martin-Benito et al., 2010). Therefore, $\delta^{18}\text{O}$ enrichment could be observed that would not be related to changes in stomatal conductance. This mechanism has also been suggested by other authors (Martin-Benito et al., 2010; Fernandes et al., 2016).

Similarly, it could be argued that thinning can also induce changes in soil water uptake depths, again especially in dry environments. Evaporation from the soil causes oxygen and deuterium enrichment near the soil surface, which generally decreases exponentially with soil depth (Barnes and Allison, 1988; Ehleringer and Dawson, 1992). Several authors reported a reduction in shallow water uptake with increasing water deficit, and shallower water uptake during relative wet conditions (Sarris et al., 2013; Grossiord et al., 2017; Barbeta and Penuelas, 2017). We hypothesize that trees on thinned stands present higher rates of shallow water uptake than in unthinned stands, at least under normal conditions. On the contrary in unthinned stands, due to enhanced competition, water is extracted more often from deeper soil layers. This would agree with our observation that thinned trees present weaker growth-climate relationship, while control trees are more restrained by the climatic conditions. This mechanism has also been suggested by Sarris et al. (2013) for thinned *P. halepensis* subsp. *brutia*.

It is clear that tree-ring $\delta^{18}\text{O}$ may not always be a good proxy of foliar conditions, and as a consequence should not be used alone to study stomatal behaviour, at least under uncontrolled conditions regarding the source of water or the microclimatic conditions at the stand level. Our findings suggest that $\delta^{18}\text{O}$ was neither constant in the soil water nor in the atmosphere after and before thinning, so that the observed increase in tree-ring $\delta^{18}\text{O}$ does not necessarily imply a reduction of stomatal conductance and photosynthetic capacity as implied by the Scheidegger et al. (2000) model. Araguas-Araguas and Diaz Teijeiro (2005) reported spatial and temporal variability both in rainfall and in soil water in the Iberian Peninsula. Previous results from La Hundo by Molina and del Campo (2012) and del Campo et al. (2014) show that thinning increased the amount of water reaching the soil surface by reducing rainfall interception, but also increased evaporation from the soil.

It is difficult to offer a physiological interpretation of the observed changes in thinned stands in our case study, at least based on isotopic data alone. We observed increasing BAI in thinned trees, which could be explained by increased carbon assimilation. We also observed a reduction in WUEi, which could be a consequence of: i) decreasing A at constant g_s ; ii) increasing g_s at constant A ; or iii) increasing g_s and A , with that in g_s proportionally larger than that in A . Therefore, our proposed mechanism for explaining increasing BAI and reduced WUEi is an increase in both g_s and A . The increase in g_s was proportionally larger than the increase in A which could explain the reduction of WUEi. This interpretation agrees with sap flow measurements in La Hundo, which revealed increasing transpiration on thinned trees (del Campo et al.,

2014, and Fernandes et al., 2016). The increase in A could be related to elevated soil water and nitrogen availability (Feeney et al., 1998; Kaye and Hart, 1998; McDowell et al., 2003; McDowell et al., 2006) and improved nutrient foliar status such as increased N content after thinning (López-Serrano et al., 2005), that generally results in a greater net CO_2 assimilation rate during photosynthesis and increased growth (Warren et al., 2001; Moreno-Gutiérrez et al., 2011). In addition, an increase in A may also occur as a result of a water-availability related increase in g_s , thus allowing for a higher CO_2 uptake (Giuggiola et al., 2016; Moreno-Gutiérrez et al., 2011).

4.3. Reliability of resistance, recovery and resilience indices

Analysis of basal growth inferred from tree-ring series revealed the benefits of thinning, and the analysis of WUEi and stable isotopes allowed interpreting the mechanisms that explain these benefits. It could be expected, therefore, that thinned trees would exhibit enhanced resistance and resilience to drought. Thinning seemed to enhance drought resistance, implying a reduced effect of drought on tree growth, what coincides with the results of regression analysis that indicated reduced climate sensitivity of thinned trees (lower regression coefficients for SPEI).

However, thinning seemed to reduce drought recovery and resilience indices, which seems counter-intuitive. In the case of drought recovery index, we are comparing tree growth after the drought and during the drought. Therefore, if drought affects thinned trees less, then it is to be expected that the recovery index would also be lower. A similar reasoning can be made with respect to drought resilience, where the years after and before the drought are compared.

One problem with drought resistance indices is that the climate conditions of the years before and after a drought are not considered, nor the magnitude of the drought. Climate is an uncontrolled factor in observational studies, and therefore it is difficult to determine how the particular conditions of the years used may affect the indices. In short, it is very different if the year before or after the drought were average years or very good in terms of moisture availability. One benefit of the regression analysis performed is that it allows considering the effect of climate on BAI, through the SPEI drought index. This allows us to redefine at least resistance and recovery indices based on the regression models, using the following expressions:

$$\text{Resistance (Rt)} = \text{BAI}_{\text{SPEI}=-2} / \text{BAI}_{\text{SPEI}=0} \quad (9)$$

$$\text{Recovery (Rc)} = \text{BAI}_{\text{SPEI}=0} / \text{BAI}_{\text{SPEI}=-2} \quad (10)$$

where $\text{BAI}_{\text{SPEI}=-2}$ is the tree growth expected for a value of the SPEI of -2 (that is, a severe drought), and $\text{BAI}_{\text{SPEI}=0}$ is the tree growth expected during normal conditions. Following this approach, we obtain values of R_t of 0.62 and 0.91 for moderately thinned trees in Tarazona and La Hundo, respectively, against 0.50 and 0.75 in the control plot. That is, thinning enhanced drought resistance in both sites following this approach. Regarding R_c , we obtain values of 1.62 and 1.10 in the thinned plot against 2.0 and 1.32 in the control plot, confirming that thinning reduced drought recovery according to the formulation of this index. The resilience index cannot be defined in a similar way, since it would involve normal conditions in the nominator and denominator.

5. Conclusions

The findings reported here contribute to the existing knowledge about the effects of thinning on tree growth and functioning, with the general conclusion that thinning had a positive effect on the two Aleppo pine stands analysed by enhancing radial growth. The most interesting result in terms of adaptive silviculture was that thinning helped to reduce growth dependence on drought (SPEI) on the remaining trees. In addition, we found that thinning decreased WUEi, implying a reduction

of water stress. Considering the climate and drought (SPEI) effects on BAI to calculate drought resistance index, we observed an increase after thinning that coincided with the observations on selected dry years. Results regarding recovery and resilience indices were compromised by the fact that growth was less affected by drought on thinned trees, so both indices had reduced values when compared to trees in the control plot. We cannot deduce, however, that recovery and resilience were lower on thinned trees, since the effect of drought (and therefore vulnerability) was clearly lower. Due to this effect, we do not recommend the use of these two indices for comparing between thinned and not thinned stands, or in general when different drought vulnerability between groups is suspected. Here we selected two sites of similar climatic conditions to increase robustness of the study, these sites are representative of semiarid conditions, although *P. halepensis* covers a large range of climatic conditions.

Further research should be undertaken to determine to what extent stable isotopes are a reliable tool to describe physiological changes after thinning under uncontrolled environmental conditions. We found that the dual-isotope theoretical model cannot be applied to our case due to non-controlled factors that affect wood $\delta^{18}\text{O}$ composition besides stomatal conductance, such as changes in water uptake depth (and therefore isotopic composition) or micro-climatic effects derived from the thinning. Based on additional physiologic measurements undertaken in one of the two sites, we hypothesize that thinning promoted an increase of both stomatal conductance (transpiration) and photosynthetic rates, with the former being proportionally larger so their ratio ($\text{WUEi} = A/g_s$) was reduced.

As a policy recommendation, our findings suggest that thinning (25 to 50% of the initial basal area removed) offers a clear advantage and can be implemented to adapt Mediterranean pine forests to climate variability, improving resistance against drought and promoting a lower dependence of growth on climate conditions. The positive effects of thinning remained at least for 20 years, especially in the heavy thinning treatment, and this value could be used a starting point in order to plan further interventions. This could be critically important to develop adaptation management plans to respond proactively to climate-related global challenges.

CRediT authorship contribution statement

Àngela Manrique-Alba: Conceptualization, Investigation, Writing - original draft. **Santiago Beguería:** Methodology, Formal analysis, Writing - review & editing, Supervision, Funding acquisition. **Antonio J. Molina:** Validation, Writing - review & editing. **María González-Sanchis:** Writing - review & editing. **Miquel Tomàs-Burguera:** Data curation, Writing - review & editing. **Antonio D. del Campo:** Writing - review & editing. **Michele Colangelo:** Resources. **J. Julio Camarero:** Conceptualization, Methodology, Writing - review & editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

Funding for this research was provided by the Spanish “Agencia Estatal de Investigación” and the European Commission in the frame of the ERA-NET WaterWorks2015 Cofunded Call (project INNOMED). This ERA-NET is an integral part of the 2016 Joint Activities developed by the Water Challenges for a Changing World Joint Programme Initiative (Water JPI) as a result of a joint collaborative effort with the Joint Programming Initiative on Agriculture, Food Security and Climate Change (FACCE JPI). A.J. Molina is beneficiary of an ‘APOSTD’ fellowship (APOSTD/2019/111) funded by the Generalitat Valenciana. JJC

acknowledges the support of FunDiver (CGL2015-69186-C2-1-R) and FORMAL (RTI2018-096884-B-C31) projects from the Spanish Ministry of Science, Innovation and Universities. The support of CEHYRFO-MED (CGL2017-86839-C3-2-R), CLICES (CGL2017-83866-C3-3-R), RESILIENT-FORESTS (LIFE17 CCA/ES/000063) and SILVADAPT.NET (RED2018-102719-T) projects is also acknowledged. We sincerely thank E. Arrechea and Alvaro Hernández (Gob. Aragón) for their support and help to study the Tarazona plantation site, and A. Sánchez-Miranda for helping processing the tree-ring samples.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2020.138536>.

References

- Allen, C.D., Macalady, A.K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Kitzberger, T., Rigling, A., Breshears, D.D., Hogg, E.H., Gonzalez, P., Fensham, R., Zhang, Z., Castro, J., Demidova, N., Lim, J.-H., Allard, G., Running, S.W., Semerci, A., Cobb, N., 2010. A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecol. Manag.* 259, 660–684.
- Anderegg, W.R.L., Schwalm, C., Biondi, F., Camarero, J.J., Koch, G., Litvak, M., Ogle, K., Shaw, J.D., Shevliakova, E., Williams, A.P., Wolf, A., Ziaco, E., Pacala, S., 2015. Pervasive drought legacies in forest ecosystems and their implications for carbon cycle models. *Science* 349, 528–532.
- Araguas-Araguas, L.J., Diaz Teijeiro, M.F., 2005. Isotope composition of precipitation and water vapour in the Iberian Peninsula. *Isotopic Composition of Precipitation in the Mediterranean Basin in Relation to Air Circulation Patterns and Climate*. IAEA-TECDOC-1453 (October 2005).
- Aussenac, G., 2000. Interactions between forest stands and microclimate: ecophysiological aspects and consequences for silviculture. *Ann. Sci.* 57, 287–301.
- Barbour, M.M., 2007. Stable oxygen isotope composition of plant tissue: a review. *Funct. Plant Biol.* 34, 83–94.
- Barnard, H.R., Brooks, J.R., Bond, B.J., 2012. Applying the dual-isotope conceptual model to interpret physiological trends under uncontrolled conditions. *Tree Physiol.* 32, 1183–1198.
- Barnes, C.J., Allison, G.B., 1988. Tracing of water movement in the unsaturated zone using stable isotopes of hydrogen and oxygen. *J. Hydrol.* 100, 143–176.
- Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* 67, 1–48.
- Battipaglia, G., De Micco, V., Fournier, T., Aronne, G., Carcaillet, C., 2014. Isotopic and anatomical signals for interpreting fire-related responses in *Pinus halepensis*. *Trees-Struct. Funct.* 28, 1095–1104.
- Barbeta, A., Penuelas, J., 2017. Increasing carbon discrimination rates and depth of water uptake favor the growth of Mediterranean evergreen trees in the ecotone with temperate deciduous forests. *Glob. Change Biol.* 23 (12), 5054–5068.
- Beguería, S., Vicente-Serrano, S.M., Reig, F., Latorre, B., 2014. Standardized precipitation evapotranspiration index (SPEI) revisited: parameter fitting, evapotranspiration models, tools, datasets and drought monitoring. *Int. J. Climatol.* 34, 3001–3023.
- Bottero, A., D’Amato, A.W., Palik, B.J., Bradford, J.B., Fraver, S., Battaglia, M.A., Asherin, L.A., 2017. Density-dependent vulnerability of forest ecosystems to drought. *J. Appl. Ecol.* 54 (6), 1605–1614.
- Breda, N., Granier, A., Aussenac, G., 1995. Effects of thinning on soil and tree water relations, transpiration and growth in an oak forest (*Quercus petraea* (Matt.) Liebl.). *Tree Physiol.* 15, 295–306.
- Brooks, J.R., Mitchell, A.K., 2011. Interpreting tree responses to thinning and fertilization using tree-ring stable isotopes. *New Phytol.* 190, 770–782.
- Camarero, J.J., Manzanedo, R.D., Sanchez-Salguero, R., Navarro-Cerrillo, R.M., 2013. Growth response to climate and drought change along an aridity gradient in the southernmost *Pinus nigra* relict forests. *Ann. For. Sci.* 70, 769–780.
- Camarero, J.J., Gazol, A., Sangüesa-Barreda, G., Oliva, J., Vicente-Serrano, S.M., 2015. To die or not to die: early warnings of tree dieback in response to a severe drought. *J. Ecol.* 103, 44–57.
- Camarero, J.J., Sanchez-Salguero, R., Sangüesa-Barreda, G., Matías, L., 2018. Tree species from contrasting hydrological niches show divergent growth and water-use efficiency. *Dendrochronologia* 52, 87–95.
- del Campo, A.D., Fernandes, T.J., Molina, A.J., 2014. Hydrology-oriented (adaptive) silviculture in a semiarid pine plantation: how much can be modified the water cycle through forest management? *Eur. J. For. Res.* 133, 879–894.
- del Campo, A.D., González-Sanchis, M., Lidón, A., Ceacero, C.J., García-Prats, A., 2018. Rainfall partitioning after thinning in two low-biomass semiarid forests: impact of meteorological variables and forest structure on the effectiveness of water-oriented treatments. *J. Hydrol.* 565, 74–86.
- del Campo, A.D., González-Sanchis, M., Molina, A.J., García-Prats, A., Ceacero, C.J., Bautista, I., 2019. Effectiveness of water-oriented thinning in two semiarid forests: the redistribution of increased net rainfall into soil water, drainage and runoff. *Forest Ecol. Manag.* 438, 163–175.
- Carnicer, J., Coll, M., Ninyerola, M., Pons, X., Sánchez, G., Peñuelas, J., 2011. Widespread crown condition decline, food web disruption, and amplified tree mortality with increased climate change-type drought. *Proc. Natl. Acad. Sci. U. S. A.* 108, 1474–1478.

- De Luis, M., Novak, K., Raventos, J., Grisar, J., Prislán, P., Čufar, K., 2011. Climate factors promoting intra-annual density fluctuations in Aleppo pine (*Pinus halepensis*) from semiarid sites. *Dendrochronologia* 29, 163–169.
- Dobbertin, M., 2005. Tree growth as indicator of tree vitality and of tree reaction to environmental stress: a review. *Eur J Forest Res* 124, 319–333.
- Ehleringer, J.R., Dawson, T.E., 1992. Water uptake by plants: perspectives from stable isotope composition. *Plant Cell Environ* 15, 1073–1082.
- Fady, B., Semerci, H., Vendramin, G.G., 2003. EUFOR-GEN Technical Guidelines for Genetic Conservation and Use for Aleppo Pine (*Pinus halepensis*) and Brutia Pine (*Pinus brutia*). International Plant Genetic Resources Institute, Rome, Italy.
- Farquhar, G., O'Leary, M., Berry, J., 1982. On the relationship between carbon isotope discrimination and the intercellular carbon dioxide concentration in leaves. *Funct. Plant Biol* 9, 121–137.
- Farquhar, G., Richards, R., 1984. Isotopic composition of plant carbon correlates with water-use efficiency of wheat genotypes. *Funct. Plant Biol* 11, 539–552.
- Farquhar, G.D., Ehleringer, J.R., Hubick, K.T., 1989. Carbon isotope discrimination and photosynthesis. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 40, 503–537.
- Feeley, S.R., Kolb, T.E., Wagner, M.R., Covington, W.W., 1998. Influence of thinning and burning restoration treatments on presettlement ponderosa pines at the Gus Pearson Natural Area. *Can. J. For. Res.* 28, 1295–1306.
- Fernandes, T.J., del Campo, A.D., Herrera, R., Molina, A.J., 2016. Simultaneous assessment, through sap flow and stable isotopes, of water use efficiency (WUE) in thinned pines shows improvement in growth, tree-climate sensitivity and WUE, but not in WUEi. *For. Ecol. Manag.* 361, 298–308.
- Francey, R.J., Farquhar, G.D., 1982. An explanation of $^{13}\text{C}/^{12}\text{C}$ variations in tree rings. *Nature* 297, 28–31.
- García de la Serrana, R., Vilagrosa, A., Alloza, J.A., 2015. Pine mortality in southeast Spain after an extreme dry and warm year: interactions among drought stress, carbohydrates and bark beetle attack. *Trees* 29, 1791–1804.
- Gessler, A., Ferrio, J.P., Hommel, R., Treydte, K., Werner, R.A., Monson, R.K., 2014. Stable isotopes in tree rings: towards a mechanistic understanding of isotope fractionation and mixing processes from the leaves to the wood. *Tree Physiol.* 34, 796–818.
- Gessler, A., Cailleret, M., Joseph, J., Schönbeck, L., Schaub, M., 2018. Drought induced tree mortality – a tree-ring isotope based conceptual model to assess mechanisms and predispositions. *New Phytol* 219, 485–490.
- Giuggiola, A., Bugmann, H., Zingg, A., Dobbertin, M., Rigling, A., 2013. Reduction of stand density increases drought resistance in xeric Scots pine forests. *For. Ecol. Manag.* 310, 827–835.
- Giuggiola, A., Ogée, J., Rigling, A., Gessler, A., Bugmann, H., Treydte, K., 2016. Improvement of water and light availability after thinning at a xeric site: which matters more? A dual isotope approach. *New Phytol* 210, 108–121.
- Grossiord, C., Sevanto, S., Dawson, T.E., Adams, H.D., Collins, A.D., ... McDowell, N.G., 2017. Warming combined with more extreme precipitation regimes modifies the water sources used by trees. *New Phytologist* 213, 584–596.
- Hernández, A., Arcechea, E., 2017. Tratamientos para el incremento de la diversidad en masas artificiales de *Pinus halepensis*. Proceedings of the 7^o CFE, Plasencia, Spain 26–30 June.
- Holmes, R.L., 1983. Computer-assisted quality control in tree-ring dating and measurement. *Tree-Ring Bull.* 43, 69–75.
- Kaye, J.P., Hart, S.C., 1998. Ecological restoration alters nitrogen transformations in a ponderosa pine–bunchgrass ecosystem. *Ecol. Appl.* 8, 1052–1060.
- Klein, T., Hemming, D., Lin, T., Grunzweig, J.M., Maseyk, K., Rotenberg, E., Yakir, D., 2005. Association between tree-ring and needle delta C-13 and leaf gas exchange in *Pinus halepensis* under semi-arid conditions. *Oecologia* 144, 45–54.
- Lloret, F., Keeling, E.G., Sala, A., 2011. Components of tree resilience: effects of successive low-growth episodes in old ponderosa pine forests. *Oikos* 120, 1909–1920.
- López-Serrano, F.R., de las Heras, J., González-Ochoa, A.I., García-Morote, F.A., 2005. Effects of silvicultural treatments and seasonal patterns on foliar nutrients in young post-fire *Pinus halepensis* forest stands. *For. Ecol. Manag.* 210, 321–336.
- McDowell, N.G., Adams, H.D., Bailey, J.D., Hess, M., Kolb, T.E., 2006. Homeostatic maintenance of ponderosa pine gas exchange in response to stand density changes. *Ecol. Appl.* 16, 1164–1182.
- Maestre, F., Cortina, J., 2004. Are *Pinus halepensis* plantations useful as a restoration tool in semiarid Mediterranean areas? *For. Ecol. Manag.* 198, 303–317.
- Mäkinen, H., Isomäki, A., 2004. Thinning intensity and growth of Scots pine stands in Finland. *For. Ecol. Manag.* 201, 311–325.
- Martin-Benito, D., Cherubini, P., del Río, M., et al., 2008. Growth response to climate and drought in *Pinus nigra* Arn. trees of different crown classes. *Trees* 22, 363–373.
- Martin-Benito, D., Del Río, M., Heinrich, I., Helle, G., Canellas, I., 2010. Response of climate-growth relationships and water use efficiency to thinning in a *Pinus nigra* afforestation. *For. Ecol. Manag.* 259, 967–975.
- Martínez-Vilalta, J., Pinol, J., 2002. Drought-induced mortality and hydraulic architecture in pine populations of the NE Iberian Peninsula. *For. Ecol. Manag.* 161, 247–256.
- McCarroll, D., Loader, N.J., 2004. Stable isotopes in tree rings. *Quat. Sci. Rev.* 23, 771–801.
- McDowell, N.G., Allen, C.D., 2015. Darcy's law predicts widespread forest mortality under climate warming. *Nat. Clim. Chang.* 5, 669–672.
- McDowell, N., Brooks, J.R., Fitzgerald, S.A., Bond, B.J., 2003. Carbon isotope discrimination and growth response of old *Pinus ponderosa* trees to stand density reductions. *Plant Cell Environ.* 26, 631–644.
- McDowell, N.G., Adams, H.D., Bailey, J.D., Hess, M., Kolb, T.E., 2006. Homeostatic maintenance of ponderosa pine gas exchange in response to stand density changes. *Ecol. Appl.* 16, 1164–1182.
- Molina, A.J., del Campo, A.D., 2012. The effects of experimental thinning on throughfall and stemflow: a contribution towards hydrology-oriented silviculture in Aleppo pine plantations. *For. Ecol. Manag.* 269, 206–213.
- Moreno-Gutiérrez, C., Barbera, G., Nicolas, E., De Luis, M., Castillo, V., Martínez-Fernández, F., Querejeta, J., 2011. Leaf $\delta^{18}\text{O}$ of remaining trees is affected by thinning intensity in a semiarid pine forest. *Plant Cell Environ.* 34, 1009–1019.
- Moreno-Gutiérrez, C., Battipaglia, G., Cherubini, P., Saurer, M., Nicolás, E., Contreras, S., Querejeta, J.L., 2012. Stand structure modulates the long-term vulnerability of *Pinus halepensis* to climatic drought in a semiarid Mediterranean ecosystem. *Plant Cell Environ.* 35, 1026–1039.
- Navarro-Cerrillo, R.M., Sánchez-Salguero, R., Herrera, R., Ceacero Ruiz, C., Moreno-Rojas, J., Delgado Manzanedo, R., López-Quintanilla, J., 2016. Contrasting growth and water use efficiency after thinning in mixed *Abies pinsapo*–*Pinus pinaster*–*Pinus sylvestris* forests. *J. Forest Sci.* 62, 53–64.
- Navarro-Cerrillo, R.M., Sánchez-Salguero, R., Rodríguez, C., Lazo, J.D., Moreno-Rojas, J.M., Palacios-Rodríguez, G., Camarero, J.J., 2019. Is thinning an alternative when trees could die in response to drought? The case of planted *Pinus nigra* and *P. sylvestris* stands in southern Spain. *For. Ecol. Manag.* 433, 313–324.
- Oliver, J., Bogino, S., Rathgeber, C., Bonnesœur, V., Bravo, F., 2014. Thinning has a positive effect on growth dynamics and growth–climate relationships in Aleppo pine (*Pinus halepensis*) trees of different crown classes. *Ann. For. Sci.* 71, 395–404.
- Increased water-use efficiency during the 20th century did not translate into enhanced tree growth. In: Peñuelas, J., Canadell, J.G., Ogaya, R. (Eds.), *Glob Ecol. Biogeogr.* 20, 597–608.
- Pemán, J., García, V., Vadell Guiral, E., 2009. In: S.E.F.C. Castilla y León, Junta (Eds.), *Reconstrucción de la estadística de la actividad repobladora desde 1879 hasta nuestros días en 5^o CFE. Montes y sociedad: saber qué hacer*, p. 669 Avila.
- Powers, M.D., Pregitzer, K.S., Palik, B.J., Webster, C.R., 2010. Wood $\delta^{13}\text{C}$, $\delta^{18}\text{O}$ and radial growth responses of residual red pine to variable retention harvesting. *Tree Physiol.* 30, 326–334.
- Prescott, C., deMontigny, L., Harper, G., 2019. Eighteen-year growth responses to thinning and fertilization of a height-repressed lodgepole pine stand in interior British Columbia. *For. Chron.* 95, 207–221.
- Quézel, P., 2000. Taxonomy and biogeography of Mediterranean pines (*Pinus halepensis* and *P. brutia*). In: Neeman, G., Trabaud, L. (Eds.), *Ecology, Biogeography and Management of *Pinus halepensis* and *P. brutia* Forest Ecosystems in the Mediterranean Basin*. Backhuys Publishers, Leiden, pp. 1–12.
- Roden, J., Siegwolf, R., 2012. Is the dual-isotope conceptual model fully operational? *Tree Physiol.* 32, 1179–1182.
- Ruano, I., Rodríguez-García, E., Bravo, F., 2013. Effects of pre-commercial thinning on growth and reproduction in post-fire regeneration of *Pinus halepensis* Mill. *Ann. Forest Sci.* 70, 357–366.
- Sánchez-Salguero, R., Navarro-Cerillo, R.M., Camarero, J.J., Fernández-Cancio, A., 2012. Selective drought-induced decline of pine species in southeastern Spain. *Clim. Chang.* 113, 767–785.
- Sánchez-Salguero, R., Linares, J.C., Camarero, J.J., Madrigal-González, J., Hevia, A., Sánchez-Miranda, Á., Ballesteros-Cánovas, J.A., Alfaro-Sánchez, R., García-Cervigón, A.I., Bigler, C., 2015. Disentangling the effects of competition and climate on individual tree growth: a retrospective and dynamic approach in Scots pine. *For. Ecol. Manag.* 358, 12–25.
- Sarris, D., Siegwolf, R., Körner, C., 2013. Inter- and intra-annual stable carbon and oxygen isotope signals in response to drought in Mediterranean pines. *Agric. For. Meteorol.* 168, 59–68.
- Saurer, M., Siegwolf, R.T.W., Schweingruber, F.H., 2004. Carbon isotope discrimination indicates improving water-use efficiency of trees in northern Eurasia over the last 100 years. *Glob Ch Biol* 10, 2109–2120.
- Scheidegger, Y., Saurer, M., Bahn, M., Siegwolf, R., 2000. Linking stable oxygen and carbon isotopes with stomatal conductance and photosynthetic capacity: a conceptual model. *Oecologia* 125, 350–357.
- Sohn, J.A., Gebhardt, T., Ammer, C., Bauhus, J., Häberle, K.-H., Matyssek, R., Grams, T.E.E., 2013. Mitigation of drought by thinning: short-term and long-term effects on growth and physiological performance of Norway spruce (*Picea abies*). *For. Ecol. Manag.* 308, 188–197.
- Sohn, J.A., Saha, S., Bauhus, J., 2016. Potential of forest thinning to mitigate drought stress: a meta-analysis. *For. Ecol. Manag.* 380, 261–273.
- Tague, C.L., Moritz, M., Hanan, E., 2019. The changing water cycle: the eco-hydrologic impacts of forest density reduction in Mediterranean (seasonally dry) regions. *WIREs Water* 6, e1350.
- Tomas-Burguera, M., Vicente-Serrano, S.M., Beguería, S., Reig, F., Latorre, B., 2019. Reference crop evapotranspiration database in Spain (1961–2014). *Earth Syst. Sci. Data* 11, 1917–1930.
- Tsamir, M., Gottleib, S., Preisler, Y., Rotenberg, E., Tatarinov, F., Yakir, D., ... Klein, T., 2019. Stand density effects on carbon and water fluxes in a semi-arid forest, from leaf to stand-scale. *For. Ecol. Manag.* 453, 117573.
- Vallejo, V.R., Alloza, J.A., 1998. The restoration of burned lands: the case of Eastern Spain. In: Moreno, J.M. (Ed.), *Large Forest Fires*. Backhuys Publishers, Leiden, The Netherlands, pp. 91–108.
- Vicente-Serrano, S.M., Beguería, S., López-Moreno, J.I., 2010. A multiscalar drought index sensitive to global warming: the standardized precipitation evapotranspiration index. *J. Clim.* 23, 1696–1718.
- Vicente-Serrano, S.M., Tomás-Burguera, M., Beguería, S., Reig-Gracia, F., Latorre, B., Peña-Gallardo, M., Luna, Y., Morata, A., González-Hidalgo, J.C., 2017. A high resolution dataset of drought indices for Spain. *Data* 2, 22.
- Warren, C.R., McGrath, J.F., Adams, M.A., 2001. Water availability and carbon isotope discrimination in conifers. *Oecologia* 127, 476–486.
- World Reference Base for Soil Resources, 2014. Update 2015. *World Soil Resources Reports*. 106. FAO, Rome, p. 2015.