



















- meat during refrigerated storage. *Int J Food Microbiol* 2010; 137:175-80.
43. Gutierrez J, Rodriguez G, Barry-Ryan C, Bourke P. Efficacy of plant essential oils against food-borne pathogens and spoilage bacteria associated with ready to eat vegetables: antimicrobial and sensory screening. *J Food Protect* 2008;71:1846-54
  44. Basavegowda N, Baek KH. Synergistic antioxidant and antibacterial advantages of essential oils for food packaging applications. *Biomolecules* 2021;2:1267.
  45. García-García R, López-Malo A, Palou E. Bactericidal action of binary and ternary mixtures of carvacrol, thymol, and eugenol against *Listeria innocua*. *J Food Sci* 2011;76:95–100.
  46. Camele I, Elshafie HS, Caputo L, De Feo V. Anti-quorum sensing and antimicrobial effect of Mediterranean plant essential oils against phytopathogenic bacteria. *Front Microbiol* 2019;10: 2619.
  47. Kerekes EB, Deák É, Takó M, et al. Anti-biofilm forming and anti-quorum sensing activity of selected essential oils and their main components on food-related micro-organisms. *J. Appl Microbiol* 2013;115:933–942.
  48. Luciardi MC, Blázquez MA, Cartagena E, et al. Mandarin essential oils inhibit quorum sensing and virulence factors of *Pseudomonas aeruginosa*. *LWT Food Sci Technol* 2016;68: 373–380.

Non-commercial use only

---

Online supplementary material:

Table S1. GC-MS of the total identified components in *O. vulgare* EO.

Table S2. GC-MS of the total identified components in *S. officinalis* EO.