

## Supplementary Material

### **Antioxidant Potential of Herbal Preparations and Components from *Galactites elegans* (All.) Nyman ex Soldano.**

Omar Tebboub,<sup>1,§</sup> Roberta Cotugno,<sup>2,§</sup> Feyza Oke-Altuntas,<sup>3</sup> Mohamed Bouheroum,<sup>1</sup> İbrahim Demirtas,<sup>4</sup> Massimiliano D'Ambola,<sup>2</sup> Nicola Malafrente,<sup>2</sup> and Antonio Vassallo<sup>5\*</sup>

<sup>1</sup> Unit de Recherche de Valorisation des Ressources Naturelles, Molécules Bioactives, Analyses Physico chimiques et Biologiques(VARENBIOMOL), Faculty of Exact Sciences, Université des Freres Mentouri Constantine1, Algeria

<sup>2</sup> Dipartimento di Farmacia, Università di Salerno, Fisciano, (SA), Italy.

<sup>3</sup> Department of Biology, Faculty of Science, Gazi University, Ankara 06500, Turkey

<sup>4</sup> Laboratory of Plant Research, Department of Chemistry, Faculty of Science, Cankiri Karatekin University, Cankiri, Turkey

<sup>5</sup> Dipartimento di Scienze, Università degli Studi della Basilicata, Potenza, (PZ), Italy

§ Equal contributing authors

\*Correspondence should be addressed to Antonio Vassallo; antonio.vassallo@unibas.it

## Contents

Figure S1. <sup>1</sup>H NMR spectrum of compound **1** (CD<sub>3</sub>OD, 600 MHz).

Figure S2. <sup>1</sup>H NMR spectrum of compound **2** (CD<sub>3</sub>OD, 600 MHz).

Figure S3. <sup>1</sup>H NMR spectrum of compound **3** (CD<sub>3</sub>OD, 600 MHz).

Figure S4. <sup>1</sup>H NMR spectrum of compound **4** (CD<sub>3</sub>OD, 600 MHz).

Figure S5. <sup>1</sup>H NMR spectrum of abietin (CD<sub>3</sub>OD, 600 MHz).

Figure S6. <sup>1</sup>H NMR spectrum of luteolin 4'-*O*-glucuronide (CD<sub>3</sub>OD, 600 MHz).

Figure S7. <sup>1</sup>H NMR spectrum of naringenin-7-*O*-neohesperidoside (CD<sub>3</sub>OD, 600 MHz).

Figure S8. <sup>1</sup>H NMR spectrum of kaempferol-3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside (CD<sub>3</sub>OD, 600 MHz).

Figure S9. <sup>1</sup>H NMR spectrum of apigenin-7-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside (CD<sub>3</sub>OD, 600 MHz).

Figure S10. <sup>1</sup>H NMR spectrum of quercitrin (CD<sub>3</sub>OD, 600 MHz).

Figure S11. <sup>1</sup>H NMR spectrum of quercetin-3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside (CD<sub>3</sub>OD, 600 MHz).

Figure S1.  $^1\text{H}$  NMR spectrum of compound **1** ( $\text{CD}_3\text{OD}$ , 600 MHz).




Figure S2.  $^1\text{H}$  NMR spectrum of compound **2** ( $\text{CD}_3\text{OD}$ , 600 MHz).




Figure S3.  $^1\text{H}$  NMR spectrum of compound **3** ( $\text{CD}_3\text{OD}$ , 600 MHz).




Figure S4.  $^1\text{H}$  NMR spectrum of compound **4** ( $\text{CD}_3\text{OD}$ , 600 MHz).




Figure S5.  $^1\text{H}$  NMR spectrum of abietin ( $\text{CD}_3\text{OD}$ , 600 MHz).




Figure S6.  $^1\text{H}$  NMR spectrum of luteolin 4'-*O*-glucuronide ( $\text{CD}_3\text{OD}$ , 600 MHz).






Figure S7.  $^1\text{H}$  NMR spectrum of naringenin-7-*O*-neohesperidoside ( $\text{CD}_3\text{OD}$ , 600 MHz).




Figure S8.  $^1\text{H}$  NMR spectrum of kaempferol-3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside ( $\text{CD}_3\text{OD}$ , 600 MHz).




Figure S9.  $^1\text{H}$  NMR spectrum of apigenin-7-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside ( $\text{CD}_3\text{OD}$ , 600 MHz).




Figure S10.  $^1\text{H}$  NMR spectrum of quercitrin ( $\text{CD}_3\text{OD}$ , 600 MHz).




Figure S11.  $^1\text{H}$  NMR spectrum of quercetin-3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside ( $\text{CD}_3\text{OD}$ , 600 MHz).

