

Article

Role of Ovarian Proteins Secreted by *Toxoneuron nigriceps* (Viereck) (Hymenoptera, Braconidae) in the Early Suppression of Host Immune Response

Rosanna Salvia ^{1,2}, Carmen Scieuzo ^{1,2}, Annalisa Grimaldi ³, Paolo Fanti ¹, Antonio Moretta ¹, Antonio Franco ^{1,2}, Paola Varricchio ⁴, S. Bradley Vinson ⁵ and Patrizia Falabella ^{1,2,*}

¹ Department of Sciences, University of Basilicata, Via dell'Ateneo Lucano 10, 85100 Potenza, Italy; r.salvia@unibas.it (R.S.); carmen.scieuzo@unibas.it (C.S.); paolo.fanti@unibas.it (P.F.); antonio.moretta@unibas.it (A.M.); antonio.franco@unibas.it (A.F.)

² Spinoff XFlies s.r.l, University of Basilicata, Via dell'Ateneo Lucano 10, 85100 Potenza, Italy

³ Department of Biotechnology and Life Science, University of Insubria, Via J.H. Dunant 3, 21100 Varese, Italy; Annalisa.Grimaldi@uninsubria.it

⁴ Department of Agricultural Sciences, University of Naples "Federico II", 80055 Portici, Italy; paolavarricchio9599@gmail.com

⁵ Department of Entomology, Texas A&M University, 370 Olsen Blvd, College Station, TX 77843-2475, USA; bvinson@tamu.edu

* Correspondence: patrizia.falabella@unibas.it

Citation: Salvia, R.; Scieuzo, C.; Grimaldi, A.; Fanti, P.; Moretta, A.; Franco, A.; Varricchio, P.; Vinson, S.B.; Falabella, P. Role of Ovarian Proteins Secreted by *Toxoneuron nigriceps* (Viereck) (Hymenoptera, Braconidae) in the Early Suppression of Host Immune Response. *Insects* **2021**, *12*, 33. <https://doi.org/10.3390/insects12010033>

Received: 10 December 2020

Accepted: 31 December 2020

Published: 5 January 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

Simple Summary: *Toxoneuron nigriceps* is an endoparasitoid of the tobacco budworm *Heliothis virescens*. Parasitoid strategies to survive involve different regulating factors that are injected into the host body together with the egg: the venom and the calyx fluid, containing a Polydnavirus (PDV) and Ovarian Proteins (OPs). The combination of these factors increases the success of parasitism. Although many studies have been reported on venom protein components and the knowledge on PDVs is increasing, little is known on OPs. These secretions are able to interfere early with the host cellular immune response, acting specifically on host haemocytes, cells involved in immune response. Our results show that OPs induce several alterations on haemocytes, including cellular oxidative stress condition and modifications of actin cytoskeleton, so inducing both a loss of haemocyte functionality and cell death. Overall, in synergy with PDV and venom, OPs positively contribute to the evasion of the host immune response by *T. nigriceps*.

Abstract: *Toxoneuron nigriceps* (Viereck) (Hymenoptera, Braconidae) is an endophagous parasitoid of the larval stages of the tobacco budworm, *Heliothis virescens* (Fabricius) (Lepidoptera, Noctuidae). During oviposition, *T. nigriceps* injects into the host body, along with the egg, the venom, the calyx fluid, which contains a Polydnavirus (*T. nigriceps* BracoVirus: *TnBV*), and the Ovarian Proteins (OPs). Although viral gene expression in the host reaches detectable levels after a few hours, a precocious disruption of the host metabolism and immune system is observed right after parasitization. This alteration appears to be induced by female secretions including *TnBV* venom and OPs. OPs, originating from the ovarian calyx cells, are involved in the induction of precocious symptoms in the host immune system alteration. It is known that OPs in braconid and ichneumonid wasps can interfere with the cellular immune response before Polydnavirus infects and expresses its genes in the host tissues. Here we show that *T. nigriceps* OPs induce several alterations on host haemocytes that trigger cell death. The OP injection induces an extensive oxidative stress and a disorganization of actin cytoskeleton and these alterations can explain the high-level of haemocyte mortality, the loss of haemocyte functionality, and so the reduction in encapsulation ability by the host.

Keywords: Ovarian Proteins; host-parasitoid interaction; *Heliothis virescens*; *Toxoneuron nigriceps*

1. Introduction

Parasitoid insects have developed in some species a great variety of adaptations in terms of physiological integration which make them similar to parasites, which constitute the most specialized forms of zoophagy [1]. Parasitoids belong to different insect orders, such as Diptera, Coleoptera, Lepidoptera, Trichoptera, and Neuroptera but they are common above all among the Hymenoptera [2]. Parasitoids develop at the expense of a single victim, the host, killing it [3]. They can be classified by different parameters, such as the number of deposited eggs per host and the parasitization mode [3]. Their host can be an egg, a larva, a pupa or, rarely, adult insects. Parasitoids are generally defined as ectoparasitoids when the juvenile stages feed on the host from the outside of the body and endoparasitoids when the development of the parasitoid takes place inside the host body [4].

During evolution, insects have developed many defense mechanisms as a result of frequent attacks by pathogens and parasites. They have an innate immunity defense, capable of recognizing and identifying a large class of foreign organisms and responding selectively and effectively to attacks [5–8]. The non-self-recognition by insects is due to the molecular interaction between the non-self-molecules and the specific receptors distributed on the host cell surface [9,10]. When the receptors recognize and bind to the proper ligands (non-self-molecules), a signal cascade is activated into the cell, promoting the immune response through the production of antimicrobial peptides (AMPs) and the melanization process [11,12].

To overcome the immune defenses of the host, the parasitoids have developed several strategies that are generally classified as active and passive [13]. The parasitic factor responsible for the active immune suppression in the host include maternal factors such as the venom and the ovarian calyx fluid, which contain the Ovarian Proteins (OPs) and in some cases Polydnavirus (PDV) or virus-like particles (VLP), and embryonic factors such as teratocytes [4,14–17].

PDVs belong to a Polydnaviridae family with unique biological traits, whose members are divided in two groups, Ichnovirus and Bracovirus, all associated with endophagous parasitoids of lepidopteran larvae [18–20]. Several genes expressed, even at an early stage of infection, by PDVs in different tissues of the host, including the haemocytes, have been shown to be implicated in the active suppression of the host immune response [21–27]. The expression of the viral gene in the host tissues does not guarantee an escape of the immune response in the early hours following oviposition. This problem is overcome by some parasitoids by injecting (together with the egg and the venom) the OPs, produced by epithelial cells of the female parasitoid reproductive traits [28,29]. The OPs have not been investigated as exhaustively as the other parasitoid maternal factors.

In this work we investigated the role of OPs of the endophagous parasitoid, the Hymenoptera Braconidae *Toxoneuron nigriceps* (Viereck) on its natural host, the noctuid moth *Heliothis virescens* (Fabricius). The parasitoid is able to lay the egg in all the larval stages of its host which, while reaching the mature larval stage, shows a slowed development, and the inability to pupate [30,31]. This alteration of the host development that allows the development of the parasitoid larva, is due to changes in the neuroendocrine equilibrium, to the redirection of the host biochemical and metabolic processes, in a condition of severe alteration of the host's immune response [32–34]. This regulation of the host physiology is induced by both maternal (fluid of the calyx and venom) and embryonic (teratocytes) factors of parasitic origin [35,36]. The expression of *T. nigriceps* BracoVirus (*TnBV*) genes into host cells determines the functional inactivation of the prothoracic glands in the mature larvae [33–38] and induces apoptosis in haemocytes [26,39]. The teratocytes, cells derived from the embryonic serosal membrane [40] contribute to the arrest of the host development [41] inducing the transformation of 20-hydroxyecdysone into inactive polar metabolites [31,32,42–44]. Among the factors of maternal origin, *TnBV* and venom have been the most studied factors [15,24,26,44]. The virus infects different tissues like the fat body, the haemocytes and the prothoracic glands, in which it actively expresses its genes.

Among the viral genes, *TnBV1* [45,46] and *TnBVank1* [24,26] seem to play an important role in the suppression of the immune response. In the parasitized host a decrease in the number of haemocytes has been observed as early as 2 h up to 24 h after the oviposition, with the lower number of haemocytes observed after 4 h [39]. Two hours after parasitization *H. virescens* haemolymph showed a high number of apoptotic cells and a very low level of cell proliferation [39]. Furthermore, as a result of parasitization, the remaining haemocytes showed strong morphological and structural changes, losing the ability to adhere to external objects and to encapsulate invading microorganisms [39]. The expression of the identified *TnBV* genes, that are responsible for morphological alterations and apoptosis in haemocytes, always occurs at least after 3–4 h [24,26]. We hypothesize that the very early suppression of the host immune response after oviposition is correlated to both a passive mechanism, due to the presence of a fibrous layer on the egg surface, and an active mechanism, due to the action of the possible synergic effect of the venom and OP mixture [47–49]. The venom composition was elucidated by using a combined transcriptomic and proteomic approach [15] and several proteins putatively involved in the alteration of the immune system of the parasitized host identified.

Currently, nothing is known about the function of OPs of *T. nigriceps* and their role in the alterations of the immune system of the host *H. virescens*. Here we present research we did on this aspect. In particular, we believe that OPs can interfere with the cellular immune response before *TnBV* infects and expresses its genes in the host tissues.

2. Materials and Methods

2.1. Insect Rearing

Toxoneuron nigriceps parasitoid was reared according to the methodology described by Vinson et al. [50]. The larvae of the host *Heliothis virescens* were reared on an artificial diet developed by Vanderzant et al. [51] (Corn Earworm Diet, Bioserve, Frenchtown, NJ, USA). The breeding temperature was maintained at $29\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$, both for non-parasitized and parasitized *H. virescens* larvae. *Toxoneuron nigriceps* adults were bred at a temperature of $25\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$. In both cases a photoperiod of 16 h of light and 8 h of darkness was set with a relative humidity of $70\% \pm 5\%$. For the following analysis *H. virescens* larvae parasitized at day one of the last instar were used.

2.2. Calyx Fluid Collection and Ovarian Protein Purification

About 80–100 adult females of *T. nigriceps*, two weeks old, previously anesthetized on ice for 10–15 min, were used for the collection of the calyx fluid, containing *T. nigriceps* BracoVirus (*TnBV*) and Ovarian Protein (OPs). From each female, immersed in a physiological solution of 1× Phosphate-Buffered Saline (PBS), the entire reproductive apparatus was removed, pulling away the ovipositor with a pair of forceps. The isolated ovaries, explanted by one or two females, were placed in a drop of 20 μL of 1× PBS (1 or 2 equivalent females) at $4\text{ }^{\circ}\text{C}$ and the ovarian calyx were dissected to allow the flow of the calyx fluid. The collected liquid was transferred into a 1.5 mL tube (Eppendorf, Hamburg, DE, catalogue number 0030120.086) and centrifuged at $2,000\times g$ for 5 min at $4\text{ }^{\circ}\text{C}$ to remove the eggs and the calyx tissues and any other debris. The supernatant, containing OPs and *TnBV*, was further purified with 0.45 μm Millex PVDF filters (Millipore, Burlington, MA, USA, catalogue number SLHVM33RS) followed by washing 3 times with 1 mL of 1× PBS. To separate *TnBV* a centrifugation was carried out at $30,000\times g$ for 1 h at $4\text{ }^{\circ}\text{C}$ and subsequently the supernatant containing the OPs was concentrated by ultrafiltration in a column with cut-off 3,000 (Millipore, Burlington, MA, USA, catalogue number UFC900308), by centrifuging the sample at $3,000\times g$ until a volume of around 500 μL was obtained. The resulting filtrate was stored at $-80\text{ }^{\circ}\text{C}$ for subsequent uses (Protocol S1).

2.3. Collection of Haemocytes from Larvae of *H. virescens*

Larvae of *H. virescens* on the third day of the last instar were anesthetized in water, sterilized in 70% ethanol (v/v) and subsequently washed in sterile deionized water. The first pair of legs was cut and the hemolymph from the wound was collected and transferred to a centrifuge tube containing 1 mL of MEAD pre-cooled solution in ice [39]. The sample was centrifuged at $400\times g$ for 7 min at 4 °C. The precipitate was washed twice with a MEAD-PBS solution (1:1). Finally, the haemocytes were gently resuspended in 1 mL of Grace Insect Medium (Sigma Aldrich, St. Louis, MO, USA, catalogue number, G8142) containing FBS (Gibco, Gaithersburg, MD, USA, catalogue number A4766801) 10% and anti-biotic-antimycotic 1% (Gibco, Gaithersburg, MD, USA, catalogue number 15240062).

An amount of 1×10^6 haemocyte cells per well were inoculated in 24-well culture plates (Corning Incorporated, New York, NY, USA, catalogue number CLS3527-100EA). OPs (deriving from 1 or 2 equivalent females) or 1× PBS (as control) collected as described above were added to the haemocytes in the culture medium and incubated at 27 °C.

2.4. Cells Viability

To evaluate cell viability after the treatment with OPs, and to compare the effect of parasitization and OPs alone, trypan blue staining (Sigma Aldrich, St Louis, MO, USA, catalogue number T8154) was used on haemocytes, collected from parasitized and non-parasitized larvae. OPs (obtained from 1 or 2 equivalent females) or 1× PBS (as control) were added to the haemocytes collected from non-parasitized larvae in the culture medium and incubated at 27 °C for 24 h. Trypan blue staining was added to the cell suspension at a final concentration of 0.04% and then haemocytes were counted by Neubauer's chamber under microscopy (Eclipse 80i, Nikon, Tokyo, Japan).

2.5. Light Microscopy Haemocyte Observations

The haemocytes incubated with OPs (deriving from 2 equivalent females) at different time (30 min, 1 h and 2 h) or with 1× PBS (negative control) were detached from the well transferred on slides and subjected to different staining methodologies: May Grünwald GIEMSA (Sigma Aldrich, St Louis, MO, USA, catalogue number MG500 and GS500), 2,7 dichlorodihydrofluorescein acetate (H₂DCFDA) (Thermo Fisher Scientific, Waltham, MA, USA, catalogue number D399), and tetramethylrhodamine isothiocyanate (TRITC)-conjugated phalloidin (Sigma Aldrich, St Louis, MO, USA, catalogue number P1951) dyes. For each analyzed parameter, evaluation was performed considering five random fields in three independent replicates in which cells with alteration were counted on the total number of cells.

Haemocytes were fixed with 4% paraformaldehyde for 10 min, washed with 1× PBS, and stained with May–Grünwald dye for 15 min followed by 30 min in 5% Giemsa stain. The cells were washed three times with 1× PBS and the slides were mounted with glycerol (Sigma Aldrich, St Louis, MO, USA, catalogue number G5516) for morphologic observation.

For H₂DCFDA staining, after fixing on slides, cells were incubated in the dark with H₂DCFDA 10 µM for 30 min at room temperature. After washing three times in 1× PBS, the slides were mounted with glycerol.

A further staining was carried out using TRITC-conjugated phalloidin staining, the haemocytes fixed on slides, were incubated with TRITC-conjugated phalloidin diluted 50 µg/mL in 1% BSA (Sigma Aldrich St Louis, MO, USA, catalogue number 05470) for 2 h at room temperature in dark conditions and then, after three washes with 1× PBS, the slides were mounted with glycerol.

For all staining methodologies, the slides were observed microscopically with Nikon Eclipse 80i equipped with a Nikon Plan Fluor 100×/0.5–1.3 Oil Iris objective and the images, five random fields obtained in three independent replicates, were recorded with a

Nikon Digital Sight DS-U1 camera, and the percentage of stained/fluorescent cells was counted on the total number of cells.

2.6. Chromatographic Sphere Staining

Sephadex Fine G50 chromatographic beads (Cytiva, Little Chalfont, England, GB catalogue number 17004202) were stained with the Congo Red dye (Sigma Aldrich, St Louis, MO, USA, catalogue number 75768). The beads were autoclaved in 1 mL of 1× PBS at 121 °C for 20 min and centrifuged at 13,000× g for 5 min. The supernatant was removed and 500 µL of a 1× PBS solution containing 20 mg/mL of Congo Red was added. The beads were autoclaved again at 121 °C for 20 min, centrifuged at 13,000× g for 5 min and repeatedly washed with 1× PBS to remove excess of dye.

2.7. Encapsulation Measurement

Before testing the effect of OPs on encapsulation, we defined the minimum time needed in *H. virescens* to obtain an adequate level of encapsulation. To assess this time we performed an assay, injecting 30 beads into 5 larvae for each, at 3 different dissection times after injection (10 min, 3 h, 6 h) for 3 replicates. The level of encapsulation reached after 6 h was found to be adequate to see the encapsulation effect in subsequent experiments. All the recovered beads were observed under the microscope (Eclipse 80i, Nikon, Tokyo, Japan).

2.8. Injections of OPs and Chromatographic Spheres in *H. virescens* Larvae and Evaluation of *In Vivo* Encapsulation Degree

The injections were performed on larvae of *H. virescens*, on the first day of the fifth instar larva. The larvae were anesthetized in water, sterilized in 70% ethanol (Sigma Aldrich St Louis, MO, USA, catalogue number 51976) (v/v) and subsequently washed in deionized water. Then, they were dried and placed on a Petri dish. Injection of the Sephadex Fine G 50 (50–150 µm) chromatographic beads was performed in the neck membrane, using a Hamilton syringe (Sigma-Aldrich, St Louis, MO, USA, catalogue number HAM80075-1EA) having a needle with 0.13 mm internal diameter and 0.47 mm outer diameter. The chromatographic spheres were injected at different times (10 min, 1 h or 3 h) after OPs injections or 5 µL of 1× PBS. After 6 h from the injection of the spheres, the time necessary for the formation of a complete haemocyte capsule (Figure S1), the larvae were anesthetized in water, sterilized in ethanol 70% (v/v) and washed in deionized water. Subsequently they were dried and placed in a drop of 1× PBS solution. The dissection was performed by cutting the head and longitudinally cutting in the ventral position. The chromatographic spheres attached to the tissues were collected and counted. The spheres were classified according to the degree of encapsulation as: 0 = unencapsulated (no haemocytes layer); 1 = the thickness of capsule is one or more than one layer, but less than a half of the bead's radius; 2 = the thickness of capsule is equal or more than a half of the bead's radius. We considered encapsulated the beads that showed after 6 h case 2, and not encapsulated when the beads showed case 0 or 1.

2.9. Statistical Analysis of Data

Statistical analysis was performed with one-way ANOVA (analysis of variance) and Bonferroni or Tukey post-hoc tests. Statistical differences were analyzed both among all treatments and between control and treated samples at the same experimental time. For the encapsulation assay we first verified that there was no statistical difference among the experimental groups in the percent of recovered beads after the dissection and then we compared the percent of encapsulated beads on the number of recovered beads.

3. Results

3.1. Cell Viability of the Haemocytes after Treatment with Ovarian Proteins

The extracted haemocytes from non-parasitized *H. virescens* larvae (control) showed 85% of cell viability. The haemocytes extracted from larvae of *H. virescens* after 24 h from the parasitization showed only 30% of cell viability. The cell viability of haemocytes cultured in the presence of *T. nigriceps* Ovarian Proteins (OPs) for 24 h did not vary according to their concentration. The cell viability percentages equal to 34% and 38% referred to, respectively, samples treated with OPs deriving from 1 or 2 equivalent *T. nigriceps* females. Table S1 and Figure 1 show the obtained results.




Figure 1. Percentage of haemocyte viability incubated with 1× PBS (control), haemocytes extracted from larvae 24 h after parasitization, haemocytes incubated with Ovarian Proteins (OPs) deriving from 1 or 2 equivalent females. Data are presented as means ± SD ($n = 3$ replicates). Statistical analysis was performed with one-way ANOVA and Tukey post-hoc test. Different letters indicate significant differences (p value < 0.0001).

3.2. Haemocyte Staining

3.2.1. May Grunwald–Giemsa Staining

The haemocytes, treated with OPs deriving from 2 equivalent females at different incubation times (30 min, 1 h and 2 h) were observed after May Grunwald–Giemsa (MGG) staining. The cells showed a different staining as a function of the cytoplasmic pH and this allowed the different cell populations to be counted in the three different examined samples (Figure 2). The percentage of cells with acidophilic cytoplasm and massive vacuolization was greater than control and increased in relation to the different times of OP exposure (Table S2, Figure 3).

Figure 3 shows that after 30 min of incubation the percentage of cells with acidophilic cytoplasm was 27%. After 1 h the percentage was raised to 34% compared to the relative control and after 2 h the percentage reached 41%.




Figure 2. Haemocytes treated with 1× PBS (a) or with OPs for 2 h (b) stained with May Grunwald–Giemsa dye. Scale bar 10 µm. Arrows indicate vacuolization process.




Figure 3. Percentage of haemocytes showing vacuolization, after treatment with OPs deriving from 2 equivalent females at different incubation times (30 min, 1 h and 2 h) and observed after May Grunwald–Giemsa (MGG) staining. Data are presented as means \pm SD ($n = 3$ replicates). Statistical analysis was performed with one-way ANOVA and Bonferroni (for control vs treatment analysis) and Tukey post-hoc test (for all sample analysis). Different letters indicate significant differences among all treatments (p value < 0.0001), asterisks indicate significant differences between control and treated samples at the same experimental time (** p value < 0.01 ; **** p value < 0.0001).

3.2.2. H₂DCFDA Staining

Haemocytes treated with OPs (2 equivalent females) at different times (30 min, 1 h and 2 h) were stained with 2,7 dichlorodihydrofluorescein acetate (H₂DCFDA). H₂DCFDA after entering into the cells, were converted into dichlorodihydrofluorescein (H₂DCF) by intracellular esterases, H₂DCF is rapidly oxidized to a highly fluorescent compound, 2,7-dichlorofluorescein (DCF), only in the presence of reactive oxygen species (ROS).

Figure 4 and Figure S2 show a strong fluorescent signal indicative of high oxidative stress in cells incubated for 2 h with OPs and a very weak fluorescent signal that could constitute the background experiment and/or could be indicative of cell physiological presence of ROS at low concentrations in control cells.




Figure 4. Healthy *in vitro* haemocytes (a) or treated with OPs for 2 h (b) and stained with 2,7 dichlorodihydrofluorescein acetate (H₂DCFDA). Red box = enlargement of cells below. Scale bar 10 μm.

Table S3 and Figure 5 report the percentage of haemocytes showing a strong fluorescent signal indicative of oxidative stress, after incubation with OPs at different times. After 30 min of incubation with Ops, 46% of haemocytes were strongly fluorescent, after 1 h of treatment 64% of cells were fluorescent, after 2 h of incubation with OPs, almost the totality of the haemocytes (90%) showed oxidative stress.




Figure 5. Percentage of haemocytes stained with H₂DCFDA showing oxidative stress, after treatment with OPs deriving from 2 equivalent females at different times of exposure (30 min, 1 h and 2 h). Data are presented as mean ± SD (*n* = 3). Statistical analysis was performed with one-way ANOVA and Bonferroni (for control vs treatment analysis) and Tukey post-hoc test (for all sample analysis). Different letters indicate significant differences among all treatments (*p* value < 0.0001), asterisks indicate significant differences between control and treated samples at the same experimental time (*p* value < 0.0001).

3.2.3. TRITC-Conjugated Phalloidin Staining

Actin filaments were detected using TRITC-Conjugated phalloidin staining in healthy haemocytes (control) or treated with OPs at different times (30 min, 1 h and 2 h) and stained with phalloidin. Figure 6 shows in the treated cells a weak and fragmented signal localized near the nucleus and along the plasma membrane while a strong and homogeneously distributed fluorescent signal is observed in control cells.




Figure 6. In vitro healthy haemocytes (a) or treated with OPs for 2 h (b) and stained with TRITC-Conjugated phalloidin. Scale bar 10 µm.

In Figure 7 and in Table S4 the percentage of cells that show cytoskeletal damage on the total number of cells is reported. After 30 min of incubation with OPs the percentage of cells with cytoskeletal damage was equal to 31% of the total haemocytes, while no damage to actin filaments was observed in control cells. After 1 h the percentage rose to 70% for the treated cells. After 2 h 80% of the treated cells showed cytoskeletal damage.




Figure 7. Percentage of haemocytes stained with phalloidin showing cytoskeletal damage, after treatment with OPs deriving from 2 equivalent females at different times of exposure (30 min, 1 h and 2 h). Data are presented as mean \pm SD ($n = 3$). Statistical analysis was performed with one-way ANOVA and Bonferroni (for control vs treatment analysis) and Tukey post-hoc test (for all sample analysis). Different letters indicate significant differences among all treatments (p value < 0.0001), asterisks indicate significant differences between control and treated samples at the same experimental time (p value < 0.0001).

3.3. In Vivo Encapsulation

To assess whether the alterations induced by OPs influenced the ability of the haemocytes to recognize and encapsulate foreign intruders, a test was performed by injecting OPs (2 females equivalent) or 1 \times PBS into *H. virescens* larvae at different times (10 min, 1 h and 3 h) before injection of chromatographic spheres used as non-self-material. Chromatographic spheres were injected also in *H. virescens* parasitized larvae. The spheres

were classified according to the level of encapsulation as: 0 = unencapsulated (no haemocytes layer); 1 = the thickness of capsule is one or more than one layer, but less than a half of the bead's radius; 2 = the thickness of capsule is equal or more than a half of the bead's radius. We considered encapsulated exclusively the spheres of type 2 (Figure S1). Figure 8 and Table S5 show a strong reduction in the encapsulation capacity in the different treatments.




Figure 8. Percentage of encapsulation of chromatographic spheres extracted after 6 h from larvae injected with 1× PBS (control), after parasitization or OP treatment at 10 min, 1 h or 3 h before injection of spheres. Data are presented as mean \pm SD ($n = 3$). Statistical analysis was performed with one-way ANOVA and Bonferroni (for control vs treatment and parasitized vs OPs analysis) and Tukey post-hoc test (for all sample analysis). Different letters indicate significant differences among all treatments (p value < 0.0001), asterisks indicate significant differences between control and treated samples at the same experimental time (p value < 0.0001), dots indicate significant differences between parasitized and OP treatment at the same experimental time (•• p value < 0.01 , • p value < 0.05).

4. Discussion

Endoparasitoid insects have developed, during evolution, mechanisms capable to alter the physiology and biochemistry of the host, and they are able to create an environment suitable for the development of their progeny [4,27,39,52]. Such alterations imply the manipulation of the host's immune system. The suppression of the immune reaction in hosts parasitized by endoparasitoid hymenopterans is a very complex syndrome, characterized by several molecular mechanisms integrating each other [39,43,53]. The maternal factors, introduced into the host during the oviposition, are the main protagonists of the physiological alterations observed in the parasitized hosts [54,55]. These factors include venom proteins, symbiotic Polydnavirus (PDV), virus-like particles (VLP) and ovarian proteins (OPs) [4,15,16,49,52,56,57]. Among these, the PDVs have received a great interest over the years as an important source of useful genes [19]. PDVs are associated exclusively with endoparasitoid wasps of lepidopteran larvae [43,58]. PDVs replicate only in the cells of the wasp ovarian calyxes and are released in the lumen of the oviduct [28,29]. Once introduced into the host, PDVs infect tissues and actively express their genes [59]. The products of these genes induce a wide range of alterations to the neuroendocrine and immune system [27]. Many genes of viral origin have been isolated and characterized [4,24,26,38,44,45,60–65], but their expression requires several hours to be able to perform the immunosuppressive effect. Because the insect's immune system is already functioning within minutes of recognizing the parasitoid, the parasitoid needs to alter its defense mechanisms in a short time [66]. Therefore, other maternal factors, such as OPs [28,49,67] and venom [15,56,68–71], or the proteins that cover and protect the parasitoid egg [72–75] must necessarily set to mediate immune-suppression in the early stages of parasitization. Luckhart and Webb [76] demonstrated the immunosuppressive intervention of OPs after 30 min from parasitization in *Camponotus sonorensis*. As already showed by Ferrarese et al.

[39], there is a reduction of the haemocyte titer in the haemolymph following parasitization, probably due also to the action of the venom and/or OPs that induce, after few hours, a high cellular mortality. The molecular mechanisms that induce such alterations have not been clarified yet.

In this study we focused on the role of OPs in the host-parasitoid system *Heliothis virescens*-*Toxoneuron nigriceps*. OPs seem to interfere with the cell-mediated immune system in the minutes following parasitization, before *T. nigriceps* BracoVirus (*TnBV*) infects host tissues and expresses its genes. The OPs induce a series of alterations affecting the haemocytes and carrying the cell to death.

In haemocytes treated with OPs many alterations are recorded, including mitochondrial alterations accompanied by ATP depletion, loss of transmembrane potential and the consequent increase in reactive oxygen species (ROS) in the cytoplasm. Mitochondria, therefore, can be a very important target in the attempt to alter cellular function. A similar picture has been described as one of the different manifestations of cell death [77]. Furthermore, several studies suggest that the alteration of the actin cytoskeleton is strongly linked to the increase in intracellular ROS levels and apoptosis [78,79]. ROS are physiologically present in cells at low concentrations and increase considerably in the case of severe mitochondrial damage [80,81]. In our system, haemocytes treated with OPs show that oxidative stress often occurs with cytoskeletal damage to actin filaments; these changes may explain the high levels of haemocyte mortality. The vacuolization observed in the haemocytes already after 30 min from exposure to OPs, confirms the observations reported by Ferrarese et al. [39] on haemocytes extracted from parasitized larvae. These high levels of oxidizing agents, together with a change in pH, probably linked to the increase of ROS, can negatively affect survival and cellular functionality [82,83]. All the observed alterations induce a loss of functionality in the haemocytes. In fact, by injecting OPs and chromatographic spheres into the larval haemocoel, a loss of encapsulation capacity by the haemocytes was observed. We observed a strong reduction of encapsulation both in parasitized larvae and when treated with OPs, as already reported for the endoparasitoid *Macrocentrus cingulum* in which OP treatment inhibited the encapsulation in a dose-dependent manner [84]. A *T. nigriceps* OP stronger effect was recorded at 1 h. This allowed the hypothesis that the inhibition of haemocyte's encapsulation capacity is caused not only by OPs but also by venom and *TnBV*. Indeed, a previous study [15], among *T. nigriceps* venom proteins, identified the Calreticulin protein, whose role in inhibition of encapsulation has been reported in several parasitoids [85–88]. On the contrary in *M. cingulum* the mix of venom and OPs did not have a different effect compared to OPs alone, suggesting that, in this case, there are no synergic effect of venom and OPs in suppressing the encapsulation by *Ostrinia furnacalis* haemocytes [84]. Moreover, we observed a higher percentage of unencapsulated beads in larvae parasitized 3 h before injection of beads compared to larvae treated with OPs. This could be due to the expression of *TnBV* genes that act on the host immune system [24,26].

The present work confirms the hypothesis that OPs have short-term immunosuppressive activity also in the host-parasitoid system *H. virescens*-*T. nigriceps*, as reported in other host-parasitoid system, such as *Chilo suppressalis*-*Cotesia chilonis* [89]. To guarantee the success of parasitization a synergic action of all parasitic factors is required. Several *TnBV* genes have been reported to be responsible for morphological alterations and apoptosis in haemocytes, but their expression occurs at least after 3–4 h [24,26]. The fibrous layer on the *T. nigriceps* egg surface [47] venom proteins [15] and OPs ensure an escape from host cellular immune responses immediately after oviposition.

Knowledge of the sequence of biologically active proteins will probably allow us to understand the mechanisms by which they act.

5. Conclusions

Little information on the role of Ovarian Proteins (OPs) is available in the literature. Most of the studies concerning host-parasitoid systems, and in particular the parasitic factors, focus on Polydnavirus (PDV), if they are present, and venom. With this study we provide valuable information on possible functions of these secretions on host haemocytes, in the host-parasitoid system *Heliothis virescens*–*Toxoneuron nigriceps*. Experiments performed on haemocytes treated with OPs showed many alterations, including the increase of reactive oxygen species (ROS) in the cytoplasm and actin cytoskeleton disruption. These changes provoke both reduction in haemocyte functionality, losing encapsulation capacity, and increasing cellular death. Our study shows that OPs have a significant action, especially regarding the functional alteration of haemocytes, the cells of the host immune response. Overall OPs, together with other maternal factors (venom and PDV) actively contribute to suppress the host immune response, supporting the development of the parasitoid larva and the success of the parasitism.

This study underlines the importance of OPs and suggests further in-depth analysis to characterize the entire protein profile of this mixture of parasitic origin, and also specific analysis to identify the active proteins responsible for the described effects.

Supplementary Materials: The following are available online at www.mdpi.com/2075-4450/12/1/33/s1, Protocol S1: Extraction and purification of *Toxoneuron nigriceps* Ovarian Proteins (OPs), Table S1: Data obtained counting haemocytes incubated with 1×PBS solution (control), haemocytes extracted from larvae 24 h after parasitization and OPs derived from 1 or 2 equivalent females, Table S2: Data obtained from the counts of healthy cells or treated with OPs at different times (30 min, 1 h and 2 h) showing vacuolization, Table S3: Data obtained from counting healthy cell or treated with OPs at different times (30 min, 1 h and 2 h) showing ROS positive cells, Table S4: Data obtained from haemocytes stained with phalloidin showing cytoskeletal damage, after OPs treatment at different exposure times (30 min, 1 h and 2 h), Table S5: Data obtained from haemocytes after injection of chromatographic spheres and after OPs treatment at different exposure times (30 min, 1 h and 2 h) or parasitization, Figure S1: Chromatographic spheres extracted from 5 larvae for each time (10 min (a), 3 h (b) and 6 h (c) after OP injection, Figure S2: Healthy haemocytes or treated with OPs for 2 h and stained with DAPI and H₂DCFDA.

Author Contributions: Conceptualization, P.F. (Patrizia Falabella); methodology, R.S., A.G., C.S.; data curation, P.F. (Paolo Fanti), P.V., R.S.; writing—original draft preparation, P.F. (Patrizia Falabella); writing—review and editing, R.S., C.S., A.G., P.F. (Paolo Fanti), A.M., A.F., P.V., S.B.V., P.F. (Patrizia Falabella); supervision, P.F. (Patrizia Falabella). All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by University of Basilicata.

Data Availability Statement: Data is contained within this article and the supplementary material.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Quicke, D.L.J. *Parasitic Wasps*; Chapman & Hall: London, UK, 1997.
2. Heraty, J. Parasitoid Biodiversity and Insect Pest Management. In *Insect Biodiversity: Science and Society*, 2nd ed.; Footitt, R.G., Adler, P.H., Eds.; John Wiley & Sons: Hoboken, NJ, USA, 2017; pp. 603–625.
3. Godfray, H.C.J. Parasitoids. In *Encyclopedia of Biodiversity*, 2nd ed.; Levin, S.A., Ed.; Elsevier: Amsterdam, The Netherlands; 2007; pp. 1–13.
4. Pennacchio, F.; Strand, M.R. Evolution of developmental strategies in parasitic Hymenoptera. *Annu. Rev. Entomol.* **2006**, *51*, 233–258.
5. Schmidt, O.; Theopold, U.; Strand, M.R. Innate immunity and evasion by insect parasitoids. *BioEssays* **2001**, *23*, 344–351.
6. Bangham, J.; Jiggins, F.; Lemaitre, B. Insect immunity: The post-genomic era. *Immunity* **2006**, *25*, 1–5.
7. Sheehan, G.; Garvey, A.; Croke, M.; Kavanagh, K. Innate humoral immune defences in mammals and insects: The same, with differences? *Virulence* **2018**, *9*, 1625–1639.
8. Brivio, M.F.; Mastore, M. When Appearance Misleads: The Role of the Entomopathogen Surface in the Relationship with Its Host. *Insects* **2020**, *11*, 387.
9. Welchman, D.P.; Aksoy, S.; Jiggins, F.; Lemaitre, B. Insect immunity: From pattern recognition to symbiont-mediated host defense. *Cell Host Microbe* **2009**, *6*, 107–114.

10. Wang, X.; Zhang, Y.; Zhang, R.; Zhang, J. The diversity of pattern recognition receptors (PRRs) involved with insect defense against pathogens. *Curr. Opin. Insect Sci.* **2019**, *33*, 105–110.
11. Sheehan, G.; Farrell, G.; Kavanagh, K. Immune priming: The secret weapon of the insect world. *Virulence* **2020**, *11*, 238–246.
12. Nakhleh, J.; Moussawi, L.E.; Osta, M.A. Chapter Three—The Melanization Response in Insect Immunity. *Adv. Insect Physiol.* **2017**, *52*, 83–109.
13. Asgari, S. Venoms from Endoparasitoids. In *Parasitoid Viruses*; Beckage, N.E., Drezen, J.M., Eds.; Academic Press: Cambridge, MA, USA, 2012; pp. 217–231.
14. Strand, M.R. Teratocytes and their functions in parasitoids. *Curr. Opin. Insect Sci.* **2014**, *6*, 68–73.
15. Laurino, S.; Grossi, G.; Pucci, P.; Flagiello, A.; Bufo, S.A.; Bianco, G.; Salvia, R.; Vinson, S.B.; Vogel, H.; Falabella, P. Identification of major *Toxoneuron nigriceps* venom proteins using an integrated transcriptomic/proteomic approach. *Insect Biochem. Mol. Biol.* **2016**, *76*, 49–61.
16. Leobold, M.; Bézier, A.; Pichon, A.; Herniou, E.A.; Volkoff, A.N.; Drezen, J.M. The domestication of a large DNA virus by the wasp *Venturia canescens* involves targeted genome reduction through pseudogenization. *Genome Biol. Evol.* **2018**, *10*, 1745–1764.
17. Salvia, R.; Grimaldi, A.; Girardello, R.; Scieuzo, C.; Scala, A.; Bufo, S.A.; Vogel, H.; Falabella, P. *Aphidius ervi* Teratocytes Release Enolase and Fatty Acid Binding Protein Through Exosomal Vesicles. *Front. Physiol.* **2019**, *10*, 715.
18. Fath-Goodin, A.; Webb, B.A. Polydnviruses: General Features. In *Encyclopedia of Virology*, 3rd ed.; Mahy, B.W.J., Van Regenmortel, M.H.V., Eds.; Academic Press: Cambridge, MA, USA, 2008; pp. 256–261.
19. Strand, M.R.; Burke, G.R. Polydnviruses: From discovery to current insights. *Virology* **2015**, *479–480*, 393–402.
20. Volkoff, A.N.; Huguet, E. Polydnviruses (Polydnviridae, Bracovirus, and Ichnovirus). In *Reference Module in Life Sciences*; Elsevier, Amsterdam, The Netherlands; 2020.
21. Lovallo, N.C.; McPherson, B.A.; Cox-Foster, D.L. Effects of the polydnvirus of *Cotesia congregata* on the immune system and development of non-habitual hosts of the parasitoid. *J. Insect Physiol.* **2002**, *48*, 517–526.
22. Yin, L.; Zhang, C.; Qin, J.; Wang, C. Polydnvirus of *Campoletis chloridae*: Characterization and temporal effect on host *Helicoverpa armigera* cellular immune response. *Arch. Insect Biochem. Physiol.* **2003**, *52*, 104–113.
23. Barat-Houari, M.; Hilliou, F.; Jousset, F.X.; Sofer, L.; Deleury, E.; Rocher, J.; Ravallec, M.; Galibert, L.; Delobel, P.; Feyereisen, R.; Fournier, P.; et al. Gene expression profiling of *Spodoptera frugiperda* hemocytes and fat body using cDNA microarray reveals polydnvirus-associated variations in lepidopteran host genes transcript levels. *BMC Genom.* **2006**, *7*, 160.
24. Falabella, P.; Varricchio, P.; Provost, B.; Espagne, E.; Ferrarese, R.; Grimaldi, A.; de Eguileor, M.; Fimiani, G.; Ursini, M.V.; Malva, C.; et al. Characterization of the IkappaB-like gene family in polydnviruses associated with wasps belonging to different Braconid subfamilies. *J. Gen. Virol.* **2007**, *88*, 92–104.
25. Gueguen, G.; Kalamarz, M.E.; Ramroop, J.; Uribe, J.; Govind, S. Polydnviral Ankyrin Proteins Aid Parasitic Wasp Survival by Coordinate and Selective Inhibition of Hematopoietic and Immune NF-kappa B Signaling in Insect Hosts. *PLoS Pathog.* **2013**, *9*, e1003580.
26. Salvia, R.; Grossi, G.; Amoresano, A.; Scieuzo, C.; Nardiello, M.; Giangrande, C.; Laurenzana, I.; Ruggieri, V.; Bufo, S.A.; Vinson, S.B.; et al. The multifunctional polydnvirus TnBVANK1 protein: Impact on host apoptotic pathway. *Sci. Rep.* **2017**, *7*, 11775.
27. Ye, X.Q.; Shi, M.; Huang, J.H.; Chen, X.X. Parasitoid polydnviruses and immune interaction with secondary hosts. *Dev. Comp. Immunol.* **2018**, *83*, 124–129.
28. Tanaka, K.; Matsumoto, H.; Hayakawa, Y. Analysis in the course of polydnvirus replication in ovarian calyx cells of the parasitoid wasp, *Cotesia kariyai* (Hymenoptera: Braconidae). *Appl. Entomol. Zool.* **2002**, *37*, 323–328.
29. Park, B.; Yonggyun, K. Polydnvirus replication at ovarian calyx in *Cotesia plutellae* and endocrine impact. *Korean J. Appl. Entomol.* **2010**, *43*, 225–231.
30. Lewis, W.J.; Vinson, S.B. Egg and larval development of *Cardiochiles nigriceps*. *Ann. Entomol. Soc. Am.* **1968**, *61*, 561–565.
31. Pennacchio, F.; Vinson, S.B.; Tremblay, E. Growth and development of *Cardiochiles nigriceps* Viereck (Hymenoptera, Braconidae) larvae and their synchronization with some changes of the hemolymph composition of their host, *Heliothis virescens* (F.) (Lepidoptera, Noctuidae). *Arch. Insect Biochem. Physiol.* **1993**, *24*, 65–77.
32. Pennacchio, F.; Vinson, S.B.; Tremblay, E.; Tanaka, T. Biochemical and developmental alterations of *Heliothis virescens* (F.) (Lepidoptera, Noctuidae) larvae induced by the endophagous parasitoid *Cardiochiles nigriceps* (Hymenoptera, Braconidae). *Arch. Insect Biochem. Physiol.* **1994**, *26*, 211–233.
33. Pennacchio, F.; Sordetti, R.; Falabella, P.; Vinson, S.B. Biochemical and ultrastructural alterations in prothoracic glands of *Heliothis virescens* (F.) (Lepidoptera: Noctuidae) last instar larvae parasitized by *Cardiochiles nigriceps* Viereck (Hymenoptera: Braconidae). *Insect Biochem. Molec. Biol.* **1997**, *27*, 439–450.
34. Pennacchio, F.; Falabella, P.; Sordetti, R.; Varricchio, P.; Malva, C.; Vinson, S.B. Prothoracic gland inactivation in *Heliothis virescens* (F.) (Lepidoptera: Noctuidae) larvae parasitized by *Cardiochiles nigriceps* Viereck (Hymenoptera: Braconidae). *J. Insect Physiol.* **1998**, *44*, 845–857.
35. Tanaka, T.; Vinson, S.B. Depression of prothoracic gland activity of *Heliothis virescens* by venom and calix fluid from the parasitoid, *Cardiochiles nigriceps*. *J. Insect Physiol.* **1991**, *37*, 139–144.
36. Pennacchio, F.; Vinson, S.B.; Tremblay, E.; Ostuni, A. Alteration of ecdysone metabolism in *Heliothis virescens* (F.) (Lepidoptera: Noctuidae) larvae induced by *Cardiochiles nigriceps* Viereck (Hymenoptera: Braconidae) teratocytes. *Insect Biochem. Mol. Biol.* **1994**, *24*, 383–394.

37. Falabella, P.; Caccialupi, P.; Varricchio, P.; Malva, C.; Pennacchio, F. Protein Tyrosine Phosphatases of *Toxoneuron nigriceps* bracovirus as potential disrupters of host prothoracic glands. *Arch. Insect Biochem. Physiol.* **2006**, *61*, 157–169.
38. Salvia, R.; Nardiello, M.; Scieuzo, C.; Scala, A.; Bufo, S.A.; Rao, A.; Vogel, H.; Falabella, P. Novel Factors of Viral Origin Inhibit TOR Pathway Gene Expression. *Front Physiol.* **2018**, *9*, 1678.
39. Ferrarese, R.; Brivio, M.; Congiu, T.; Falabella, P.; Grimaldi, A.; Mastore, M.; Perletti, G.; Pennacchio, F.; Sciacca, L.; Tettamanti, G.; et al. Early suppression of immune response in *Heliothis virescens* larvae by the endophagous parasitoid *Toxoneuron nigriceps*. *Invert. Surv. J.* **2005**, *2*, 60–68.
40. Pennacchio, F.; Vinson, S.B.; Tremblay, E. Morphology and ultrastructure of the serosal cells (teratocytes) in *Cardiochiles nigriceps* Viereck (Hymenoptera: Braconidae) embryos. *Int. J. Insect Morphol. Embryol.* **1994**, *23*, 93–104.
41. Pennacchio, F.; Vinson, S.B.; Tremblay, E. Host regulation effects of *Heliothis virescens* (F.) larvae induced by teratocytes of *Cardiochiles nigriceps* Viereck (Lepidoptera, Noctuidae—Hymenoptera, Braconidae). *Arch. Insect Biochem. Physiol.* **1992**, *19*, 177–192.
42. Pennacchio, F.; Vinson, S.B.; Malva, C. Regulation of host endocrine system by the endophagous braconid *Cardiochiles nigriceps* and its polydnavirus. In *Endocrine Interactions of Insect Parasites and Pathogens*; Edwards, J.P., Weaver, R.J., Eds.; BIOS Scientific Publishers: Oxford, UK, 2001; pp. 123–132.
43. Grimaldi, A.; Caccia, S.; Congiu, T.; Ferrarese, R.; Tettamanti, G.; Rivas-Pena, M.; Perletti, G.; Valvassori, R.; Giordana, B.; Falabella, P.; et al. Structure and function of the extraembryonic membrane persisting around the larvae of the parasitoid *Toxoneuron nigriceps*. *J. Insect Physiol.* **2006**, *52*, 870–880.
44. Falabella, P. The mechanism utilized by *Toxoneuron nigriceps* in inhibiting the host immune system. *Invert. Surviv. J.* **2018**, *15*, 240–255.
45. Varricchio, P.; Falabella, P.; Sordetti, R.; Graziani, F.; Malva, C.; Pennacchio, F. *Cardiochiles nigriceps* polydnavirus: Molecular characterization and gene expression in parasitized *Heliothis virescens* larvae. *Insect Biochem. Mol. Biol.* **1999**, *29*, 1087–1096.
46. Lapointe, R.; Wilson, R.; Vilaplana, L.; O'reilly, D.R.; Falabella, P.; Douris, V.; Bernier-Cardou, M.; Pennacchio, F.; Iatrou, K.; Malva, C.; et al. Expression of a *Toxoneuron nigriceps* polydnavirus-encoded protein causes apoptosis-like programmed cell death in lepidopteran insect cells. *J. Gen. Virol.* **2005**, *86*, 963–971.
47. Davies, D.H.; Vinson, S.B. Passive evasion by eggs of braconid parasitoid *Cardiochiles nigriceps* of encapsulation in vitro by hemocytes of host *Heliothis virescens*. Possible role for fibrous layer in immunity. *J. Insect Physiol.* **1986**, *32*, 1003–1010.
48. Hu, J.; Zhu, X.X.; Fu, W.J. Passive evasion of encapsulation in *Macrocentrus cingulum* Brischke (Hymenoptera: Braconidae), a polyembryonic parasitoid of *Ostrinia furnacalis* Guenee (Lepidoptera: Pyralidae). *J. Insect Physiol.* **2003**, *49*, 367–375.
49. Teng, Z.; Wu, H.; Ye, X.; Xiong, S.; Xu, G.; Wang, F.; Fang, Q.; Ye, G. An Ovarian Protein Involved in Passive Avoidance of an Endoparasitoid to Evade Its Host Immune Response. *J. Proteome Res.* **2019**, *18*, 2695–2705.
50. Vinson, S.B.; Guillot, F.S.; Hays, D.B. Rearing of *Cardiochiles nigriceps* in the laboratory, with *Heliothis virescens* as hosts. *Ann. Entomol. Soc. Am.* **1973**, *66*, 1170–1172.
51. Vanderzant, E.S.; Richardson, C.D.; Fort, S.W., Jr. Rearing of the bollworm on artificial diet. *J. Econ. Entomol.* **1962**, *55*, 140.
52. Beckage, N.E.; Gelman, D.B. Wasp parasitoid disruption of host development: Implication for new biologically based strategies for insect control. *Annu. Rev. Entomol.* **2004**, *49*, 299–330.
53. Schmidt, O. Insect Immune Recognition and Suppression. In *Insect Immunology*; Beckage, N.E., Ed.; Academic Press: Cambridge, MA, USA, 2008; pp. 271–294.
54. Mrinalini; Werren, J.H. Parasitoid Wasps and Their Venoms. In *Evolution of Venomous Animals and Their Toxins*; Gopalakrishnakone, P., Malhotra, A., Eds.; Toxinology; Springer: Berlin/Heidelberg, Germany, 2015; pp. 1–26.
55. Moreau, S.J.; Asgari, S. Venom Proteins from Parasitoid Wasps and Their Biological Functions. *Toxins* **2015**, *7*, 2385–2412.
56. Teng, Z.W.; Xiong, S.J.; Xu, G.; Gan, S.Y.; Chen, X.; Stanley, D.; Yan, Z.C.; Ye, G.Y.; Fang, Q. Protein Discovery: Combined Transcriptomic and Proteomic Analyses of Venom from the Endoparasitoid *Cotesia chilonis* (Hymenoptera: Braconidae). *Toxins* **2017**, *9*, 135.
57. Volkoff, A.N.; Cusson, M. The Unconventional Viruses of Ichneumonid Parasitoid Wasps. *Viruses* **2020**, *12*, 1170.
58. Strand, M.R. Chapter 12—Polydnavirus Gene Products that Interact with the Host Immune System. In *Parasitoid Viruses*; Beckage, N.E., Drezen, J.M., Eds.; Academic Press: Cambridge, MA, USA, 2012; pp. 149–161.
59. Bitra, K.; Zhang, S.; Strand, M.R. Transcriptomic profiling of *Microplitis demolitor* bracovirus reveals host, tissue and stage-specific patterns of activity. *J. Gen. Virol.* **2011**, *92*, 2060–2071.
60. Johner, A.; Lanzrein, B. Characterization of two genes of the polydnavirus of *Chelonus inanitus* and their stage-specific expression in the host *Spodoptera littoralis*. *J. Gen. Virol.* **2002**, *83*, 1075–1085.
61. Falabella, P.; Varricchio, P.; Gliotti, S.; Tranfaglia, A.; Pennacchio, F.; Malva, C. *Toxoneuron nigriceps* polydnavirus encodes a putative aspartyl protease highly expressed in parasitised host larvae. *Insect Mol. Biol.* **2003**, *12*, 9–17.
62. Provost, B.; Varricchio, P.; Arana, E.; Espagne, E.; Falabella, P.; Hugué, E.; La Scaleia, R.; Cattolico, L.; Poirié, M.; Malva, C.; et al. Bracovirus contain a large multigene family coding for a protein tyrosine phosphatase. *J. Virol.* **2004**, *78*, 13090–13103.
63. Dupas, S.; Wanjiru Gitau, C.; Branca, A.; Le Rü, B.P.; Silvain, J.F. Evolution of a Polydnavirus Gene in Relation to Parasitoid–Host Species Immune Resistance. *J. Hered.* **2008**, *99*, 491–499.
64. Desjardins, C.A.; Gundersen-Rindal, D.E.; Hostetler, J.B.; Tallon, L.J.; Fadrosch, D.W.; Fuester, R.W.; Pedroni, M.J.; Haas, B.J.; Schatz, M.C.; Jones, K.M.; et al. Comparative genomics of mutualistic viruses of *Glyptapanteles* parasitic wasps. *Genome Biol.* **2008**, *9*, R183.

65. Legeai, F.; Santos, B.; Robin, S.; Bretaudeau, A.; Dikow, R.; Lemaitre, C.; Jouan, V.; Ravallec, M.; Drezen, J.M.; Tagu, D.; et al. Conserved and specific genomic features of endogenous polydnviruses revealed by whole genome sequencing of two ichneumonid wasps. 2019, in press.
66. Carton, Y.; Poirié, M.; Nappi, A. Insect immune resistance to parasitoids. *Insect Science* **2008**, *15*, 67–87.
67. Webb, B.A.; Luckhart, S. Evidence for an early immunosuppressive role for related *Campoletis sonorensis* venom and ovarian proteins in *Heliothis virescens*. *Arch. Insect Biochem. Physiol.* **1994**, *26*, 147–163.
68. Vincent, B.; Kaeslin, M.; Roth, T.; Heller, M.; Poulain, J.; Cousserans, F.; Schaller, J.; Poirié, M.; Lanzrein, B.; Drezen, J.; et al. The venom composition of the parasitic wasp *Chelonus inanitus* resolved by combined expressed sequence tags analysis and proteomic approach. *BMC Genom.* **2010**, *11*, 693.
69. Goecks, J.; Mortimer, N.T.; Mobley, J.A.; Bowersock, G.J.; Taylor, J.; Schlenke, T.A. Integrative approach reveals composition of endoparasitoid wasp venoms. *PLoS ONE* **2013**, *8*, e64125.
70. Yan, Z.C.; Fang, Q.; Wang, L.; Liu, J.; Zhu, Y.; Wang, F.; Li, F.; Werren, J.H.; Ye, G. Insights into the venom composition and evolution of an endoparasitoid wasp by combining proteomic and transcriptomic analyses. *Sci. Rep.* **2016**, *6*, 19604.
71. Yang, L.; Lin, Z.; Fang, Q.; Wang, J.; Yan, Z.; Zou, Z.; Song, Q.; Ye, G. Identification and characterization of serine protease inhibitors in a parasitic wasp, *Pteromalus puparum*. *Sci. Rep.* **2017**, *7*, 15755.
72. Asgari, S.; Schmidt, O. Passive protection of eggs from parasitoid, *Cotesia rubecula*, in the host *Pieris rapae*. *J. Insect Physiol.* **1994**, *40*, 789–795.
73. Beckage, N.E. Parasitoids and polydnviruses. *Bioscience* **1998**, *48*, 305–311.
74. Han, L.B.; Yin, L.H.; Huang, L.Q.; Wang, C.Z. Differential immunosuppression by *Campoletis chloridae* eggs and ichnovirus in larvae of *Helicoverpa armigera* and *Spodoptera exigua*. *J. Invertebr. Pathol.* **2015**, *130*, 88–96.
75. Yin, C.; Li, M.; Hu, J.; Lang, K.; Chen, Q.; Liu, J.; Guo, D.; He, K.; Dong, Y.; Luo, J.; et al. The genomic features of parasitism, Polyembryony and immune evasion in the endoparasitic wasp *Macrocentrus cingulum*. *BMC Genom.* **2018**, *19*, 420.
76. Luckhart, S.; Webb, B.A. Interaction of a wasp ovarian protein and polydnvirus in host immune suppression. *Dev. Comp. Immunol.* **1996**, *20*, 1–21.
77. Bras, M.; Queenan, B.; Susin, S.A. Programmed Cell Death via Mitochondria: Different Modes of Dying. *Biochemistry* **2005**, *70*, 231–239.
78. Gourlay, C.W.; Ayscough, K.R. A Role for Actin in Aging and Apoptosis. *Biochem. Soc. Trans.* **2005**, *33*, 1260–1264.
79. Redza-Dutordoir, M.; Averill-Bates, D.A. Activation of apoptosis signalling pathways by reactive oxygen species. *Biochim. Biophys. Acta.* **2016**, *1863*, 2977–2992.
80. Li, T.S.; Marbán, E. Physiological levels of reactive oxygen species are required to maintain genomic stability in stem cells. *Stem Cells* **2010**, *28*, 1178–1185.
81. Chen, Y.; Zhou, Z.; Min, W. Mitochondria, Oxidative Stress and Innate Immunity. *Front. Physiol.* **2018**, *9*, 1487.
82. Stefani, M.; Dobson, C.M. Protein aggregation and aggregate toxicity: New insights into protein folding, misfolding diseases and biological evolution. *J. Mol. Med.* **2003**, *81*, 678–699.
83. Milkovic, L.; Cipak, G.A.; Cindric, M.; Mouthuy, P.A.; Zarkovic, N. Short Overview of ROS as Cell Function Regulators and Their Implications in Therapy Concepts. *Cells* **2019**, *8*, 793.
84. Li, Y.; Lu, J.F.; Feng, C.J.; Ke, X.; Fu, W.J. Role of venom and ovarian proteins in immune suppression of *Ostrinia furnacalis* (Lepidoptera: Pyralidae) larvae parasitized by *Macrocentrus cingulum* (Hymenoptera: Braconidae), a polyembryonic parasitoid. *Insect Sci.* **2007**, *14*, 93–100.
85. Choi, J.Y.; Whitten, M.M.; Cho, M.Y.; Lee, K.Y.; Kim, M.S.; Ratcliffe, N.A.; Lee, B.L. Calreticulin enriched as an early-stage encapsulation protein in wax moth *Galleria mellonella* larvae. *Dev. Comp. Immunol.* **2002**, *26*, 335–343.
86. Asgari, S.; Schmidt, O. Is cell surface calreticulin involved in phagocytosis by insect hemocytes? *J. Insect. Physiol.* **2003**, *49*, 545–550.
87. Zhu, J.Y.; Fang, Q.; Wang, L.; Hu, C.; Ye, G.Y. Proteomic analysis of the venom from the endoparasitoid wasp *Pteromalus puparum* (Hymenoptera: Pteromalidae). *Arch. Insect Biochem. Physiol.* **2010**, *75*, 28–44.
88. Wang, L.; Fang, Q.; Qian, C.; Wang, F.; Yu, X.Q.; Ye, G. Inhibition of host cell encapsulation through inhibiting immune gene expression by the parasitic wasp venom calreticulin. *Insect Biochem. Mol. Biol.* **2013**, *43*, 936–946.
89. Teng, Z.W.; Xu, G.; Gan, S.Y.; Chen, X.; Fang, Q.; Ye, G.Y. Effects of the endoparasitoid *Cotesia chilonis* (Hymenoptera: Braconidae) parasitism, venom, and calyx fluid on cellular and humoral immunity of its host *Chilo suppressalis* (Lepidoptera: Crambidae) larvae. *J. Insect Physiol.* **2016**, *85*, 46–56.