



Astaxanthin anticancer effects are mediated through multiple molecular mechanisms: A systematic review

Immacolata Faraone^{a,b,1}, Chiara Sinisgalli^{a,b,1}, Angela Ostuni^{a,b}, Maria Francesca Armentano^{a,b},
Monica Carmosino^{a,b}, Luigi Milella^{a,*}, Daniela Russo^{a,b}, Fabiana Labanca^{a,*}, Haroon Khan^c

^a Dipartimento di Scienze, Università della Basilicata, Via dell'Ateneo Lucano, 10 85100 Potenza, Italy

^b BioActiPlant s.r.l., Via dell'Ateneo lucano, 10 85100, Potenza, Italy

^c Department of Pharmacy, Abdul Wali Khan University, Mardan, 23200, Pakistan

ARTICLE INFO

Keywords:

Proapoptotic
Cancer prevention
Cell cycle
Malignancy
Survivina
Staxanthin

ABSTRACT

During the latest decades, the interest on the effectiveness of natural compounds and their impact on human health constantly increased, especially on those demonstrating to be effective on cancer. Molecules coming from nature are currently used in chemotherapy like Taxol, Vincristine or Vinblastine, and several other natural substances have been showed to be active in reducing cancer cell progression and migration. Among them, astaxanthin, a xanthophyll red colored carotenoid, displayed different biological activities including, anti-inflammatory, antioxidant, proapoptotic, and anticancer effects. It can induce apoptosis through downregulation of antiapoptotic protein (Bcl-2, p-Bad, and survivin) expression and upregulation of proapoptotic ones (Bax/Bad and PARP). Thanks to these mechanisms, it can exert anticancer effects towards colorectal cancer, melanoma, or gastric carcinoma cell lines. Moreover, it possesses antiproliferative activity in many experimental models and enhances the effectiveness of conventional chemotherapeutic drugs on tumor cells underling its potential future use. This review provides an overview of the current knowledge on the anticancer potential of astaxanthin by modulating several molecular targets. While it has been clearly demonstrated its multitarget activity in the prevention and regression of malignant cells in *in vitro* or in preclinical investigations, further clinical studies are needed to assess its real potential as anticancer in humans.

1. Introduction

Cancer is the second-leading cause of death in the world, by generation of an uncontrollable proliferation of abnormal cells able to grow

in situ or also able to overrun contiguous tissues and to extent to other organs in the body, through the circulatory or lymphatic systems [1]. GLOBOCAN estimated that the incidence and mortality worldwide are continuously rising year by year reaching 18.1 million new cases and

Abbreviations: Akt, protein kinase B; anti-PCNA, proliferating cell nuclear antigen; Bax, Bcl-2 associated protein; Bcl-2, B-cell lymphoma 2; BDNF, brain-derived neurotrophic factor; bw, body weight; CdC2, cell division cycle 2; CDK, cyclin-dependent kinase; COX-2, cyclooxygenase-2; DMH, dimethyl hydrazine; ECM, extracellular matrix; EMT, epithelial-mesenchymal transition; ERK, extracellular signal-regulated kinases; FAK, focal adhesion kinase; Fas-L, Fas ligand; FoxO3, Forkhead box O3; GSK-3 β , Glycogen synthase kinase 3; GSTM2, glutathione S-transferase M2; HAS, human serum albumin; HDL, high-density lipoprotein; hENT1, human equilibrative nucleoside transporter; HO-1, Heme oxygenase; HPV, human papilloma virus; HSV, herpes simplex virus; IAP, inhibitor of apoptosis proteins; IKK β , inhibitor of nuclear factor kappa-B kinase; IL-, Interleukin-; JAK1, Janus kinase 1; MAPK, mitogen-activated protein kinase; miRNAs, microRNA; MMP, matrix metallo proteinases; mTOR, Mammalian target of rapamycin; NF-kB, nuclear factor kappa-light-chain-enhancer of activated B cells; NMBA, N-nitrosomethylbenzylamine; NQO1, NAD(P)H quinone oxidoreductase 1; Nrf2/ARE, nuclear factor-erythroid 2-related factor 2/antioxidant response elements; p70S6K, p70 ribosomal protein S6 kinase; pBCEC, porcine brain capillary endothelial cells; pCNA, proliferating cell nuclear antigen; PI3K, phosphatidylinositol 3-kinases; PPAR- γ , peroxisome proliferator-activated receptor; PSA, Prostate Specific Antigen; ROS, oxygen reactive species; RRM1-2, ribonucleotide reductase subunit M1 and M2; RTKs, receptor tyrosine kinases; SIRT1, sirtuin 1; SOD, superoxide dismutase; STAT3, Signal transducer and activator of transcription 3; TGF-, transforming growth factor-; TLR-4, toll-like receptors 4; TNF, tumor necrosis factor; TRAIL, TNF-related apoptosis-inducing ligand; VEGF, Vascular Endothelial Growth Factor; VEGFR2, Vascular Endothelial Growth Factor receptor; Wnt, Wingless-INT; XPC, xeroderma pigmentosum complementation group C; ZEB1, Zinc finger E-box binding homeobox 1

* Corresponding authors.

E-mail addresses: luigi.milella@unibas.it (L. Milella), fabianalabanca@outlook.it (F. Labanca).

¹ These authors contributed equally to this work.

<https://doi.org/10.1016/j.phrs.2020.104689>

Received 28 November 2019; Received in revised form 4 February 2020; Accepted 10 February 2020

Available online 11 February 2020

1043-6618/ © 2020 Elsevier Ltd. All rights reserved.

9.6 million deaths in 2018. Moreover, the incidence to develop cancer in men is higher than women (one in 5 men and one in 6 women worldwide) as well as the mortality (one in 8 men and one in 11 women) [2]. Worldwide, it is estimated to be 43.8 million the number of people alive within 5 years of a cancer diagnosis (the 5-year prevalence). For decades, surgery, chemotherapy, radiotherapy and molecular target therapy represented the only antitumoral strategies available. Nowadays, significant efforts have revolutionized these traditional cancer treatments with the advent of promising strategies as cancer immunotherapy and precision medicine. Nevertheless, chemotherapy is still widely used and the importance of chemical compounds, in particular coming from natural sources, is widely recognized, like Taxol, Vincristine and Vinblastine. Moreover, the interest on the comprehension of the mechanistic aspects that lead to the beginning and the growth of cancer is increasing [3–5].

Natural products are gaining success in our society and almost half of the best-selling pharmaceuticals were derived or inspired from nature. Several testing procedures as preclinical, clinical phases I to III, and preregistration, included 547 FDA-approved natural products and derivatives. They derived mainly from plants (47 %), bacteria (30 %), and fungi (23 %) [6].

Among natural substances, several have been demonstrated to act specifically on biological targets involved in prevention or treatment of malignant cell as astaxanthin (3,3'-dihydroxy- β , β '-carotene-4,4'-dione) (Fig. 1). It was discovered for the first time in lobsters, in 1938, and it was employed in aquaculture for the pigmentation. In 1991, it gained approval as a supplement for food, thanks to its antioxidant features, biological and physiological activities [7]. The polar-nonpolar-polar layout allows astaxanthin to precisely fit into the cell membrane [8,9]. Moreover, the hydroxyl (OH) and the carbonyl (C = O) present in each ionic ring of astaxanthin make the molecule more polar, esterifiable, and highly antioxidant [10].

In particular, astaxanthin is one of the most important terpene unsaturated compound [11] derived from β -carotene by 3-hydroxylation and 4-ketolation, catalysed by β -carotene hydroxylase and β -carotene, in both ionone end groups, changing the colour from yellow to red [12]. This reaction occurs naturally only in living organisms, such as few

bacteria, fungi, and some unicellular green algae, because the ketolation is limited to these species [12,13]. There are many different natural molecular isomers of astaxanthin with unique three-dimensional shape (Fig. 1). In detail, the presence of two stereogenic carbon atoms in C3 and C3' position give three stereoisomers for astaxanthin: a pair of enantiomers (3S,3'S- and 3R,3'R-astaxanthin) and an optically inactive *meso* form (3R,3'S astaxanthin) [14,15]. Given the higher stability, the predominant natural astaxanthin form usually used as a dietary supplement and in clinical trials is the all-*trans* 3S,3'S astaxanthin [8,15]. Nowadays, a large proportion of astaxanthin is synthetically produced starting from the ketoisophorone obtained from petroleum [10]. Only a small part of commercial astaxanthin is extracted from *Haematococcus pluvialis* and *Xanthophyllomyces dendrorhous*, providing the bioactive 3S,3'S stereoisomer [12]. In astaxanthin-producing organisms, the free form of the molecule is uncommon because it is particularly unstable and susceptible to oxidation; in fact, the most common form of astaxanthin is present as acyl monoesters or diesters (esterified with one or two fatty acids such as palmitic, oleic, stearic or linoleic acid) or conjugated with proteins or lipoproteins. In *H. pluvialis*, the 99 % of astaxanthin is as acyl monoesters and diesters, which make up astaxanthin currently available in commercial field [8].

Indeed, the microalgae *H. pluvialis* represents the primary natural source of astaxanthin containing up to 4% of the total cellular dry weight of 3S,3'S isomer. It is commonly used for human applications such as food and beverages, dietary supplements, and cosmetics; in fact, in humans, astaxanthin intake derives almost exclusively from seafood [8]. Other natural sources are represented by *X. dendrorhous* (*Phaffia rhodozyma*) and *Zantophyllumiss dendrorhousis* yeasts, *Chlorococcum* sp., *Chlorella zofingiensis* algae, marine *Agrobacterium aurantiacum* bacterium and other marine organisms such as salmon, lobster, crab and shrimp [7]. On the other hand, the synthetic form of molecule obtained from yeast (mutated *X. dendrorhous*) and bacteria sources (aerobic bacteria *Paracoccus carotinifaciens*) are predominantly used in the aquaculture sector [16]. Astaxanthin displayed several interesting biological activities including, proapoptotic, anticancer, antioxidant, and antiinflammatory effects, alleviating endothelial dysfunction, but its role in the prevention, effectiveness and regression on malignant cells is




Fig. 1. Astaxanthin and isomers.




Fig. 2. Flow diagram of the systematic review literature search results based on PRISMA statement.

still under investigation [9].

In several cancers there is a deregulation of different pathways, these are responsible of cancer cell hyperproliferation, diffusion and consequently on their subsistence. In cancer, several aberrant signaling pathways contribute to its pathology, and a multi-targeted treatment could be able to modulate different pathways and up or downregulate different genes [17]. Biologically active natural products are able to target several signaling pathways simultaneously, showing advantageous in cancer therapy.

This review provides a detailed and comprehensive knowledge of the recent achievements in the field of astaxanthin mechanistic effects on the different phases of cancer prevention, malignant cell transformation and therapy analyzing the available research papers available up to now (Fig. 1).

2. Study analysis

The diagram describes the selection process of the bibliographic sources (Fig. 2). The initial search provided 2285 studies (1390 from Scopus and 895 from PubMed), out of which 1511 were duplicates among all keywords used. Out of those 774 studies, 619 were off topic basing on the exclusion criteria. Out of those 154, 113 do not contain experimental data about effective concentration. The final reference list of 41 articles comprises 6 major countries [China 24.5 %, Japan 20.4 %, India 18.4 %, Usa 12.2 %, South Korea 10.2 %, others (Iran, Italy, Malaysia, Taiwan, Thailand) 14.3 %] as shown in Fig. 3. Statistical analysis showed that the considered references have been published




Fig. 3. Representation of distribution of author origin country.




Fig. 4. Number of works published on astaxanthin and cancer.

mostly in the last few years (Fig. 4). Astaxanthin anticancer effects have been evaluated *in vitro* on several cell lines. Among them, eight cell lines belong to colon carcinoma, and HCT116 were the most studied ($n = 3$). Six different hepatocarcinoma cell lines have been studied, while no *in vitro* studies have been reported on tongue and urinary cancer. Regarding these tumors, 2 studies on rats and 1 study on mice have been analyzed. Moreover, *in vivo* studies regard breast carcinoma ($n = 2$), colon carcinoma ($n = 2$), and hepatocarcinoma ($n = 3$). Our systematic review included 27 *in vitro* studies and 14 *in vivo* reports. *In vitro* studies included biochemical and molecular analysis, comprising colorimetric and enzymatic assays, western blot, flow cytometry analysis, and immunofluorescence techniques. *In vivo* experiments were carried out on animal model as mice, rats and hamster in order to determine tumor weight, metastasis formation, and apoptosis index. Regarding the methodological quality assessment, the *in vivo* studies were selected via standard checklist for preclinical trials [18]. As reported in Fig. 5, all studies describing the objectives, outcomes and main findings have been considered in this review. In addition, doses, routes of administration and frequency were cited, but none of the studies showed sample size calculations and no information on blinding was reported. Studies on randomization of animals, blinding and reported outcome measurements were generally considered of higher methodological quality. Limitations of these studies are due to the lack of these important information and the unclear results; the experimental design of some animal studies might be improved.

To better describe and understand *in vivo* and *in vitro* results of




Fig. 5. Methodological quality assessment of included studies. Dark gray bars indicate the proportion of articles that met each criterion; light gray bars indicate the proportion of studies that did not and white bars indicate the proportion of studies with unclear answers.

anticancer properties of astaxanthin, we summarized them as reported below.

3. Astaxanthin pharmacological properties/anti-cancer effects *in vitro*

It is broadly accepted that elevated inflammation levels, increased oxidative stress, aging, as well as mitochondrial dysfunctions are responsible for major chronic degenerative diseases [19,20]. Due to its antioxidant potential, astaxanthin has gained considerable importance as a valuable therapeutic candidate. It shows a wide field of clinical applications, sharing the same oxidative and inflammatory etiology. Among them, the most relevant are neuroprotection, gastric ulcer care, stroke injury amelioration, kidney and liver protection, which are strictly related with cancer prevention. In fact, astaxanthin treatment reduces the release of interleukins, COX-2 and nitric oxide in lipopolysaccharide-stimulated BV2 microglia [21] and pretreatment with astaxanthin (50 μM) improves stretch injury in astrocyte model by reducing inflammatory mediators [22]. The subcellular location of astaxanthin revealed its preference for the mitochondrial membrane, suggesting its role in modulating cytochrome *c* and pro-apoptotic factors release and, thus, the protective role against oxidative stress. Astaxanthin, indeed, by activating PPAR- γ and catalase, decreases mitochondrial and intracellular oxygen ROS levels and IL-8 expression in gastric epithelial cells [23]. The ability of astaxanthin in the modulation of many inflammatory-related pathways is responsible of its preventive effects on ischemia-reperfusion injury in steatotic liver cells [24], as well as against liver fibrosis in human LX-2 cells [25]. Furthermore, the activation of Nrf2/ARE signaling by astaxanthin explains the marked attenuation of renal fibrosis in glomerular mesangial cells [26], increasing the transcription of the activity of the phase II detoxifying/antioxidant enzymes activities, such as NAD(P)H, NQO1, GSTM2, HO-1 and SOD and reducing the incidence of neoplastic transformation [1]. Moreover, it has been demonstrated that astaxanthin attenuates acute kidney injury and reduces apoptosis in tubular epithelial cells, through the inhibition of the histone deacetylase SIRT1 [27] and the related FoxO3 transcription factor [28]. Astaxanthin may also improve the nephropathy-related complications, as well as peritoneal fibrosis, limiting epithelial-mesenchymal transition, thanks to the anti-inflammatory, antioxidative and mitochondrial ROS scavenging activities in peritoneal mesothelial cells [29]. The enhancement of the antioxidant defense system, as well as superoxide dismutase and catalase, has been also demonstrated [30].

Given this evidence, it is easily intuible that astaxanthin can exert

anti-cancer effects, as cancer stems from an unbalance in oxidative status, starting with a trigger inflammatory lesion. Oxidative stress, indeed, is strictly linked to carcinogenesis, due to the related cellular damage that can occur, deriving from DNA mutations and protein aberrations. Since ever, the chemopreventive properties of natural substances have been recognized. In particular, since the '90 s years, the researchers focused the attention on the relationship between natural xanthophylls, like astaxanthin, and cancer, demonstrating protective effects against well-known cancerogenic B1 aflatoxin, through the activation of detoxification mechanisms [31] and against tongue and large bowel carcinogenesis [32]. These properties have been mainly ascribed to the radical scavenging activity [33] and reduction of superoxide and nitric oxide generation [34] back then. It has been demonstrated that astaxanthin exhibits higher antioxidant properties rather than α -tocopherol and β -carotene [35] and reduces DNA damage deriving from UVA radiation exposure [36]. Evidence of the decrease of oxidative markers at doses of 10 nM, in rat kidney fibroblasts, in 1998, has led to the study of this carotenoid as candidate to contrast skin photodamage from the past [35] to date [37], also considering its non-toxicity on normal human dermal fibroblasts [38], probably due to the non-interaction with tyrosinase activity [39]. By contrast, astaxanthin demonstrated to promote the reepithelization of wounds, by increasing the number and the recruiting of keratinocytes in a dose-dependent manner [40]. Recently, researchers have been focused on the role of astaxanthin towards skin cancer, finding that it may inhibit the neoplastic transformation of mouse epidermal cell line JB6 P + cells, with a decrease of cell viability by up to 94 % after five day-treatment with 50 μM of astaxanthin [41]. These results confirmed the study of Chen, et al. [42] on two melanoma cell lines, A375 and A2058, who showed inhibition of proliferation with 168 μM astaxanthin by up 50 % and 80 %, respectively. The antiproliferative activity of astaxanthin extends to multiple cancer lines, including human hepatocarcinoma (CBRH-7919), rat breast cancer cells (SHZ-88), and mouse Lewis lung carcinoma cells, with higher sensitivity for the first cell line, with IC₅₀ values around 39 μM [43]. As regards this last tumor type, astaxanthin can suppress cancer proliferation also in two human non-small cancer lung cells, bronchioloalveolar cell carcinoma (A549), and squamous cell carcinoma (H1703), reducing, with a concentration of 20 μM , the cell colony-forming ability by 50.29 % and 39.71 %, respectively [44]. The beneficial implications of astaxanthin on liver cancer development have been also investigated in other human hepatoma cell lines, reporting that astaxanthin significantly inhibits cell growth in HepG2 cell line at a concentration of 42 μM [45], while in LM3 and SMMC-7721 with IC₅₀ = 100 μM [46]. However, all reported effects are dose- and time-




Fig. 6. Multiple mechanisms of astaxanthin for cancer therapy.

dependent, as it is clearly evident in a study conducted on hepatoma cells (H22), in which low- (20 μM) and high-dose (40 μM) of astaxanthin have been tested after 12 and 24 h, showing that the inhibitory effect is more pronounced in high-dose group and it increases with time. Interestingly, the lowest dose of astaxanthin (20 μM) inhibited cell growth with the same efficacy of cisplatin, a conventional anticancer drug, used as control [47]. By *in silico* molecular docking, Suganya and Anuradha [48] demonstrated that astaxanthin can reach higher scores compared with that obtained with a standard chemotherapeutic agent, sorafenib, in hepatocellular carcinoma. Previous results report time- (24 – 72 h) and dose-dependent (4 – 16 μM) antiproliferative properties of astaxanthin against HCT116 and HT-29 [49], as well as of the entire *H. pluvialis* extract (5–25 $\mu\text{g}/\text{mL}$). This extract showed 100 % inhibition on colon cancer cell viability at concentrations of 0.1 mg/mL, with a lethal concentration in 50 % of cells of 27.03 $\mu\text{g}/\text{mL}$ lower than that of doxorubicin (37.6 $\mu\text{g}/\text{mL}$), used as control [50]. At the same time, these studies deny earlier results that do not ascribe a strong inhibitory potential to astaxanthin ($\text{IC}_{50} > 50\mu\text{M}$), when compared to that of other natural antioxidants, like curcumin ($\text{IC}_{50} = 17.6 \mu\text{M}$) and resveratrol ($\text{IC}_{50} = 15.8 \mu\text{M}$) [51]. The toxicity of astaxanthin on normal human oral keratinocytes (NOKs) has been investigated in parallel with the chemopreventive effects on colorectal adenocarcinoma cells (Caco-2), concluding that concentration of astaxanthin between 8–16 mM may inhibit Caco-2 cells proliferation, being at the same time not harmful to NOKs [52]. Furthermore, even nanoemulsions containing 25 $\mu\text{g}/\text{mL}$ of astaxanthin may decrease by up to 50 % the cell viability of HT-29 and stomach gastric cancer (AGS) cell lines [53]. By contrast, another study reports that the viability of AGS was not affected by astaxanthin administration, whereas other gastric carcinoma cell lines, KATO-III and SNU-1, showed a dose-dependent decrease [54]. Further studies will be necessary to establish if that depends on the bioavailability of formulation or other mechanisms are involved. In 2 human esophageal cancer cell lines (TE-1, TE-4), concentrations of astaxanthin higher than 13 μM in TE-1 cell line and between 42–84 μM in TE-4 inhibited cancer cell proliferation [55]. Moreover, in leukemia cells (K562), astaxanthin revealed the most powerful cell growth inhibitor among other tested carotenoids, with a strong activity even at low concentrations (5 and 10 μM) [56]. Even if at higher concentrations, 24 h of astaxanthin treatment may suppress the viability of oral squamous carcinoma (SCC131 and SCC4) cells with IC_{50} of 700 μM and 720 μM , respectively [57]. The multiple

mechanisms at the basis of the anticancer activity of astaxanthin also include epigenetic modifications and chromatin remodeling. In prostate cancer, the control of oncosuppressor gene silencing by DNA methylation is a key strategy to protect cells from carcinogenesis [58]. In addition, the prevention and treatment of prostate carcinoma pass through the suppression of 5- α -reductase enzyme. Many natural sources showed to inhibit this enzyme activity. Among them, astaxanthin is linked to a reduction of 98 % in 5- α -reductase activity at 502 μM *in vitro*, in association with a cancer growth inhibition of 24 % at 0.2 μM [59]. Although from first observation astaxanthin seemed not significantly suppress the cell viability of breast cancer (MCF-7 and MDA-MB-468) cells [60], further investigation has demonstrated the cytotoxic activity of *H. pluvialis* extract on breast adenocarcinoma [50] suggesting a possible synergistic effect. Another study evaluated the astaxanthin nanoparticle anticancer potential against MDA-MB-231, revealing $\text{IC}_{50} = 84 \mu\text{M}$ [61], while astaxanthin induces cell death in MCF-7 at concentrations of 33 mM [62]. These inconsistencies may be probably due to the synthetic origin of astaxanthin, which revealed 90 fold less potent than natural astaxanthin as demonstrated by Régnier, et al. [63] on human umbilical vein endothelial cells. However, astaxanthin has been recognized as breast cancer chemopreventive agent, [62], on the other hand, astaxanthin does not present toxicity on normal breast epithelial cells [64]. Astaxanthin also revealed an excellent adjuvant to conventional therapies, thanks to its capability to reduce irradiation-related cytogenetic effects on human peripheral blood lymphocytes at concentrations of 20.0 $\mu\text{g}/\text{mL}$ [65] and for its photothermal effect. Considering its chemical structure, indeed, xanthophylls may strongly absorb the light and convert into heat in cancer cells, as demonstrated in squamous cell carcinoma (VX2) and macrophage (246.7) cell lines. This treatment led to the reduction of adverse reactions and a more site-specific action [66]. In conclusion, astaxanthin has revealed anticancer activity by multiple mechanisms, as well as apoptosis induction, cell growth inhibition, and cell cycle interferences that will be clarified in the following sections (Fig. 6, Table 1).

4. Astaxanthin induces cell cycle arrest in tumor cells

Cell cycle machinery drives and controls cell proliferation, but what happens if something goes wrong? The elucidation of the potential mechanism involved in cell viability inhibition and the changes in cell cycle progression of cancer cells are an increased investigated topic.

Table 1Description of the main mechanisms involved in anticancer activity and effective dose in different *in vitro* and *in vivo* experimental models.

Cancer	<i>In vitro</i>		<i>In vivo</i>		Mechanisms	Ref.
	Cell line	Cytotoxic concentration	Experimental model	Dose		
Breast carcinoma	MCF-7	IC ₅₀ = 20 ± 0.2 μM			↓ cyclin D1; ↑ cell number in G0/G1 phase; ↓ migration; ↑ apoptosis; ↑ p53; ↓ Bax; ↓ Bcl-2; ↑ ROS	(64, 67, 94)
	MDA-MB-231	IC ₅₀ = 84 μM	Sprague-Dawley rats	50 % (45.5 ± 0.6 mg)	↑ apoptosis; ↓ migration ↑ adiponectin	(61) (143)
			Mice	25 mg (single dose)	↓ microelements in tumor site	(144)
Cervical adenocarcinoma	HeLa	IC ₅₀ ~ 50 μM			↑ cell number in G2/M phase	(80)
Colon carcinoma	HCT116	IC ₅₀ = 10-16 μM			↑ cell number in G2/M phase; ↑ p21; ↑ p27; ↑ p53; ↓ CDK4 and CDK6; ↑ apoptosis; ↑ caspase-3; ↑ PARP; ↑ EGFR; ↑ p-p38; ↑ p-JNK; ↑ pERK1/2; ↓ invadopodia; ↓ EMT; ↑ E-cadherin; ↓ vimentin; ↓ cortactin; ↓ MMP2; ↑ miR-29a-3p; ↓ ZEB1; ↓ MYC; ↓ cyclin D1	(49, 76, 108)
	CT-26	50 - 100 μM			↓ invadopodia; ↓ EMT; ↑ E-cadherin; ↓ vimentin; ↓ cortactin; ↓ MMP2; ↑ miR-29a-3p; ↓ ZEB1; ↓ MYC	(108)
	HT-29	IC ₅₀ = 10-16 μM			↓ cell viability	(49)
	CaCo-2	IC ₅₀ = 8.4 - 16.8 mM			↓ cell viability	(52)
	LS-174, WiDr, SW-480	53.0 ± 3.1 % (41.9 μM) 37.8 ± 2.1 % (41.9 μM) 40.5 ± 2.2 % (41.9 μM)			↓ cell viability	(76)
	LS-180	100 - 200 μM	Db/db mice	340 μM	↑ apoptosis; ↑ Bax; ↑ caspase 3; ↓ Bcl2 ↓ oxidative stress	(93) (146)
Esophageal carcinoma	ECA 109 T13	IC ₅₀ = 295.3 μM IC ₅₀ = 216.1 μM	Wistar male rats	15 mg/kg	↓ ERK-2; ↓ NFκB; ↓ COX-2 ↑ cell number in G2/M phase; ↓ cyclin B1; ↓ Cdc2; ↑ apoptosis	(147) (79)
	TE-1	IC ₅₀ > 13.4 μM			↓ cell proliferation	(55)
Gastric adenocarcinoma	TE-4	IC ₅₀ = 41.9 - 83.8 μM				
	KATO-III	IC ₅₀ ≤ 100 μM			↑ cell number in G0/G1 phase; ↓ p-ERK; ↑ p27 ^{Kip-1}	(54)
	SNU-1					
Hepatocarcinoma	HepG2	IC ₅₀ ~ 41.9 μM			↑ cell number in G0/G1 phase; ↑ apoptosis; ↑ caspase-3; ↑ p53; ↓ c-fos-c-jun	(45) (47)
	H22	IC ₅₀ = 40 μM			↑ cell number in G2 phase	(47)
	CBRH-7919	IC ₅₀ = 39 μM			↑ apoptosis; ↓ JAK1-STAT3; ↓ Bcl-2	(43)
	LM3	IC ₅₀ = 100 μM			↑ cell number in G2/M phase; ↑ apoptosis; ↓ NF-κB p65; ↓ Wnt/β-catenin	(46)
	SMMC-7721					
	AH109A	5 μM	KM mice	2 - 4 μg/kg	↓ invasion	(107)
			Db/db mice	340 μM	↓ tumor weight	(47)
			DBA mice	100 mg/kg (4days) 1 mg/kg (14days)	↓ oxidative stress; ↑ adiponectin ↓ lipid peroxidation; ↓ metastasis	(134) (135)
Leukemia	K562	IC ₅₀ = 25 μM			↓ metastasis; ↑ p21 stimulation; ↓ cyclin D1;	(56, 68, 69)
	HL-60				↑ PPARγ; ↑ cell number in G0/G1 phase; ↑ Nrf2; ↑ apoptosis	
Lung carcinoma	A549				↑ cell number in G0/G1 phase;	(44, 75)
	H1703				↑ p38 MAPK; ↑ apoptosis	
Oral squamous cell carcinoma	SCC131	IC ₅₀ = 700 μM			↑ p21 stimulation; ↓ cyclin D1;	(57)
	SCC4	IC ₅₀ = 720 μM			↓ angiogenesis (↓ VEGF, VEGF2, HIF-1α); ↑ apoptosis	
			Hamster	15 mg/kg bw	↓ tumor growth; ↓ PI3 K/Akt/NF-κB/STAT-3	(99, 103, 138)
Prostate cancer	LNcap-FGC	24 % (0.17 μM)	xenograft PC-3 nude mice	41.7 % (100 mg/kg bw)	↓ cancer cell growth ↑ apoptosis; ↑ cleaved caspase-3; ↑ miR-375 and miR-487b	(59) (142)

(continued on next page)

Table 1 (continued)

Cancer	<i>In vitro</i>		<i>In vivo</i>		Mechanisms	Ref.
Skin cancer	JB6 P+	91.4 % (50 μ M)			↑ Nrf2; ↑ cell number in G1 phase; ↓ MMP1, 2 and 9; ↓ migration; ↑ apoptosis	(41, 42)
	A375	50 % (167.5 μ M)				
	A2058	80 % (167.5 μ M)				
Skin cancer Tongue neoplasm			Mice	850 μ M	↓ tyrosine nitration	(139)
			Rats	200 μ g/kg bw	↓ tyrosine activity	(140)
Urinary bladder cancer			male F344 rats	170 μ M	↓ polyamine levels	(137)
			male ICR mice	85 μ M	↓ cancer cell growth	(141)

Akt, protein kinase B; Bax, Bcl-2 associated protein; Bcl-2, B-cell lymphoma 2; bw, body weight; CdC2, cell division cycle 2; CDK, cyclin-dependent kinase; COX-2, cyclooxygenase-2; ERK, extracellular signal-regulated kinases; IL-, Interleukin-; JAK1, Janus kinase 1; MAPK, mitogen-activated protein kinase; miRNAs, microRNA; MMP, matrix metallo proteinases; NF-kB, nuclear factor kappa-light-chain-enhancer of activated B cells; Nrf2/Are, nuclear factor-erythroid 2-related factor 2/antioxidant response elements; P13 K, phosphatidylinositol 3-kinases; PPAR- γ , peroxisome proliferator-activated receptor; ROS, oxygen reactive species; STAT3, Signal transducer and activator of transcription 3; TNF, tumor necrosis factor; VEGF, Vascular Endothelial Growth Factor; VEGFR2, Vascular Endothelial Growth Factor receptor; Wnt, Wingless-INT; ZEB1, Zinc finger E-box binding homeobox 1.

Astaxanthin induced the inhibition of cell proliferation by cell cycle arrest in various ways. Its effects might result in cell cycle arrest at G0/G1 phase as well as at G2-M phase in different cancer cells.

In HepG2 hepatoma cells, a remarkable arrest induction at G0/G1 phase was found after 24 h of astaxanthin treatment (25 and 42 μ M), showing a strong reduction of cells in S and G2-M phases [45]. These findings were evidenced in other cancer types and the mechanism involved in the cell cycle arrest was investigated. Generally, the cell cycle progression is regulated by a set of CDKs. They are serine/threonine kinases that control cell cycle progression and modulate their catalytic activities by interaction with cyclins and CDK inhibitors. In malignant cells, changes in expression of CDKs, overexpression of cyclins, and under expression of CDK inhibitors have been commonly reported. The altered CDK activity represents a selective advantage for cancer cell growing. Several studies reported that astaxanthin affects the activity of CDK, repressing tumor progression. Astaxanthin as carotenoid compounds has been reported to restrain tumor cell growth stimulating p21 and inhibiting cyclin D1 [67]. As cyclin D1 controls G1 to S-phase shift, its suppression may be associated to a delay in of tumor cell line proliferation. A CDK inhibitor as p21 is responsible of G1 arrest and, consequently, cell cycle arrest [56].

In human leukemia K562 and HL-60 cells, astaxanthin activates PPAR γ , resulting in p21 stimulation and cyclin D1 inhibition, leading to cell cycle arrest at G0/G1 phase [68,69]. PPAR γ , belonging to the nuclear receptor superfamily, acts as a transcription factor and exerts strong anti-proliferative effects in a great range of neoplastic cells [70]. Among ligands that could target PPAR γ receptors, rosiglitazone arises as an anticancer agent towards a large set of cancers including colon, breast, prostate, and lung [71]. For instance, Liu, et al. [72] reported that rosiglitazone (> 40 μ M) induced apoptosis in K562 cells, depending by time and dose. Interestingly, the same effects and in the same cells were observed under astaxanthin treatment, with a significant and dose-dependently induction of cellular apoptosis and PPAR γ protein expression in K562 cells. These effects might be limited by administrating antagonist molecules of PPAR γ receptor [56]. Moreover, the association of astaxanthin and PPAR γ was firstly described in 2005 in fibroblasts [73], when the astaxanthin-related expression of connexin 43, involved in early carcinogenesis stages, was inhibited by PPAR γ antagonists [74].

In human gastric adenocarcinoma, KATO-III and SNU-1 cells, the cell cycle arrest was shown at the G0/G1 phase reducing cell proliferation at the S phase (range 0–100 μ M) [54]. As ERK and Akt play central roles in cell growth and survival, their inhibition in KATO-III and SNU-1 cells by astaxanthin (100 μ M) is reflected in a decrease of protein synthesis and cell proliferation, due to cyclin D1/CDK4 complex inhibition. In both cell lines, astaxanthin increased, in a concentration-dependent manner, the protein level of p27, an inhibitor of CDK, by suppression of cyclin E/CDK2 complex. These processes might induce the cell cycle arrest at the G0/G1 phase. Expression levels of p-Akt, p-Rb

and cyclin D1 were not influenced by astaxanthin in KATO-III and SNU-1 cells [54]. No effects were reported on cell cycle of AGS and MKN-45 cancer cell lines when astaxanthin has been tested at different concentration (0–100 μ M) [54]. The inconsistent results among cell lines might derive from their diverse features, showing different molecular characteristics, including genetic and epigenetic changes (cell cycle regulator genes, growth factors etc.). The antiproliferative effect on oral squamous cell carcinoma models was confirmed by the reduction of the expression of cyclin D1 along with the increase in the nuclear p21, colony formation inhibition, in both SCC131 and SCC4 oral cancer cells (0–700 μ M) [57]. Astaxanthin induced G1 arrest in melanoma A2058 and A375 cells [42] and in A549 lung cancer cells measured by flow cytometric analysis [75].

Cell growth arrest might be observed after 24 h treatment with astaxanthin on colon cancer HCT116, HT-29, LS-174, WiDr, SW-480 cells [76]. In HCT116 treated with astaxanthin, the level of cyclin D1, p27 and p21 was similarly affected by the treatment with *H. pluvialis* extract [76,77]. Studies of the growth-inhibitory effects of astaxanthin (synthetic, S: meso: R = 1:2:1) and its stereoisomers (R- and S-astaxanthin) on HCT116 and HT-29 colon cancer cells were deepened. Astaxanthin stereoisomers, from *H. pluvialis* and *P. rhodozyma*, were tested without significant difference in the cell inhibition. From flow cytometry analysis, astaxanthin and its stereoisomers caused an increased number of HCT116 cells in G2/M after 48 h treatment to 1.4–2.4-times of the control cells. The cell cycle arrest is accompanied with a decrease of cells in G1/S phase and is associated to up-regulated expression of p21, and p27, as well as down-regulation of CDK4 and CDK6 [49].

In human breast cancer, astaxanthin isolated from shrimp, showed the cell cycle arrest at the G0/G1 phase associated to a down-regulation of cyclin D1 in MCF-7 cells [67], whereas Atalay, et al. [78] reported an accumulation of G2/M increase on the same cell line. However, in both reports the antiproliferative effects of astaxanthin is confirmed and differences in the astaxanthin origin, the vehicle used for the treatment, time and dose could be involved. Besides, effect of a combination of carotenoids (astaxanthin, lutein and β -carotene) on MCF-7 cell cycle progression has been studied suggesting that the co-administered carotenoids reduced cyclin D1 expression of G0/G1 phase more than individually. It is important to underline as no effects of astaxanthin on cell proliferation and cell cycle regulation have been reported in normal breast epithelial cells (MCF-10A) [67].

A reduction in Cyclin B1 and Cdc2 expression triggers G2/M cell cycle arrest as reported in ECA 109 esophageal squamous cell carcinoma [79] treated with astaxanthin. A significant increase of cells in the G2/M phase has been recorded as antiproliferative mechanism in other cancer cells. Unlike the HepG2, astaxanthin arrested hepatoma H22 cells in G2 phase [47], according to flow cytometry analysis. *In vitro* H22 cells showed G2 phase cells increased and G0/G1 and S phase cells decreased when low-dose and high-dose of astaxanthin were used, as also confirmed by *in vivo* studies [47]. The role of vehicle during

treatment is significant in modern drug/nutraceutical delivery, in fact, a synergism between the vehicle and active ingredients might ensure the therapeutic efficiency. Sowmya, et al. [80] investigated the role of the vehicles, THF, DMSO, FBS, for astaxanthin delivery on *in vitro* cervical adenocarcinoma HeLa cells. Results revealed that DMSO seems to be a better vehicle for astaxanthin keto-carotenoid delivery and the accumulation of G2/M phase is coupled with a cell reduction in S phase [80].

These findings showed that cell cycle arrest is a significant anti-cancer aspect induced by astaxanthin (Fig. 6). However, the anti-cancer mechanisms in cell cycle progression remain still unclear in some cancer cell lines.

5. Pro-apoptotic effects of astaxanthin

Programmed cell death is one of the possible fates of a cell. It is an essential mechanism during embryogenesis and cell differentiation, used to eliminate cells no longer needed by the body or damaged by factors such as oxidative stress, mitochondrial and/or DNA injury, growth factor deprivation, virus, radiation, chemotherapy. Apoptosis is characterized by a well-defined sequence of morphological and molecular changes, resulting in cell contraction, condensation and fragmentation itself, giving rise to the so-called *apoptotic bodies*, and finally death. Numerous intra- and/or extra-cellular molecules are involved: membrane receptors, which essentially bind factors or cytokines external to cells (e.g. TNF, Fas-L, TRAIL, *extrinsic via of apoptosis*), and/or proapoptotic (e.g. Bax, Bam, Bid) and antiapoptotic (e.g. IAP and Bcl2) proteins, most of which have mitochondrial origin (*intrinsic pathway of apoptosis*). To carry out the cell death program, the activity of proteolytic enzymes, called caspases, is essential [81,82]. A defective apoptotic process in any part of this complex mechanism is often involved in cancer, affecting the neoplastic transformation, the resistance to anticancer drugs and the possible metastatization [83–85]. Certainly, one of the therapeutic strategies to counteract the proliferation of tumor cells consists in aiming at the apoptotic mechanism. It is also shown that many molecules of natural origin have anti-tumor properties, aiming precisely at the intracellular mechanisms of programmed cell death, restoring, more or less consistently, a normal cellular homeostasis [86–88]. They have the undoubted advantage of high bioavailability and relatively less potent side effects. Many different phytochemical compounds of natural origin are studied and used in various types of cancer and among these many carotenoids [89–92], including astaxanthin. Its proapoptotic activity on cancer cells, has been highlighted in several studies, both *in vitro* and *in vivo* models. Specifically, an increase in astaxanthin-induced apoptosis was elucidated in colorectal carcinoma cell lines HCT-116, HT-29 [49,53,76], LS-174, WiDr, SW480 [76] and LS-180 [93], AGS gastric cells [53], breast cancer cells MCF7 [67,94], MDA-MB-231 [61,95] and MDA-MB-468 [60], human hepatocellular carcinoma cell lines LM3, SMM-7721 [46], HepG2 [96], rat HCC cells CBRH-7919 [43,97] and mouse HCC cells H22 [47], lung cancer cells A549 [75], melanoma cell lines A375 and A2058 [42], ovarian carcinoma cells SKOV3 [98], esophageal squamous cancer cells ESCC [79], rabbit squamous cancer cells VX2 [66], rat oral squamous carcinoma cells SCC131 and human oral squamous carcinoma cells SCC4 [57], chronic myelogenous leukemia (CML) cells K562 [56]. *In vivo* studies have also been carried out, which have helped to believe in an effective antitumor activity of astaxanthin through its proapoptotic effect, as reported successively.

Astaxanthin also showed its proapoptotic activity by down-regulating the expression of antiapoptotic Bcl-2, p-Bad, and survivin proteins and upregulating proapoptotic Bax/Bad and PARP proteins, together with the efflux of Smac/Diablo and cytochrome *c* into the cytosol. The inhibition of these pathways was then further confirmed in LM3 and SMMC-7721 human HCC cell lines [46], in which it was shown an activation of caspases 3 and 9, a downregulation of Bcl2 and an upregulation of Bax, markers of an apoptotic process, probably

caused by NF- κ B p65 and Wnt/ β -catenin down-regulation *via* negative activation of PI3 K/Akt and ERK signaling pathways. Interestingly, PI3 K/Akt signaling pathway and its downstream effectors have been shown to be affected by astaxanthin in various reports. For example, addition of astaxanthin prevents cell proliferation in SCC131 and SCC4 oral cancer cells by inhibiting PI3 K/Akt and NF- κ B pathways [57]. Moreover, astaxanthin exerted chemopreventive effects in the hamster buccal pouch (HBP) carcinogenesis model by inducing intrinsic apoptosis *via* the abrogation of the upstream kinase PI-3 K/Akt, and its downstream signaling pathway NF- κ B [99]. Activated Akt, migrating from the cytoplasm to the cell membrane, activates its downstream pathways, such as mTOR, p70S6K, MMP-2 and NF- κ B. mTOR and p70S6K enhance protein synthesis and cellular proliferation, while NF- κ B reduces apoptosis-related genes transcription and upregulates the IGF-IR and MMP2 expression, associated with invasion and metastasis. Finally, in recent studies, the central role of this pathway, as astaxanthin target, has still been highlighted. Su, et al. [98] showed that astaxanthin combined treatment with human serum albumin (HAS) was able to inhibit NF- κ B expression and its translocation to nucleus, promoting a cytotoxic effect in ovarian cancer cells SKOV3. Moreover, Kowshik, et al. [57] confirmed that the PI3 K/NF- κ B /STAT3 axis is a specific target of astaxanthin in promoting cellular apoptotic process. The involvement of STAT3 was also examined by Song, et al. [97], resulting that astaxanthin could induce the apoptosis of rat hepatocellular carcinoma CBRH-7919 cells by inhibiting the expression of STAT3 and its upstream activator JAK1, so dysregulating the expression of genes involved in apoptosis [100].

In LS-180 colorectal cancer cells, astaxanthin increased expression of Bax and caspase 3 and decreased that of Bcl2, triggering the apoptotic process [93]. It was showed that a nanoemulsion of astaxanthin and alpha-tocopherol (a type of vitamin E) was able to induce mitochondria-mediated apoptosis in HT-29 and AGS colorectal and gastric cancer cells [53] through ROS generation and damaged mitochondrial membrane potential, opening up encouraging outlook for the use of new formulations well balanced between high efficacy, high absorption rate and low toxicity. A different extraction procedure was used to obtain astaxanthin from shrimp by-products and use it to induce apoptosis in HepG2 hepatocellular carcinoma cells through the upregulation of p53 expression, the activation of caspase 3 and a down-regulation of c-Jun and c-Fos [96]. Some carotenoids were tested, also in combination with the anthracycline drug doxorubicin, against MDA-MB-468 [60], MCF7 and MDA-MB-231 breast cancer cells [95], to induce oxidative stress-mediated apoptosis, showing that lutein and beta-carotene were more potent cell growth inhibitors and more effective in triggering the apoptotic process compared to fucoxanthin and astaxanthin. However, another study revealed a synergistic efficacy of astaxanthin plus β -carotene and lutein carotenoids on molecular events in MCF-7 cells, inducing enhanced cytotoxicity and oxidative stress, higher cellular uptake/accumulation of astaxanthin along with β -carotene and lutein to synergistically trigger apoptosis [67]. In HCT116 and HT-29 colon cancer cells, different stereoisomers of astaxanthin stimulated apoptosis, as suggested by the analysis of increased levels of caspase 3, PARP, EGFR, p-p38, p-JNK and pERK1/2 [49]. A formulation as gold nanoparticles of astaxanthin [61] showed cytotoxic effect against MDA-MB-231 breast cancer cell, inducing apoptosis and revealing itself also useful as contrast agent in photoacoustic imaging.

6. Astaxanthin attenuates cancer diffusion

Cancer diffusion includes a complex cascade of events as angiogenesis, migration and invasion of cancer cells to new tissues and promising anti-cancer molecules might be able to modulate and attenuate these multiple hallmarks.

In the last years, the cell migration targeting has become a therapeutic approach worth to be investigated for cancer treatment [101]. *In vitro* wound healing and invasion assays were performed to evaluate

the anticancer effect of astaxanthin. MCF7, MDA-MB-231 breast cancer cell lines were associated to a direct diminution in distance migrated when treated with astaxanthin (10–50 μM), than normal breast epithelial cells [64]. It has been found that 42 μM astaxanthin is associated with a reduction in cell migration to 50.2 % in A375 and 78.1 % in A2058 through the suppression of MMPs 1,2,9 expression [42].

MMPs, zinc-binding endopeptidases, play a crucial role in cancer and they can promote the cancer metastasis by breaking the ECM [102].

The effect of astaxanthin in tumor angiogenesis has been poorly investigated. Angiogenesis is the physiological mechanism for the formation of new blood vessels from preexisting ones during normal processes as embryogenesis, menstruation, and wound healing. In case of tumor cells, the release of angiogenic stimulators lead to an uncontrolled vessel growth. Several reports suggested that astaxanthin might mediate their inhibition. Indeed, angiogenesis was limited in oral squamous cell carcinoma after astaxanthin treatment [57,103]. Astaxanthin significantly decreases the expression of MMP-2, MMP-9, VEGF, VEGFR2 and HIF-1 α nuclear translocation in 131 and SCC4 oral squamous carcinoma cells by JAK-2/STAT3 signaling pathway [57]. Normally, JAK/STAT pathway is involved in the growth and development processes, while in case of cancer, the dysregulation is associated to over proliferation and metastasis [104–106]. Recently, it has been reported that astaxanthin supplementation might affect this pathway and downstream events, in the HBP carcinogenesis model [103]. Accordingly, astaxanthin downregulates the expression of STAT-3 target genes involved in invasion (MMP-2 and MMP-9) in this model.

In rat ascites hepatoma cells (AH109A), 5 μM of astaxanthin inhibited the invasion, allowing to counteract the metastatic process [107].

Astaxanthin induced the EMT in CT26 and HCT116 colon cancer cells by modulation of the expression level of specific markers; in particular astaxanthin increased the expression of epithelial marker E-cadherin and reduced the mesenchymal and invadopodia markers vimentin and cortactin, respectively [108]. One fancy hypothesis about carcinogenesis confers a role to miRNAs, small noncoding RNAs responsible of post-transcriptional control of gene expression. Interestingly, astaxanthin increased miR-29a-3p and miR-200a expression in colorectal cancer cells inhibiting the ability of these cells to colonize surrounding tissues [108]. mRNA and the protein expression of MMP2 were suppressed in HCT116 and CT26 cells. These results indicated that increased miR-29a-3p by astaxanthin suppressed MMP2 expression by direct targeting 3'UTR of mRNA and affected the metastatic processes of colon cancer cells. In addition, the increased expression of miR-200a in CT26 cells, and the related ZEB1 restore, reduced invasion and migration extension. ZEB1 stimulates EMT and promotes metastasis by E-cadherin repression. In colon cancer cells, astaxanthin administration reduced ZEB1 mRNA and MYC oncogenic transcription factor protein and mRNA expression [108].

Since it has been demonstrated that viral infections may be one of the major causes of tumor occurrence, researchers have been focused on new natural approaches in counteracting virus-associated cancer [109]. It is well known that HPV is one of the main etiologic factors for cervical and testicular neoplasia, while HSV have implications on nasopharyngeal cancer. In this context, natural carotenoids with antiviral properties might represent a useful tool in virus-associated tumor treatment. Donà, et al. [110] revealed that astaxanthin is able to preserve sperm by reducing the amount of HPV16-L1 bound onto membranes, showing its antiviral potential. Given the promising results of carotenoids [111] in decreasing the persistence of HPV and the important antiviral activities of astaxanthin-rich *H. pluvialis* extract [112], further investigations should be addressed to evaluate the correlation between astaxanthin and decrement of testicular and cervix cancer development risk.

7. Astaxanthin chemosensitizing effects

Chemosensitization is a strategy to potentiate the cytotoxic effect of anticancer drugs. Astaxanthin as other natural compounds showed an important synergistic effect with antineoplastic drug in addition to their cancer chemoprevention activity. For example, co-treatment of breast cancer cells (MDA-MB, MCF-7) with astaxanthin and a minimal cytotoxic concentration of doxorubicin significantly reduced cell viability compared to control cells [95]. Particularly, astaxanthin allowed to reduce doxorubicin dose (0.12 μM in MCF-7 and 0.28 μM in MDA-MB-231 cells) and IC₅₀ value of co-treatment (6.6 \pm 0.1 μM in MCF-7 cells and 9.5 \pm 0.2 μM in MDA-MB-231 cells) was significantly lower than astaxanthin alone (23.5 \pm 0.1 μM in MCF-7 cells and 26.6 \pm 0.2 μM in MDA-MB-231 cells) [95]. AlQahtani, et al. [113] reported as astaxanthin treatment increased percentage of cells in G2/M, the best phase for the activity of doxorubicin due to maximum expression of its target enzyme topoisomerase II. Moreover, the carotenoid increased doxorubicin cellular uptake in tumor cells by inhibition of P-glycoprotein pump, important for absorption, distribution and elimination of the drug [113]. Antioxidant and anti-inflammatory properties of astaxanthin are also important to reduce doxorubicin neurotoxicity due to increase of ROS production and proinflammatory cytokines expression. Astaxanthin showed neuroprotective effect in male albino rats treated with combination of astaxanthin (25 mg/kg/week) and doxorubicin (2 mg/kg/week). Particularly, it restored hippocampal function and cholinesterase activity and reduced ROS and proinflammatory cytokines production [114].

The synergistic effect of astaxanthin has also been demonstrated with other anticancer drugs. Astaxanthin enhanced cytotoxic effect of erlotinib in human lung carcinoma cells A549 and H1975 by reducing expression of XPC, an important protein in recognizing DNA damage via activation of p38 MAPK [115]. Different combination of astaxanthin and carbendazim, fungicide and also antitumor agent against a variety of mammalian carcinoma, decreased cell viability of MCF-7 human breast carcinoma cells increased after 24 h of treatment. Particularly, different doses of astaxanthin (8, 17, 25 μM) combined with carbendazim (15 μM) led to a higher anti-proliferative effect than carbendazim alone. Moreover, co-treatment increased G2/M phase arrest compared to the control but not to carbendazim alone [78]. Astaxanthin thanks to its antioxidant activity can protect normal tissue from damage of oxidative stress of different antineoplastic drugs. Several studies [116–118] demonstrated as treatment with astaxanthin reduced cisplatin nephrotoxicity and gonadal toxicity in male cancer patients treated with cyclophosphamide. Further, astaxanthin treatment reduced ROS generation in C57BL/6 mice exposed to 4 Gy radiation and improved hematopoietic injury [119].

Astaxanthin resensitized gemcitabine-resistant human pancreatic cancer cells to gemcitabine. Co-treatment with the carotenoid increased cell chemosensitizer by acting on key target of gemcitabine resistance like human equilibrative nucleoside transporter (hENT1) and ribonucleotide reductase subunit M1(RRM1) and M2 (RRM2) and inhibited the gemcitabine-induced epithelial-mesenchymal transition phenotype by promoting expression of epithelial cell markers instead of mesenchymal ones [120]. Indeed, these studies evidence as astaxanthin enhanced tumor cells sensitization to chemotherapy and limit its adverse reactions.

8. Astaxanthin in pre-clinical studies

From the first assumptions about the correlation between blood carotenoids levels and cancer prevention in the 1980s [121], many evidence have been collected to confirmation of the *in vivo* anticancer properties of astaxanthin (Table 1). The inflammatory nature of the major chronic pathologies, as well as cardiovascular and neurodegenerative diseases, diabetes and cancer makes the xanthophyll astaxanthin, with its antioxidative properties, one of the most promising

therapeutic agents in this context. Hyperglycemia, indeed, is associated with an increment of oxidative stress, with consequent tissue and cellular damages. Astaxanthin revealed *in vivo* protective properties in preserving pancreatic cell functions on diabetic mice model (db/db) [122]. Moreover, it restored diabetes-dependent liver complications in rats [123], as well as reduced renal impairment in db/db mice [124,125] and protected neurons from hyperglycemic damage, preventing depression in mice [126]. By counteracting oxidative stress, astaxanthin has a pivotal role in the context of neurodegenerative disorders and aging. Astaxanthin treatment with a dose of 10 mg/kg/day improved learning and memory impairments in mice models of cerebral ischemia-reperfusion [127] and modulated inflammatory mediators release [21]. Astaxanthin treatment (25 mg/kg) might restore brain histopathological architecture and protect against doxorubicin-induced memory impairment [114]. In the same way, the antioxidant properties of astaxanthin are responsible for cardioprotective effects on rats [128], antiatherosclerotic properties on hyperlipidemic rabbits [129], prevention of bone loss in ovariectomized mice [130], as already shown in previous reported *in vitro* experiments. Moreover, oral administration of astaxanthin (100 mg/kg) reduced skin severity symptoms in murine model of atopic dermatitis [131]. To confirm the *in vitro* implications of astaxanthin in eye care, *in vivo* treatment with oral astaxanthin (100 mg/kg) also attenuated retinal damage [132]. As a microenvironment of chronic inflammation is strictly linked to the promotion of carcinogenesis, the chemopreventive role of astaxanthin might be related to the reduction of inflammation. As reported by Chiu et al. (2016), indeed, the reduction of hepatoma development in rats is associated with the reduction of hepatotoxicity [133]. Shao et al. (2016) confirmed their own *in vitro* results also on mice, showing an inverse correlation between astaxanthin dose and tumor weight [47]. In light of the results of Ohno et al. (2016), astaxanthin might help in liver chemoprevention in obese people. In fact, they found that dietary supplementation with 340 μ M for twenty weeks markedly attenuated hepatocarcinogenesis in db/db mice [134]. Astaxanthin treatment might allow stopping cancer cells at an early stage of invasiveness. Therefore, 14-days oral treatment with 1 mg/kg/day of astaxanthin in DBA mice is associated to a strong inhibition of metastases, while the administration of 100 mg/kg/day for 4-day exerted a modulatory effect on natural killer cells that have a crucial role in antiproliferative activity and the inhibition of cancer metastatic process, improving antitumor immune responses [135]. As shown in *in vitro* data on LX-2 cells, the role of PPARs in liver carcinogenesis is pivotal. *In vivo* astaxanthin treatment might modulate the expression of PPAR- α and PPAR- γ in liver and gastric tissue, respectively, with an improvement of oxidative status and profound implications in chemoprevention and carcinogenesis [136]. As already reported by *in vitro* studies, astaxanthin *in vivo* contrasted DNA damages induced by aflatoxin B1 [31], as well as suppressed tongue neoplasm development, when supplementation with 170 μ M was included in rats diet [137]. With the involvement of the same pathways observed *in vitro*, the *in vivo* chemoprevention of astaxanthin has been demonstrated in the hamster buccal pouch model of oral carcinogenesis by dietary supplementation of 15 mg/kg of astaxanthin which resulted in a significant delay in tumor development [57,138]. The *in vitro* chemotherapeutic effects of astaxanthin against skin cancer were also confirmed in skin carcinogenesis mice model with a decreased number of papillomas after topical application of 850 μ M of astaxanthin [139]. Once again, the modulation of tyrosinase activity might explain the reduction of skin tumor growth exerted by astaxanthin (200 μ g/kg) on rats [140]. The first proofs of chemopreventive efficacy of xanthophylls on urinary bladder carcinogenesis date from the 1990s, when Tanaka, et al. [141] observed that 20-weeks administration of 85 μ M in male ICR mice markedly reduced the incidence of preneoplastic lesions. The ability of astaxanthin to exert anti-tumorigenic activity was also observed in nude PC-3 prostate cancer mice. In this model astaxanthin significantly inhibited tumor growth by increasing the levels of miR-375 and miR-487b in tumor tissues at high (100 mg/kg) doses with a

decrease in cancer growth by up to 41 % by apoptosis induction [142].

High doses of astaxanthin revealed efficient also in mammary cancer suppression. Supplementation with 0.4 %, but not with 0.04 %, halved the incidence of palpable mammary carcinoma and reduced tumor weight [143]. In breast cancer bearing-mice, a single intratumoral injection of an aqueous solution containing 25 mg of astaxanthin reduced microelements concentration in tumor tissues, contributing to provide anticancer effects [144]. Nakao, et al. [145] demonstrated that astaxanthin pretreatment in mice prevented mammary tumor development, but it was unable to stop cancer progression in late stages [145]. Astaxanthin treatment with 340 μ M for 8 weeks reduced oxidative stress associated to precancerous lesions in colon cancer-bearing db/db mice [146]. Doses of 15 mg/kg are associated with a decrement of inflammatory mediators, involved in cancer progression, revealing astaxanthin as a helpful chemotherapeutic agent in rat colon carcinogenesis model [147]. Most recently, it was demonstrated that astaxanthin in rats may reduce NF κ B levels, preventing progression of NMBA-induced esophageal cancer in early stage of the onset [148]. The results of *in vivo* investigations on the safety of synthetic astaxanthin were in agreement with those obtained *in vitro*, showing no genotoxicity. The increment of benign liver adenoma in female rats after astaxanthin supplementation seemed to be due to a rodent specific sensitivity, not shared with humans [149]. By contrast, it exerted a protective role against germ cell toxicity under cisplatin therapy. Indeed, 5-weeks of oral treatment with 25 mg/kg of astaxanthin is associated with sperm DNA repairing in male mice [116]. Astaxanthin reconfirmed, also *in vivo*, to act as a valuable aid to conventional chemotherapy in case of drug-resistance, thanks to the photothermal effects. As shown in VX2 tumor rabbit models, the intratumoral dose of 300 μ g/mL of astaxanthin followed by laser irradiation is associated with complete and selective tumor eradication [66].

Astaxanthin is often called the “king of carotenoids”, thanks to the well-recognized benefits on human health, but despite its “regal” title, even that *in vitro* and preclinical anti-cancer activity has been widely investigated. To the best of our knowledge, there are no clinical trials or well-defined clinical studies assessing the efficacy of astaxanthin administration reported in patients suffering of any kind of cancer [150].

9. Conclusion and future perspectives

This review evidences the recent advances in the detailed description of anti-cancer effects of astaxanthin. The *in vitro* studies reported the diverse targets through which astaxanthin reduces the tumorigenesis, mainly through the cell proliferation inhibition, cell cycle arrest and apoptosis induction. *In vivo* preclinical studies, were able to assess the efficacy of astaxanthin treatment in model organisms. Furthermore, astaxanthin enhanced tumor cells sensitization to chemotherapy and limit its adverse reactions. In this context, astaxanthin could be considered as a potential candidate to apply in combining therapy in combating cancer in resistance developing patients to conventional therapy.

Further *in vivo* experiments are required to deeply investigate the anti-cancer effects of astaxanthin.

Given the lack of clinical evidence, randomized, double-blind, placebo-controlled trials in cancer patients are necessary in order to evaluate the anticancer astaxanthin effectiveness in a more reproducible and safe way. Finally, from scientific results, astaxanthin arises as a valuable alternative in anticancer therapy.

Acknowledgements

This work was supported by the Regione Basilicata; the Fondazione Enrico Mattei DGR n. 1490 del 4/12/2014, vs rep. n. 163 n8; and the Regional Project ALIMINTEGRA, GO NUTRIBAS financed on 16.1 PSR Basilicata founding ex D.G.R. n° 312/17 CUP: C31G18000210002.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.phrs.2020.104689>.

References

- [1] R. Hu, S. CL-L, R. Yu, K. A-NT, Regulation of NF-E2-related factor 2 signaling for cancer chemoprevention: antioxidant coupled with antiinflammatory, *Antioxid. Redox Signal.* 13 (11) (2010) 1679–1698.
- [2] F. Bray, J. Ferlay, I. Soerjomataram, R.L. Siegel, L.A. Torre, A. Jemal, Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries, *CA Cancer J. Clin.* 68 (6) (2018) 394–424.
- [3] A.L. Demain, P. Vaishnav, Natural products for cancer chemotherapy, *Microb. Biotechnol.* 4 (6) (2011) 687–699.
- [4] M.F. Armentano, F. Bisaccia, R. Miglionico, D. Russo, N. Nolfi, M. Carosino, P.B. Andrade, P. Valentao, M.S. Diop, L. Milella, Antioxidant and proapoptotic activities of *Sclerocarya birrea* [A. Rich.] Hochst. methanolic root extract on the hepatocellular carcinoma cell line HepG2, *Biomed Res. Int.* 2015 (2015) 561589.
- [5] D. Lamorte, I. Faraone, I. Laurenzana, L. Milella, S. Trino, L. De Luca, L. Del Vecchio, M.F. Armentano, C. Sinigaglia, L. Chiumbento, D. Russo, F. Bisaccia, P. Musto, A. Caivano, Future in the past: azorella glabra wedd. As a source of new natural compounds with antiproliferative and cytotoxic activity on multiple myeloma cells, *Int. J. Mol. Sci.* 19 (11) (2018).
- [6] E. Patridge, P. Gareiss, M.S. Kinch, D. Hoyer, An analysis of FDA-approved drugs: natural products and their derivatives, *Drug Discov. Today* 21 (2) (2016) 204–207.
- [7] S. Fakhri, F. Abbaszadeh, L. Dargahi, J.M. Astaxanthin, A mechanistic review on its biological activities and health benefits, *Pharmacol. Res.* 136 (2018) 1–20.
- [8] K.P. Astaxanthin, Cell membrane nutrient with diverse clinical benefits and anti-aging potential, *Altern. Med. Rev.* 16 (4) (2011).
- [9] R.R. Ambati, S.-M. Phang, S. Ravi, Aswathanarayana RG. Astaxanthin: sources, extraction, stability, biological activities and its commercial applications—a review, *Mar. Drugs* 12 (1) (2014) 128–152.
- [10] L.M.A.J. Seabra, L.F.C. Pedrosa, Astaxanthin structural and functional aspects, *Rev Nutr, Campinas* 23 (6) (2010) 1041–1050.
- [11] C. Xueding, C. Yanfan, X. Xiaona, Y. Dan, D. Cheng, C. Mayun, Astaxanthin prevents against lipopolysaccharide-induced acute lung injury and sepsis via inhibiting activation of MAPK/NF- κ B, *Am. J. Transl. Res.* 11 (3) (2019) 1884–1894.
- [12] P. Zhou, M. Li, B. Shen, Z. Yao, Q. Bian, L. Ye, H. Yu, Directed coevolution of beta-carotene ketolase and Hydroxylase and its application in temperature-regulated biosynthesis of astaxanthin, *J. Agric. Food Chem.* 67 (4) (2019) 1072–1080.
- [13] F.J. Pashkow, D.G. Watumull, C.L. Campbell, Astaxanthin: a novel potential treatment for oxidative stress and inflammation in cardiovascular disease, *Am J Card.* 101 (10) (2008) S58–S68.
- [14] I. Higuera-Ciapara, L. Felix-Valenzuela, F.M. Goycoolea, Astaxanthin: a review of its chemistry and applications, *Crit. Rev. Food Sci. Nutr.* 46 (2) (2006) 185–196.
- [15] R.K. Saini, Y.S. Keum, Microbial platforms to produce commercially vital carotenoids at industrial scale: an updated review of critical issues, *J. Ind. Microbiol. Biotechnol.* 46 (5) (2019) 657–674.
- [16] M. Shah, R. Mahfuzur, Y. Liang, J.J. Cheng, M. Daroch, Astaxanthin-producing green microalga *Haematococcus pluvialis*: from single cell to high value commercial products, *Front. Plant Sci.* 7 (2016) 531.
- [17] L.S. Steelman, W.H. Chappell, S.L. Abrams, C.R. Kempf, J. Long, P. Laidler, S. Mijatovic, D. Maksimovic-Ivanic, F. Stivala, M.C. Mazzarino, M. Donia, P. Fagone, G. Malaponte, F. Nicoletti, M. Libra, M. Milella, A. Tafuri, A. Bonati, J. Basecke, L. Cocco, C. Evangelisti, A.M. Martelli, G. Montalto, M. Cervello, J.A. McCubrey, Roles of the Raf/MEK/ERK and PI3K/PTEN/Akt/mTOR pathways in controlling growth and sensitivity to therapy-implications for cancer and aging, *Aging-Us.* 3 (3) (2011) 192–222.
- [18] R.G. De Oliveira Júnior, A.F.C. Adrielly, J.R.G. da Silva Almeida, R. Grougnet, V. Thiéry, L. Picot, Sensitization of tumor cells to chemotherapy by natural products: a systematic review of preclinical data and molecular mechanisms, *Fitoterapia* 129 (2018) 383–400.
- [19] A.M. Salvia, F. Cuvillo, S. Coluzzi, R. Nuccorini, I. Attolico, S.P. Pascale, F. Bisaccia, M. Pizzuti, A. Ostuni, Expression of some ATP-binding cassette transporters in acute myeloid leukemia, *Hematol. Rep.* 9 (4) (2017) 137–141.
- [20] H.D. Choi, Y.K. Youn, W.G. Shin, Positive effects of astaxanthin on lipid profiles and oxidative stress in overweight subjects, *Plant Foods Hum. Nutr.* 66 (4) (2011) 363–369.
- [21] B. Grimmig, S.-H. Kim, K. Nash, P.C. Bickford, R.D. Shytle, Neuroprotective mechanisms of astaxanthin: a potential therapeutic role in preserving cognitive function in age and neurodegeneration, *Geroscience* 39 (1) (2017) 19–32.
- [22] M. Zhang, Z. Cui, H. Cui, Y. Wang, C. Zhong, Astaxanthin protects astrocytes against trauma-induced apoptosis through inhibition of NKCC1 expression via the NF- κ B signaling pathway, *BMC Neurosci.* 18 (1) (2017) 42–42.
- [23] S.H. Kim, J.W. Lim, H. Kim, Astaxanthin inhibits mitochondrial dysfunction and Interleukin-8 expression in *Helicobacter pylori*-infected gastric epithelial cells, *Nutrients* 10 (9) (2018).
- [24] S. Li, T. Takahara, M. Fujino, Y. Fukuhara, T. Sugiyama, X.-K. Li, S. Takahara, Astaxanthin prevents ischemia-reperfusion injury of the steatotic liver in mice, *PLoS One* 12 (11) (2017) e0187810.
- [25] Y. Lee, S. Hu, Y.-K. Park, J.-Y. Lee, Health benefits of carotenoids: a role of carotenoids in the prevention of non-alcoholic fatty liver disease, *Prev. Nutr. Food Sci.* 24 (2) (2019) 103.
- [26] X. Xie, Q. Chen, J. Tao, Astaxanthin promotes Nrf2/ARE signaling to inhibit HG-Induced renal fibrosis in GMCs, *Mar. Drugs* 16 (4) (2018).
- [27] D. Gao, H. Wang, Y. Xu, D. Zheng, Q. Zhang, W. Li, Protective effect of astaxanthin against contrast-induced acute kidney injury via SIRT1-p53 pathway in rats, *Int. Urol. Nephrol.* 51 (2) (2019) 351–358.
- [28] N. Liu, J. Chen, D. Gao, W. Li, D. Zheng, Astaxanthin attenuates contrast agent-induced acute kidney injury *in vitro* and *in vivo* via the regulation of SIRT1/FOXO3a expression, *Int. Urol. Nephrol.* 50 (6) (2018) 1171–1180.
- [29] K. Hara, C. Hamada, K. Wakabayashi, R. Kanda, K. Kaneko, S. Horikoshi, Y. Tomino, Y. Suzuki, Scavenging of reactive oxygen species by astaxanthin inhibits epithelial-mesenchymal transition in high glucose-stimulated mesothelial cells, *PLoS One* 12 (9) (2017) e0184332-e0184332.
- [30] J. Li, J. Sun, B. Li, Z. Liu, Astaxanthin protects ARPE-19 cells against oxidative stress injury induced by hydrogen peroxide, *Biotechnol. Equip.* 32 (5) (2018) 1277–1284.
- [31] S. Gradelet, P. Astorg, A.-M. Le Bon, R. Bergès, M. Suschetet, Modulation of aflatoxin B1 carcinogenicity, genotoxicity and metabolism in rat liver by dietary carotenoids: evidence for a protective effect of CYP1A inducers, *Cancer Lett.* 114 (1–2) (1997) 221–223.
- [32] H. Mori, T. Tanaka, S. Sugie, N. Yoshimi, T. Kawamori, Y. Hirose, M. Ohnishi, Chemoprevention by naturally occurring and synthetic agents in oral, liver, and large bowel carcinogenesis, *J. Cell. Biochem.* 67 (S27) (1997) 35–41.
- [33] A. Mortensen, L.H. Skibsted, J. Sampson, C. Rice-Evans, S.A. Everett, Comparative mechanisms and rates of free radical scavenging by carotenoid antioxidants, *FEBS Lett.* 418 (1–2) (1997) 91–97.
- [34] A. Murakami, M. Nakashima, T. Koshiba, T. Maoka, H. Nishino, M. Yano, T. Sumida, O.K. Kim, K. Koshimizu, H. Ohigashi, Modifying effects of carotenoids on superoxide and nitric oxide generation from stimulated leukocytes, *Cancer Lett.* 149 (1–2) (2000) 115–123.
- [35] I. O'Connor, N. O'Brien, Modulation of UVA light-induced oxidative stress by β -carotene, lutein and astaxanthin in cultured fibroblasts, *J. Derm. Sci.* 16 (3) (1998) 226–230.
- [36] M. Santocono, M. Zurria, M. Berrettini, D. Fedeli, G. Falcioni, Influence of astaxanthin, zeaxanthin and lutein on DNA damage and repair in UVA-irradiated cells, *J. Photochem. Photobiol. B, Biol.* 85 (3) (2006) 205–215.
- [37] K.N. Singh, S. Patil, H. Barkate, Protective effects of astaxanthin on skin: recent scientific evidence, possible mechanisms, and potential indications, *J. Cosmet. Dermatol.* (2019).
- [38] S. Chintong, W. Phatvej, U. Rerk-Am, Y. Waiyapri, W. Klaypradit, *In vitro* antioxidant, Antityrosinase, and cytotoxic activities of astaxanthin from shrimp waste, *Antioxidants* 8 (5) (2019) 128.
- [39] T. Niwano, S. Terazawa, H. Nakajima, G. Imokawa, The stem cell factor-stimulated melanogenesis in human melanocytes can be abrogated by interrupting the phosphorylation of MSK1: evidence for involvement of the p38/MSK1/CREB/MITF axis, *Arch. Dermatol. Res.* 310 (3) (2018) 187–196.
- [40] D. Ritto, S. Tanasawet, S. Singkhorn, W. Klaypradit, P. Hutamekalin, V. Tipmanee, W. Sukketsiri, Astaxanthin induces migration in human skin keratinocytes via Rac1 activation and RhoA inhibition, *Nutr. Res. Pract.* 11 (4) (2017) 275–280.
- [41] Y. Yang, I. Yang, M. Cao, S. Z-y, R. Wu, Y. Guo, M. Fang, A.-N. Kong, Fucoxanthin elicits epigenetic modifications, Nrf2 activation and blocking transformation in mouse skin JB6 P+ cells, *AAPS J.* 20 (2) (2018) 32.
- [42] Y.-T. Chen, C.-J. Kao, H.-Y. Huang, S.-Y. Huang, C.-Y. Chen, Y.-S. Lin, Z.-H. Wen, Wang H-MD, Astaxanthin reduces MMP expressions, suppresses cancer cell migrations, and triggers apoptotic caspases of *in vitro* and *in vivo* models in melanoma, *J. Funct. Food.* 31 (2017) 20–31.
- [43] S. X-d, Z. J-j, W. M-r, L. W-b, G. X-b, C. Lv, Astaxanthin induces mitochondria-mediated apoptosis in rat hepatocellular carcinoma CBRH-7919 cells, *Biol. Pharm. Bull.* 34 (6) (2011) 839–844.
- [44] J.C. Ko, J.C. Chen, T.J. Wang, H.Y. Zheng, W.C. Chen, P.Y. Chang, Y.W. Lin, Astaxanthin down-regulates Rad51 expression via inactivation of AKT kinase to enhance mitomycin C-induced cytotoxicity in human non-small cell lung cancer cells, *Biochem. Pharmacol.* 105 (2016) 91–100.
- [45] S. Nagaraj, M.G. Rajaram, P. Arulmurugan, A. Baskaraboopathy, K. Karuppasamy, K.R. Jayappriyan, R. Sundararaj, R. Rengasamy, Antiproliferative potential of astaxanthin-rich alga *Haematococcus pluvialis* Flowot on human hepatic cancer (HepG2) cell line, *Biomed Prev Nutr.* 2 (3) (2012) 149–153.
- [46] J. Li, W. Dai, Y. Xia, K. Chen, S. Li, T. Liu, R. Zhang, J. Wang, W. Lu, Y. Zhou, Astaxanthin inhibits proliferation and induces apoptosis of human hepatocellular carcinoma cells via Inhibition of NF- κ B P65 and Wnt/ β -catenin *in vitro*, *Mar. Drugs* 13 (10) (2015) 6064–6081.
- [47] Y. Shao, Y. Ni, J. Yang, X. Lin, J. Li, L. Zhang, Astaxanthin inhibits proliferation and induces apoptosis and cell cycle arrest of mice H22 hepatoma cells, *Med. Sci. Monit.* 22 (2016) 2152–2160.
- [48] V. Suganya, V. Anuradha, *In silico* molecular docking of astaxanthin and sorafenib with different apoptotic proteins involved in hepatocellular carcinoma, *Biocatal. Agric. Biotechnol.* 19 (2019) 101076.
- [49] X. Liu, M. Song, Z. Gao, X. Cai, W. Dixon, X. Chen, Y. Cao, H. Xiao, Stereoisomers of astaxanthin inhibit human colon cancer cell growth by inducing G2/M cell cycle arrest and apoptosis, *J. Agr Food Chem.* 64 (41) (2016) 7750–7759.
- [50] F.K. El-Baz, R.A. Hussein, K. Mahmoud, S.M. Abdo, Cytotoxic activity of carotenoid rich fractions from *Haematococcus pluvialis* and *Dunaliella salina* microalgae and the identification of the phytoconstituents using LC-DAD/ESI-MS, *Phytother. Res.* 32 (2) (2018) 298–304.
- [51] M. Ogasawara, T. Matsunaga, H. Suzuki, Differential effects of antioxidants on the *in vitro* invasion, growth and lung metastasis of murine Colon Cancer cells, *Biol. Pharm. Bull.* 30 (1) (2007) 200–204.

- [52] K. Wayakanon, K. Rueangyotchanthana, P. Wayakanon, C. Suwannachart, The inhibition of Caco-2 proliferation by astaxanthin from *Xanthophyllomyces dendrorhous*, *J Med Micro*. 67 (4) (2018) 507–513.
- [53] K. Shanmugapriya, H. Kim, H.W. Kang, *In vitro* antitumor potential of astaxanthin nanoemulsion against cancer cells via mitochondrial mediated apoptosis, *Int. J. Pharm.* 560 (2019) 334–346.
- [54] J.H. Kim, J.-J. Park, B.J. Lee, M.K. Joo, H.J. Chun, S.W. Lee, Y.-T. Bak, Astaxanthin inhibits proliferation of human gastric Cancer cell lines by interrupting cell cycle progression, *Gut Liver* 10 (3) (2016) 369–374.
- [55] L. S-a, J. Lee, W. Jung, E. Lim, M. Joo, B. Lee, J.-J. Park, J. Kim, Y.-T. Bak, S.W. Jung, S. Lee, Anticancer effects of astaxanthin and α -tocopherol in esophageal Cancer cell lines, *Korean J Helicobacter Up Gastrointest Res.* 11 (2011) 170.
- [56] X. Zhang, W.E. Zhao, L. Hu, L. Zhao, J. Huang, Carotenoids inhibit proliferation and regulate expression of peroxisome proliferators-activated receptor gamma (PPARgamma) in K562 cancer cells, *Arch. Biochem. Biophys.* 512 (1) (2011) 96–106.
- [57] J. Kowshik, R. Nivetha, S. Ranjani, P. Venkatesan, S. Selvamuthukumar, V. Veeravarmal, S. Nagini, Astaxanthin inhibits hallmarks of cancer by targeting the PI3K/NF-kappaBeta/STAT3 signalling axis in oral squamous cell carcinoma models, *IUBMB Life* 71 (10) (2019) 1595–1610.
- [58] Y. Yang, F. Fuentes, L. Shu, C. Wang, D. Pung, W. Li, C. Zhang, Y. Guo, A.-N. Kong, Epigenetic CpG methylation of the promoter and reactivation of the expression of GSTP1 by astaxanthin in human prostate LNCaP cells, *AAPS J.* 19 (2) (2017) 421–430.
- [59] M.L. Anderson, A preliminary investigation of the enzymatic inhibition of 5 α -Reductase and growth of prostatic carcinoma cell line LNCap-FGC by natural astaxanthin and saw palmetto lipid extract in vitro, *J. Herb. Pharmacother.* 5 (1) (2005) 17–26.
- [60] X. Gong, J. Smith, H. Swanson, L. Rubin, Carotenoid lutein selectively inhibits breast cancer cell growth and potentiates the effect of chemotherapeutic agents through ROS-mediated mechanisms, *Molecules* 23 (4) (2018) 905.
- [61] S. Bharathiraja, P. Manivasagan, N. Quang Bui, Y.-O. Oh, I.G. Lim, S. Park, J. Oh, Cytotoxic induction and photoacoustic imaging of breast Cancer cells using astaxanthin-reduced gold nanoparticles, *Nanomaterials Basel (Basel)* 6 (4) (2016) 78.
- [62] N. Jafari, S. Nazari, S.T. Enferadi, Parthenolide reduces metastasis by inhibition of vimentin expression and induces apoptosis by suppression elongation factor α - 1 expression, *Phytomedicine*. 41 (2018) 67–73.
- [63] P. Régnier, J. Bastias, V. Rodriguez-Ruiz, N. Caballero-Casero, C. Caballo, D. Sicilia, A. Fuentes, M. Maire, M. Crepin, D. Letourneur, Astaxanthin from *Haematococcus pluvialis* prevents oxidative stress on human endothelial cells without toxicity, *Mar. Drugs* 13 (5) (2015) 2857–2874.
- [64] B. McCall, C. McPartland, R. Moore, A. Frank-Kamenetskii, B. Booth, Effects of astaxanthin on the proliferation and migration of breast cancer cells in vitro, *Antioxidants*. 7 (10) (2018) 135.
- [65] M. Pilinska, D. Eurinnyi, S. Rushkovsky, I. Dybska, Genoprotective properties of astaxanthin revealed by ionizing radiation exposure in vitro on human peripheral blood lymphocytes, *Probl Radiat Med Radiobiol.* 21 (2016) 141–148.
- [66] V.P. Nguyen, S.W. Kim, H. Kim, H. Kim, K.H. Seok, M.J. Jung, A. Y-c, H.W. Kang, Biocompatible astaxanthin as a novel marine-oriented agent for dual chemophotothermal therapy, *PLoS One* 12 (4) (2017) e0174687.
- [67] P.R. Sowmya, B.P. Arathi, K. Vijay, V. Baskaran, R. Lakshminarayana, Astaxanthin from shrimp efficiently modulates oxidative stress and allied cell death progression in MCF-7 cells treated synergistically with beta-carotene and lutein from greens, *Food Chem. Toxicol.* 106 (Pt A) (2017) 58–69.
- [68] H. Zhao, H. Gu, H. Zhang, J.H. Li, W.E. Zhao, PPARgamma-dependent pathway in the growth-inhibitory effects of K562 cells by carotenoids in combination with rosiglitazone, *Biochim. Biophys. Acta* 1840 (1) (2014) 545–555.
- [69] J. Bechelli, M. Coppage, K. Rosell, J. Liesveld, Cytotoxicity of algae extracts on normal and malignant cells, *Leuk. Res. Treatment* 2011 (2011) 373519–373519.
- [70] H.P. Koeffler, Peroxisome proliferator-activated receptor gamma and cancers, *Clin. Cancer Res.* 9 (1) (2003) 1–9.
- [71] C.M. Lyon, D.M. Klinge, G.M.J. Do KC, C.L. Thomas, L.A. Damiani, T.H. March, C.A. Stidley, S.A. Belinsky, Rosiglitazone prevents the progression of preinvasive lung cancer in a murine model, *Carcinogenesis*. 30 (12) (2009) 2095–2099.
- [72] J.J. Liu, T. Hu, X.Y. Wu, C.Z. Wang, Y. Xu, Y. Zhang, R.Z. Xiao, D.J. Lin, R.W. Huang, Q. Liu, Peroxisome proliferator-activated receptor-gamma agonist rosiglitazone-induced apoptosis in leukemia k562 cells and its mechanisms of action, *Int. J. Toxicol.* 28 (2) (2009) 123–131.
- [73] A.L. Vine, J.S. Bertram, Upregulation of connexin 43 by retinoids but not by non-provitamin A carotenoids requires RARs, *Nutr. Cancer* 52 (1) (2005) 105–113.
- [74] T.J. King, L.H. Fukushima, A.D. Hieber, K.A. Shimabukuro, W.A. Sakr, J.S. Bertram, Reduced levels of connexin43 in cervical dysplasia: inducible expression in a cervical carcinoma cell line decreases neoplastic potential with implications for tumor progression, *Carcinogenesis*. 21 (6) (2000) 1097–1109.
- [75] C. Wu, J. Zhang, T. Liu, G. Jiao, C. Li, B. Hu, Astaxanthin inhibits proliferation and promotes apoptosis of A549 lung cancer cells via blocking JAK1/STAT3 pathway, *Cell. Mol. Immunol.* 32 (6) (2016) 784–788.
- [76] P. Palozza, C. Torelli, A. Boninsegna, R. Simone, A. Catalano, M.C. Mele, N. Picci, Growth-inhibitory effects of the astaxanthin-rich alga *Haematococcus pluvialis* in human colon cancer cells, *Cancer Lett.* 283 (1) (2009) 108–117.
- [77] R. Niranjana, R. Gayathri, S. Nimish Mol, T. Sugawara, T. Hirata, K. Miyashita, P. Ganesan, Carotenoids modulate the hallmarks of cancer cells, *J Funct Food*. 18 (2015) 968–985.
- [78] P.B. Atalay, G. Kuku, B.G. Tuna, Effects of carbendazim and astaxanthin co-treatment on the proliferation of MCF-7 breast cancer cells, *In Vitro Cell. Dev. Biol. Anim.* 55 (2) (2019) 113–119.
- [79] X. Qian, C. Tan, B. Yang, F. Wang, Y. Ge, Z. Guan, J. Cai, Astaxanthin increases radiosensitivity in esophageal squamous cell carcinoma through inducing apoptosis and G2/M arrest, *Dis. Esophagus* 30 (6) (2017) 1–7.
- [80] P.R. Sowmya, B.P. Arathi, K. Vijay, V. Baskaran, R. Lakshminarayana, Role of different vehicles in carotenoids delivery and their influence on cell viability, cell cycle progression, and induction of apoptosis in HeLa cells, *Mol. Cell. Biochem.* 406 (1–2) (2015) 245–253.
- [81] D. Tang, R. Kang, T.V. Berghe, P. Vandenabeele, G. Kroemer, The molecular machinery of regulated cell death, *Cell Res.* (2019) 1.
- [82] G.C. Cavalcante, A.P. Schaan, G.F. Cabral, M.N. Santana-da-Silva, P. Pinto, A.F. Vidal, Á. Ribeiro-dos-Santos, A cell's fate: an overview of the molecular biology and genetics of apoptosis, *Int. J. Mol. Sci.* 20 (17) (2019) 4133.
- [83] C.M. Pfeffer, Singh AT. Apoptosis: a target for anticancer therapy, *Int. J. Mol. Sci.* 19 (2) (2018) 448.
- [84] R.S. Wong, Apoptosis in cancer: from pathogenesis to treatment, *J. Exp. Clin. Cancer Res.* 30 (1) (2011) 87.
- [85] A. Ostuni, P. Lara, M.F. Armentano, R. Miglionico, A.M. Salvia, M. Monnich, M. Carosino, F.M. Lasorsa, M. Monne, I. Nilsson, F. Bisaccia, The hepatitis B x antigen anti-apoptotic effector URG7 is localized to the endoplasmic reticulum membrane, *FEBS Lett.* 587 (18) (2013) 3058–3062.
- [86] F.M. Millimouno, J. Dong, L. Yang, J. Li, X. Li, Targeting apoptosis pathways in cancer and perspectives with natural compounds from mother nature, *Cancer Prev. Res. Phila. (Phila)* 7 (11) (2014) 1081–1107.
- [87] K. von Schwarzenberg, A.M. Vollmar, Targeting apoptosis pathways by natural compounds in cancer: marine compounds as lead structures and chemical tools for cancer therapy, *Cancer Lett.* 332 (2) (2013) 295–303.
- [88] S. Chirumbolo, G. Björklund, R. Lysiuk, A. Vella, L. Lenchyk, T. Uppyr, Targeting cancer with phytochemicals via their fine tuning of the cell survival signaling pathways, *Int. J. Mol. Sci.* 19 (11) (2018) 3568.
- [89] A. Milani, M. Basirnejad, S. Shahbazi, A. Bolhassani, Carotenoids: biochemistry, pharmacology and treatment, *Br. J. Pharmacol.* 174 (11) (2017) 1290–1324.
- [90] P. Palozza, S. Serini, F. Di Nicuolo, G. Calviello, Modulation of apoptotic signalling by carotenoids in cancer cells, *Arch. Biochem. Biophys.* 430 (1) (2004) 104–109.
- [91] E. Kotake-Nara, M. Kushi, H. Zhang, T. Sugawara, K. Miyashita, A. Nagao, Carotenoids affect proliferation of human prostate cancer cells, *J. Nutr.* 131 (12) (2001) 3303–3306.
- [92] E. Kotake-Nara, M. Terasaki, A. Nagao, Characterization of apoptosis induced by fucoxanthin in human promyelocytic leukemia cells, *Biosci Biotech Biochem.* 69 (1) (2005) 224–227.
- [93] M. Hormozi, S. Ghoreishi, P. Baharvand, Astaxanthin induces apoptosis and increases activity of antioxidant enzymes in LS-180 cells, *Artif. Cells Nanomed. Biotechnol.* 47 (1) (2019) 891–895.
- [94] I.T.N. Teo, C.H. Chui, J.C.O. Tang, F.Y. Lau, G.Y.M. Cheng, R.S.M. Wong, S.H.L. Kok, C.H. Cheng, A.S.C. Chan, K.P. Ho, Antiproliferation and induction of cell death of *Phaffia rhodozyma* (*Xanthophyllomyces dendrorhous*) extract fermented by brewer malt waste on breast cancer cells, *Int. J. Mol. Med.* 16 (5) (2005) 931–936.
- [95] K. Vijay, S. PR-R, B.P. Arathi, S. Shilpa, H.J. Shwetha, M. Raju, V. Baskaran, R. Lakshminarayana, Low-dose doxorubicin with carotenoids selectively alters redox status and upregulates oxidative stress-mediated apoptosis in breast cancer cells, *Food Chem. Toxicol.* 118 (2018) 675–690.
- [96] C.M. Messina, S. Manuguerra, G. Renda, A. Santulli, Biotechnological applications for the sustainable use of marine by-products: *in vitro* antioxidant and pro-apoptotic effects of astaxanthin extracted with supercritical CO₂ from *parapeneus longirostris*, *Mar. Biotechnol.* (2019) 1–12.
- [97] X. Song, M. Wang, L. Zhang, J. Zhang, X. Wang, W. Liu, X. Gu, C. Lv, Changes in cell ultrastructure and inhibition of JAK1/STAT3 signaling pathway in CBRH-7919 cells with astaxanthin, *Toxicol. Mech. Methods* 22 (9) (2012) 679–686.
- [98] X.-Z. Su, R. Chen, C.-B. Wang, X.-L. Ouyang, Y. Jiang, M.-Y. Zhu, Astaxanthin combine with human serum albumin to abrogate cell proliferation, migration, and drug-resistant in human ovarian carcinoma SKOV3 cells, *Anticancer Agents Med. Chem.* 19 (6) (2019) 792–801.
- [99] K. Kavitha, J. Kowshik, T.K.K. Kishore, A.B. Baba, S. Nagini, Astaxanthin inhibits NF- κ B and Wnt/ β -catenin signaling pathways via inactivation of Erk/MAPK and PI3K/Akt to induce intrinsic apoptosis in a hamster model of oral cancer, *Biochim. Biophys. Acta* 1830 (10) (2013) 4433–4444.
- [100] E. Bousoik, H. Montazeri Aliabadi, “Do we know jack” about JAK? A closer look at JAK/STAT signaling pathway, *Front. Oncol.* 8 (2018) 287.
- [101] X. Wang, C.C. Decker, L. Zechner, S. Krstin, M. Wink, *In vitro* wound healing of tumor cells: inhibition of cell migration by selected cytotoxic alkaloids, *BMC Pharmacol. Toxicol.* 20 (1) (2019) 4.
- [102] L. Zhang, H. Wang, Multiple mechanisms of anti-cancer effects exerted by astaxanthin, *Mar. Drugs* 13 (7) (2015) 4310–4330.
- [103] J. Kowshik, A.B. Baba, H. Giri, G.D. Reddy, M. Dixit, S. Nagini, Astaxanthin inhibits JAK/STAT-3 signaling to abrogate cell proliferation, invasion and angiogenesis in a hamster model of oral cancer, *PLoS One* 9 (10) (2014) e109114.
- [104] T. Doi, T. Ishikawa, T. Okayama, K. Oka, K. Mizushima, T. Yasuda, N. Sakamoto, K. Katada, K. Kamada, K. Uchiyama, O. Handa, T. Takagi, Y. Naito, Y. Itoh, The JAK/STAT pathway is involved in the upregulation of PD-L1 expression in pancreatic cancer cell lines, *Oncol. Rep.* 37 (3) (2017) 1545–1554.
- [105] W. Ren, S. Shen, Z. Sun, P. Shu, X. Shen, C. Bu, F. Ai, X. Zhang, A. Tang, L. Tian, G. Li, X. Li, J. Ma, Jak-STAT3 pathway triggers DICER1 for proteasomal degradation by ubiquitin ligase complex of CUL4A(DCAF1) to promote colon cancer development, *Cancer Lett.* 375 (2) (2016) 209–220.
- [106] Z.P. Wang, H.F. Jin, R.D. Xu, Q.B. Mei, D.M. Fan, Triptolide downregulates Rac1

- and the JAK/STAT3 pathway and inhibits colitis-related colon cancer progression, *Exp. Mol. Med.* 41 (10) (2009) 717–727.
- [107] Y. Kozuki, Y. Miura, K. Yagasaki, Inhibitory effects of carotenoids on the invasion of rat ascites hepatoma cells in culture, *Cancer Lett.* 151 (1) (2000) 111–115.
- [108] H.Y. Kim, Y.M. Kim, S. Hong, Astaxanthin suppresses the metastasis of colon cancer by inhibiting the MYC-mediated downregulation of microRNA-29a-3p and microRNA-200a, *Sci. Rep.* 9 (1) (2019) 9457.
- [109] M. Shaimerdenova, O. Karapina, D. Mektapbayeva, K. Alibek, D. Akilbekova, The effects of antiviral treatment on breast cancer cell line, *Infect Agent Cancer.* 12 (2017) 18.
- [110] G. Donà, A. Andrisani, E. Tibaldi, A. Brunati, C. Sabbadin, D. Armanini, G. Ambrosini, E. Ragazzi, L. Bordin, Astaxanthin prevents human papillomavirus L1 protein binding in human sperm membranes, *Mar. Drugs* 16 (11) (2018) 427.
- [111] C.E. Peterson, R.L. Sedjo, F.G. Davis, C.A. Beam, A.R. Giuliano, Combined anti-oxidant carotenoids and the risk of persistent human papillomavirus infection, *Nutr. Cancer* 62 (6) (2010) 728–733.
- [112] S. Santoyo, L. Jaime, M. Plaza, M. Herrero, I. Rodriguez-Meizoso, E. Ibañez, G. Reglero, Antiviral compounds obtained from microalgae commonly used as carotenoid sources, *J. Appl. Phycol.* 24 (4) (2011) 731–741.
- [113] A.A. AlQahtani, O. A-MM, H.M. Al-Kreathy, S.E. Al-harthy, H.S. Al-malky, M.S.A. Nasser, F.O. Kamel, M.N. Alaama, Z.A. Damanhour, Chemosensitizing effects of marine astaxanthin on the anti-cancer activity of doxorubicin in tumor bearing mice, *Cancer Res.* 15 (1) (2019) 1–8.
- [114] S.E. El-Agamy, A.K. Abdel-Aziz, S. Wahdan, A. Esmat, S.S. Azab, Astaxanthin ameliorates doxorubicin-induced cognitive impairment (Chemobrain) in experimental rat model: impact on oxidative, inflammatory, and apoptotic machineries, *Mol. Neurobiol.* 55 (7) (2018) 5727–5740.
- [115] J.-C. Chen, C.-H. Wu, Y.-S. Peng, H.-Y. Zheng, Y.-C. Lin, P.-F. Ma, T.-C. Yen, T.-Y. Chen, Y.-W. Lin, Astaxanthin enhances erlotinib-induced cytotoxicity by p38 MAPK mediated xeroderma pigmentosum complementation group C (XPC) down-regulation in human lung cancer cells, *Toxicol. Res.* 7 (6) (2018) 1247–1256.
- [116] D. Tripathi, G. Jena, Astaxanthin inhibits cytotoxic and genotoxic effects of cyclophosphamide in mice germ cells, *Toxicology* 248 (2–3) (2008) 96–103.
- [117] S. Sreeja, C.K.K. Nair, Prevention of cisplatin-induced renal toxicity in swiss albino mice by astaxanthine, *Journal Pharmacol Toxicol.* 9 (1) (2014) 25–36.
- [118] G. Akca, H. Eren, L. Tumkaya, T. Mercantepe, M.O. Horsanali, E. Deveci, E. Dil, A. Yilmaz, The protective effect of astaxanthin against cisplatin-induced nephrotoxicity in rats, *Biomed. Pharmacother.* 100 (2018) 575–582.
- [119] X.-L. Xue, X.-D. Han, Y. Li, X.-F. Chu, W.-M. Miao, J.-L. Zhang, S.-J. Fan, Astaxanthin attenuates total body irradiation-induced hematopoietic system injury in mice via inhibition of oxidative stress and apoptosis, *Stem Cell Res. Ther.* 8 (1) (2017) 7.
- [120] T. Yan, H.Y. Li, J.S. Wu, Q. Niu, W.H. Duan, Q.Z. Han, W.M. Ji, T. Zhang, W. Lv, Astaxanthin inhibits gemcitabine-resistant human pancreatic cancer progression through EMT inhibition and gemcitabine resensitization, *Oncology Lett.* 14 (5) (2017) 5400–5408.
- [121] R. Peto, R. Doll, J.D. Buckley, M. Sporn, Can dietary beta-carotene materially reduce human cancer rates? *Nature.* 290 (5803) (1981) 201.
- [122] K. Uchiyama, Y. Naito, G. Hasegawa, N. Nakamura, J. Takahashi, T. Yoshikawa, Astaxanthin protects β -cells against glucose toxicity in diabetic db/db mice, *Redox Rep.* 7 (5) (2002) 290–293.
- [123] A. Sila, Z. Kamoun, Z. Ghilissi, M. Makni, M. Nasri, Z. Sahnoun, N. Nedjar-Arroume, A. Bougatef, Ability of natural astaxanthin from shrimp by-products to attenuate liver oxidative stress in diabetic rats, *Pharmacol. Rep.* 67 (2) (2015) 310–316.
- [124] Y. Naito, K. Uchiyama, W. Aoi, G. Hasegawa, N. Nakamura, N. Yoshida, T. Maoka, J. Takahashi, T. Yoshikawa, Prevention of diabetic nephropathy by treatment with astaxanthin in diabetic db/db mice, *Biofactors.* 20 (1) (2004) 49–59.
- [125] Q. Chen, J. Tao, G. Li, D. Zheng, Y. Tan, R. Li, L. Tian, Z. Li, H. Cheng, X. Xie, Astaxanthin ameliorates experimental diabetes-induced renal oxidative stress and fibronectin by upregulating connexin43 in glomerular mesangial cells and diabetic mice, *Eur. J. Pharmacol.* 840 (2018) 33–43.
- [126] X.-Y. Zhou, F. Zhang, H. X-t, J. Chen, T. R-x, Z. K-y, Y. Song, Depression can be prevented by astaxanthin through inhibition of hippocampal inflammation in diabetic mice, *Brain Res. Bull.* 1657 (2017) 262–268.
- [127] Y. Xue, Z. Qu, J. Fu, J. Zhen, W. Wang, Y. Cai, W. Wang, The protective effect of astaxanthin on learning and memory deficits and oxidative stress in a mouse model of repeated cerebral ischemia/reperfusion, *Brain Res. Bull.* 131 (2017) 221–228.
- [128] R. Nakao, O.L. Nelson, J.S. Park, B.D. Mathison, P.A. Thompson, B.P. Chew, Effect of astaxanthin supplementation on inflammation and cardiac function in BALB/c mice, *Anticancer Res.* 30 (7) (2010) 2721–2725.
- [129] G. Hussein, U. Sankawa, H. Goto, K. Matsumoto, H. Watanabe, Astaxanthin, a carotenoid with potential in human health and nutrition, *J. Nat. Prod.* 69 (3) (2006) 443–449.
- [130] Y.-H. Hwang, K.-J. Kim, S.-J. Kim, S.-K. Mun, S.-G. Hong, Y.-J. Son, S.-T. Yee, Suppression effect of astaxanthin on osteoclast formation in vitro and bone loss in vivo, *Int. J. Mol. Sci.* 19 (3) (2018) 912.
- [131] Y. Yoshihisa, T. Andoh, K. Matsunaga, M.U. Rehman, T. Maoka, T. Shimizu, Efficacy of astaxanthin for the treatment of atopic dermatitis in a murine model, *PLoS One* 11 (3) (2016) e0152288.
- [132] Y. Nakajima, Y. Inokuchi, M. Shimazawa, K. Otsubo, T. Ishibashi, H.H. Astaxanthin, A dietary carotenoid, protects retinal cells against oxidative stress in-vitro and in mice in-vivo, *J. Pharm. Pharmacol.* 60 (10) (2008) 1365–1374.
- [133] C.-H. Chiu, C.-C. Chang, S.-T. Lin, C.-C. Chyau, R.Y. Peng, Improved hepatoprotective effect of liposome-encapsulated astaxanthin in lipopolysaccharide-induced acute hepatotoxicity, *Int. J. Mol. Sci.* 17 (7) (2016) 1128.
- [134] T. Ohno, M. Shimizu, Y. Shirakami, T. Miyazaki, T. Ideta, T. Kochi, M. Kubota, H. Sakai, T. Tanaka, H. Moriwaki, Preventive effects of astaxanthin on diethylnitrosamine-induced liver tumorigenesis in C57/BL/KsJ-db/db obese mice, *Hepatol. Res.* 46 (3) (2016) E201–E209.
- [135] H. Kurihara, H. Koda, S. Asami, Y. Kiso, T. Tanaka, Contribution of the anti-oxidative property of astaxanthin to its protective effect on the promotion of cancer metastasis in mice treated with restraint stress, *Life Sci.* 70 (21) (2002) 2509–2520.
- [136] C.-I. Choi, Astaxanthin as a peroxisome proliferator-activated receptor (PPAR) modulator: its therapeutic implications, *Mar. Drugs* 17 (4) (2019) 242.
- [137] T. Tanaka, H. Makita, M. Ohnishi, H. Mori, K. Satoh, A. Hara, Chemoprevention of rat oral carcinogenesis by naturally occurring xanthophylls, astaxanthin and canthaxanthin, *Cancer Res.* 55 (18) (1995) 4059–4064.
- [138] K. Kavitha, P. Thiagarajan, J. Rathna, R. Mishra, S. Nagini, Chemopreventive effects of diverse dietary phytochemicals against DMBA-induced hamster buccal pouch carcinogenesis via the induction of Nrf2-mediated cytoprotective antioxidant, detoxification, and DNA repair enzymes, *Biochimie.* 95 (8) (2013) 1629–1639.
- [139] T. Maoka, H. Tokuda, N. Suzuki, H. Kato, H. Etoh, Anti-oxidative, anti-tumor-promoting, and anti-carcinogenic activities of nitroastaxanthin and nitrolutein, the reaction products of astaxanthin and lutein with peroxynitrite, *Mar. Drugs* 10 (6) (2012) 1391–1399.
- [140] A.R. Rao, H. Sindhuja, S.M. Dharmesh, K.U. Sankar, R. Sarada, G.A. Ravishankar, Effective inhibition of skin cancer, tyrosinase, and antioxidative properties by astaxanthin and astaxanthin esters from the green alga *Haematococcus pluvialis*, *J. Agr Food Chem.* 61 (16) (2013) 3842–3851.
- [141] T. Tanaka, Y. Morishita, M. Suzui, T. Kojima, A. Okumura, H. Mori, Chemoprevention of mouse urinary bladder carcinogenesis by the naturally occurring carotenoid astaxanthin, *Carcinogenesis.* 15 (1) (1994) 15–19.
- [142] X. Ni, H. Yu, S. Wang, C. Zhang, S. Shen, Astaxanthin inhibits PC-3 xenograft prostate tumor growth in nude mice, *Mar. Drugs* 15 (3) (2017) 66.
- [143] T. Yuri, K. Yoshizawa, Y. Emoto, Y. Kinoshita, M. Yuki, A. Tsubura, Effects of dietary xanthophylls, Canthaxanthin and astaxanthin on N-Methyl-N-nitrosourea-induced rat mammary carcinogenesis, *In Vivo (Brooklyn)* 30 (6) (2016) 795–800.
- [144] S. Safaverdi, F. Roshani, M. Rashti, S. Golkhoo, Z.M. Hassan, L. Langroudi, Micro-pix analysis in invasive ductal carcinoma tissues after treatment of astaxanthin, *Iran J Radiat Res.* 7 (2009) 33–39.
- [145] R. Nakao, O.L. Nelson, J.S. Park, B.D. Mathison, P.A. Thompson, B.P. Chew, Effect of dietary astaxanthin at different stages of mammary tumor initiation in BALB/c mice, *Anticancer Res.* 30 (6) (2010) 2171–2175.
- [146] T. Kochi, M. Shimizu, T. Sumi, M. Kubota, Y. Shirakami, T. Tanaka, H. Moriwaki, Inhibitory effects of astaxanthin on azoxymethane-induced colonic preneoplastic lesions in C57/BL/KsJ-db/db mice, *BMC gastroent* 14 (2014) 212.
- [147] P. Nagendraprabhu, G. Sudhandiran, Astaxanthin inhibits tumor invasion by decreasing extracellular matrix production and induces apoptosis in experimental rat colon carcinogenesis by modulating the expressions of ERK-2, NFkB and COX-2, *Invest. New Drugs* 29 (2) (2011) 207–224.
- [148] L. Cui, F. Xu, M. Wang, L. Li, T. Qiao, H. Cui, Z. Li, C. Sun, Dietary natural astaxanthin at an early stage inhibits N-nitrosomethylbenzylamine-induced esophageal cancer oxidative stress and inflammation via downregulation of NFkB and COX2 in F344 rats, *Onco. Ther.* 12 (2019) 5087–5096.
- [149] J.A. Edwards, P. Bellion, P. Beilstein, R. Rumbeli, J. Schierle, Review of genotoxicity and rat carcinogenicity investigations with astaxanthin, *Regul. Toxicol. Pharmacol.* 75 (2016) 5–19.
- [150] P. Jean Soon, C. Jong Hee, K. Yoo Kyung, L.L. Larry, P.C. Boon, Astaxanthin decreased oxidative stress and inflammation and enhanced immune response in humans, *Nutr. Metab. (Lond)* 5 (7) (2010) 18.